

William M Shafer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2061694/publications.pdf>

Version: 2024-02-01

127
papers

7,731
citations

53660

45
h-index

58464

82
g-index

173
all docs

173
docs citations

173
times ranked

5684
citing authors

#	ARTICLE	IF	CITATIONS
1	Gonococcal Clinical Strains Bearing a Common <i>gdhR</i> Single Nucleotide Polymorphism That Results in Enhanced Expression of the Virulence Gene <i>lctP</i> Frequently Possess a <i>mtrR</i> Promoter Mutation That Decreases Antibiotic Susceptibility. <i>MBio</i> , 2022, 13, e0027622.	1.8	4
2	A Single Amino Acid Substitution in Elongation Factor G Can Confer Low-Level Gentamicin Resistance in <i>Neisseria gonorrhoeae</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2022, 66, e0025122.	1.4	4
3	Structures of <i>Neisseria gonorrhoeae</i> MtrR-operator complexes reveal molecular mechanisms of DNA recognition and antibiotic resistance-conferring clinical mutations. <i>Nucleic Acids Research</i> , 2021, 49, 4155-4170.	6.5	13
4	A community-driven resource for genomic epidemiology and antimicrobial resistance prediction of <i>Neisseria gonorrhoeae</i> at Pathogenwatch. <i>Genome Medicine</i> , 2021, 13, 61.	3.6	63
5	Reply: Evidence of Recent Genomic Evolution in Gonococcal Strains With Decreased Susceptibility to Cephalosporins or Azithromycin in the United States, 2014–2016. <i>Journal of Infectious Diseases</i> , 2020, 221, 852-853.	1.9	2
6	Azithromycin susceptibility of <i>Neisseria gonorrhoeae</i> in the USA in 2017: a genomic analysis of surveillance data. <i>Lancet Microbe</i> , The, 2020, 1, e154-e164.	3.4	42
7	Drug Targeting a Gonococcal Virulence Factor Exploits Host Antimicrobial Peptides in Clearance of Infection. <i>Journal of Infectious Diseases</i> , 2020, 222, 1585-1586.	1.9	1
8	Transcriptional control of the gonococcal <i>ompA</i> gene by the MisR/MisS two-component regulatory system. <i>Scientific Reports</i> , 2020, 10, 9425.	1.6	2
9	Developing target product profiles for <i>Neisseria gonorrhoeae</i> diagnostics in the context of antimicrobial resistance: An expert consensus. <i>PLoS ONE</i> , 2020, 15, e0237424.	1.1	21
10	Cryo-EM Structures of a Gonococcal Multidrug Efflux Pump Illuminate a Mechanism of Drug Recognition and Resistance. <i>MBio</i> , 2020, 11, .	1.8	50
11	Genomic evolution of <i>Neisseria gonorrhoeae</i> since the preantibiotic era (1928–2013): antimicrobial use/misuse selects for resistance and drives evolution. <i>BMC Genomics</i> , 2020, 21, 116.	1.2	57
12	Identification of a <i>Neisseria gonorrhoeae</i> Histone Deacetylase: Epigenetic Impact on Host Gene Expression. <i>Pathogens</i> , 2020, 9, 132.	1.2	14
13	The serogroup B meningococcal outer membrane vesicle-based vaccine 4CMenB induces cross-species protection against <i>Neisseria gonorrhoeae</i> . <i>PLoS Pathogens</i> , 2020, 16, e1008602.	2.1	49
14	Title is missing!. , 2020, 16, e1008602.		0
15	Title is missing!. , 2020, 16, e1008602.		0
16	Title is missing!. , 2020, 16, e1008602.		0
17	Title is missing!. , 2020, 16, e1008602.		0
18	Structural, Biochemical, and <i>In Vivo</i> Characterization of MtrR-Mediated Resistance to Innate Antimicrobials by the Human Pathogen <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2019, 201, .	1.0	13

#	ARTICLE	IF	CITATIONS
19	Could Dampening Expression of the <i>Neisseria gonorrhoeae</i> <i>mtrCDE</i> -Encoded Efflux Pump Be a Strategy To Preserve Currently or Resurrect Formerly Used Antibiotics To Treat Gonorrhea?. <i>MBio</i> , 2019, 10, .	1.8	18
20	Evidence of Recent Genomic Evolution in Gonococcal Strains With Decreased Susceptibility to Cephalosporins or Azithromycin in the United States, 2014–2016. <i>Journal of Infectious Diseases</i> , 2019, 220, 294-305.	1.9	38
21	Transcriptional regulation of a gonococcal gene encoding a virulence factor (L-lactate permease). <i>PLoS Pathogens</i> , 2019, 15, e1008233.	2.1	12
22	Copper Ions and Coordination Complexes as Novel Carbapenem Adjuvants. <i>Antimicrobial Agents and Chemotherapy</i> , 2018, 62, .	1.4	31
23	Mechanistic Basis for Decreased Antimicrobial Susceptibility in a Clinical Isolate of <i>Neisseria gonorrhoeae</i> Possessing a Mosaic-Like <i>mtr</i> Efflux Pump Locus. <i>MBio</i> , 2018, 9, .	1.8	70
24	Mosaic Drug Efflux Gene Sequences from Commensal <i>Neisseria</i> Can Lead to Low-Level Azithromycin Resistance Expressed by <i>Neisseria gonorrhoeae</i> Clinical Isolates. <i>MBio</i> , 2018, 9, .	1.8	19
25	Structure-Function Relationships of the Neisserial EptA Enzyme Responsible for Phosphoethanolamine Decoration of Lipid A: Rationale for Drug Targeting. <i>Frontiers in Microbiology</i> , 2018, 9, 1922.	1.5	16
26	<i>cis</i> - and <i>trans</i> -Acting Factors Influence Expression of the <i>norM</i> -Encoded Efflux Pump of <i>Neisseria gonorrhoeae</i> and Levels of Gonococcal Susceptibility to Substrate Antimicrobials. <i>Antimicrobial Agents and Chemotherapy</i> , 2018, 62, .	1.4	4
27	Whole-Genome Sequencing of a Large Panel of Contemporary <i>Neisseria gonorrhoeae</i> Clinical Isolates Indicates that a Wild-Type <i>mtrA</i> Gene Is Common: Implications for Inducible Antimicrobial Resistance. <i>Antimicrobial Agents and Chemotherapy</i> , 2017, 61, .	1.4	7
28	Control of <i>gdhR</i> Expression in <i>Neisseria gonorrhoeae</i> via Autoregulation and a Master Repressor (<i>MtrR</i>) of a Drug Efflux Pump Operon. <i>MBio</i> , 2017, 8, .	1.8	14
29	<i>Neisseria gonorrhoeae</i> : Drug Resistance, Mouse Models, and Vaccine Development. <i>Annual Review of Microbiology</i> , 2017, 71, 665-686.	2.9	166
30	Multidrug-resistant gonorrhea: A research and development roadmap to discover new medicines. <i>PLoS Medicine</i> , 2017, 14, e1002366.	3.9	129
31	Antimicrobial Resistance Expressed by <i>Neisseria gonorrhoeae</i> : A Major Global Public Health Problem in the 21st Century. , 2016, , 213-237.		5
32	The <i>MisR</i> Response Regulator Is Necessary for Intrinsic Cationic Antimicrobial Peptide and Aminoglycoside Resistance in <i>Neisseria gonorrhoeae</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 4690-4700.	1.4	21
33	Efflux Pumps in <i>Neisseria gonorrhoeae</i> : Contributions to Antimicrobial Resistance and Virulence. , 2016, , 439-469.		10
34	Does the Cervicovaginal Microbiome Facilitate Transmission of <i>Neisseria gonorrhoeae</i> From Women to Men? Implications for Understanding Transmission of Gonorrhea and Advancing Vaccine Development. <i>Journal of Infectious Diseases</i> , 2016, 214, 1615-1617.	1.9	4
35	The genes that encode the gonococcal transferrin binding proteins, <i>TbpB</i> and <i>TbpA</i> , are differentially regulated by <i>MisR</i> under iron-replete and iron-depleted conditions. <i>Molecular Microbiology</i> , 2016, 102, 137-151.	1.2	5
36	Antimicrobial Resistance Expressed by <i>Neisseria gonorrhoeae</i> : A Major Global Public Health Problem in the 21st Century. <i>Microbiology Spectrum</i> , 2016, 4, .	1.2	178

#	ARTICLE	IF	CITATIONS
37	The Transcriptional Repressor, MtrR, of the mtrCDE Efflux Pump Operon of <i>Neisseria gonorrhoeae</i> Can Also Serve as an Activator of β -Lactamase Gene (<i>glnE</i>) Expression. <i>Antibiotics</i> , 2015, 4, 188-197.	1.5	11
38	Genetic Resistance Determinants, In Vitro Time-Kill Curve Analysis and Pharmacodynamic Functions for the Novel Topoisomerase II Inhibitor ETX0914 (AZD0914) in <i>Neisseria gonorrhoeae</i> . <i>Frontiers in Microbiology</i> , 2015, 6, 1377.	1.5	44
39	Population structure of <i>Neisseria gonorrhoeae</i> based on whole genome data and its relationship with antibiotic resistance. <i>PeerJ</i> , 2015, 3, e806.	0.9	67
40	Copper(II)-Bis(Thiosemicarbazonato) Complexes as Antibacterial Agents: Insights into Their Mode of Action and Potential as Therapeutics. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 6444-6453.	1.4	59
41	Future treatment of gonorrhoea – novel emerging drugs are essential and in progress?. <i>Expert Opinion on Emerging Drugs</i> , 2015, 20, 357-360.	1.0	13
42	On the in vivo significance of bacterial resistance to antimicrobial peptides. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2015, 1848, 3101-3111.	1.4	42
43	Structure and Function of <i>Neisseria gonorrhoeae</i> MtrF Illuminates a Class of Antimetabolite Efflux Pumps. <i>Cell Reports</i> , 2015, 11, 61-70.	2.9	44
44	Characterization of a spermine/spermidine transport system reveals a novel DNA sequence duplication in <i>Neisseria gonorrhoeae</i> . <i>FEMS Microbiology Letters</i> , 2015, 362, fmv125.	0.7	7
45	Overproduction of the MtrCDE Efflux Pump in <i>Neisseria gonorrhoeae</i> Produces Unexpected Changes in Cellular Transcription Patterns. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 724-726.	1.4	13
46	Phosphoethanolamine Modification of <i>Neisseria gonorrhoeae</i> Lipid A Reduces Autophagy Flux in Macrophages. <i>PLoS ONE</i> , 2015, 10, e0144347.	1.1	22
47	<i>Neisseria gonorrhoeae</i> Modulates Iron-Limiting Innate Immune Defenses in Macrophages. <i>PLoS ONE</i> , 2014, 9, e87688.	1.1	52
48	Crystal Structure of the Open State of the <i>Neisseria gonorrhoeae</i> MtrE Outer Membrane Channel. <i>PLoS ONE</i> , 2014, 9, e97475.	1.1	51
49	Crystal Structure of the <i>Neisseria gonorrhoeae</i> MtrD Inner Membrane Multidrug Efflux Pump. <i>PLoS ONE</i> , 2014, 9, e97903.	1.1	65
50	Phosphoethanolamine Decoration of <i>Neisseria gonorrhoeae</i> Lipid A Plays a Dual Immunostimulatory and Protective Role during Experimental Genital Tract Infection. <i>Infection and Immunity</i> , 2014, 82, 2170-2179.	1.0	38
51	Phase-Variable Expression of <i>ptA</i> Modulates the Resistance of <i>Neisseria gonorrhoeae</i> to Cationic Antimicrobial Peptides. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 4230-4233.	1.4	21
52	Importance of Multidrug Efflux Pumps in the Antimicrobial Resistance Property of Clinical Multidrug-Resistant Isolates of <i>Neisseria gonorrhoeae</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 3556-3559.	1.4	96
53	Antimicrobial Resistance in <i>Neisseria gonorrhoeae</i> in the 21st Century: Past, Evolution, and Future. <i>Clinical Microbiology Reviews</i> , 2014, 27, 587-613.	5.7	894
54	Identification of Regulatory Elements That Control Expression of the <i>tbpA</i> Operon in <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2014, 196, 2762-2774.	1.0	15

#	ARTICLE	IF	CITATIONS
55	Challenges with gonorrhea in the era of multi-drug and extensively drug resistance “are we on the right track?. Expert Review of Anti-Infective Therapy, 2014, 12, 653-656.	2.0	30
56	Spermine impairs biofilm formation by <i>Neisseria gonorrhoeae</i> . FEMS Microbiology Letters, 2013, 343, 64-69.	0.7	36
57	Mechanisms and Significance of Bacterial Resistance to Human Cationic Antimicrobial Peptides. , 2013, , 219-254.		5
58	Lipid A’s Structure Mediates <i>Neisseria gonorrhoeae</i> Fitness during Experimental Infection of Mice and Men. MBio, 2013, 4, e00892-13.	1.8	56
59	Phosphoethanolamine Residues on the Lipid A Moiety of <i>Neisseria gonorrhoeae</i> Lipooligosaccharide Modulate Binding of Complement Inhibitors and Resistance to Complement Killing. Infection and Immunity, 2013, 81, 33-42.	1.0	46
60	MtrR Control of a Transcriptional Regulatory Pathway in <i>Neisseria meningitidis</i> That Influences Expression of a Gene (<i>nadA</i>) Encoding a Vaccine Candidate. PLoS ONE, 2013, 8, e56097.	1.1	7
61	Dueling Regulatory Properties of a Transcriptional Activator (MtrA) and Repressor (MtrR) That Control Efflux Pump Gene Expression in <i>Neisseria gonorrhoeae</i> . MBio, 2012, 3, e00446-12.	1.8	22
62	MpeR Regulates the <i>mtr</i> Efflux Locus in <i>Neisseria gonorrhoeae</i> and Modulates Antimicrobial Resistance by an Iron-Responsive Mechanism. Antimicrobial Agents and Chemotherapy, 2012, 56, 1491-1501.	1.4	28
63	Impact of Fluoroquinolone Resistance Mutations on Gonococcal Fitness and In Vivo Selection for Compensatory Mutations. Journal of Infectious Diseases, 2012, 205, 1821-1829.	1.9	73
64	Lipooligosaccharide Structure is an Important Determinant in the Resistance of <i>Neisseria Gonorrhoeae</i> to Antimicrobial Agents of Innate Host Defense. Frontiers in Microbiology, 2011, 2, 30.	1.5	26
65	Experimental Gonococcal Infection in Male Volunteers: Cumulative Experience with <i>Neisseria gonorrhoeae</i> Strains FA1090 and MS11mkC. Frontiers in Microbiology, 2011, 2, 123.	1.5	102
66	The evolution of infectious agents in relation to sex in animals and humans: brief discussions of some individual organisms. Annals of the New York Academy of Sciences, 2011, 1230, 74-107.	1.8	5
67	Antibiotic resistance in <i>Neisseria gonorrhoeae</i> : origin, evolution, and lessons learned for the future. Annals of the New York Academy of Sciences, 2011, 1230, E19-28.	1.8	174
68	Taking the Gonococcus-Human Relationship to a Whole New Level: Implications for the Coevolution of Microbes and Humans. MBio, 2011, 2, e00067-11.	1.8	2
69	A Novel Mechanism of High-Level, Broad-Spectrum Antibiotic Resistance Caused by a Single Base Pair Change in <i>Neisseria gonorrhoeae</i> . MBio, 2011, 2, .	1.8	77
70	Efflux Pumps of the Resistance“Nodulation”Division Family: A Perspective of their Structure, Function, and Regulation in Gram-Negative Bacteria. Advances in Enzymology and Related Areas of Molecular Biology, 2011, 77, 109-146.	1.3	42
71	Off-Target Gene Regulation Mediated by Transcriptional Repressors of Antimicrobial Efflux Pump Genes in <i>Neisseria gonorrhoeae</i> . Antimicrobial Agents and Chemotherapy, 2011, 55, 2559-2565.	1.4	16
72	The Iron-Repressed, AraC-Like Regulator MpeR Activates Expression of <i>fetA</i> in <i>Neisseria gonorrhoeae</i> . Infection and Immunity, 2011, 79, 4764-4776.	1.0	32

#	ARTICLE	IF	CITATIONS
73	The Human Host Defense Peptide LL-37 Interacts with <i>Neisseria meningitidis</i> Capsular Polysaccharides and Inhibits Inflammatory Mediators Release. <i>PLoS ONE</i> , 2010, 5, e13627.	1.1	28
74	Human Antimicrobial Peptide LL-37 Induces MefE/Mel-Mediated Macrolide Resistance in <i>Streptococcus pneumoniae</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2010, 54, 3516-3519.	1.4	45
75	Altered Growth, Pigmentation, and Antimicrobial Susceptibility Properties of <i>Staphylococcus aureus</i> Due to Loss of the Major Cold Shock Gene <i>cspB</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2010, 54, 2283-2290.	1.4	38
76	Biologic Activities of the TolC-Like Protein of <i>Neisseria meningitidis</i> as Assessed by Functional Complementation in <i>Escherichia coli</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2010, 54, 506-508.	1.4	5
77	Polyamines Can Increase Resistance of <i>Neisseria gonorrhoeae</i> to Mediators of the Innate Human Host Defense. <i>Infection and Immunity</i> , 2010, 78, 3187-3195.	1.0	44
78	Phosphoethanolamine Substitution of Lipid A and Resistance of <i>Neisseria gonorrhoeae</i> to Cationic Antimicrobial Peptides and Complement-Mediated Killing by Normal Human Serum. <i>Infection and Immunity</i> , 2009, 77, 1112-1120.	1.0	102
79	MtrR Modulates <i>rpoH</i> Expression and Levels of Antimicrobial Resistance in <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2009, 191, 287-297.	1.0	58
80	Crystallization and preliminary X-ray diffraction analysis of the multidrug efflux transporter NorM from <i>Neisseria gonorrhoeae</i> . <i>Acta Crystallographica Section F: Structural Biology Communications</i> , 2008, 64, 289-292.	0.7	5
81	Clinically relevant mutations that cause derepression of the <i>Neisseria gonorrhoeae</i> MtrC-MtrD-MtrE Efflux pump system confer different levels of antimicrobial resistance and <i>in vivo</i> fitness. <i>Molecular Microbiology</i> , 2008, 70, 462-478.	1.2	185
82	Phenotypic and Genotypic Analyses of <i>Neisseria gonorrhoeae</i> Isolates That Express Frequently Recovered PorB PIA Variable Region Types Suggest that Certain P1a Porin Sequences Confer a Selective Advantage for Urogenital Tract Infection. <i>Infection and Immunity</i> , 2008, 76, 3700-3709.	1.0	24
83	Functional Cloning and Characterization of the Multidrug Efflux Pumps NorM from <i>Neisseria gonorrhoeae</i> and YdhE from <i>Escherichia coli</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2008, 52, 3052-3060.	1.4	76
84	Expression of the MtrC-MtrD-MtrE Efflux Pump in <i>Neisseria gonorrhoeae</i> and Bacterial Survival in the Presence of Antimicrobials. , 2008, , 55-63.		3
85	Two ABC Transporter Operons and the Antimicrobial Resistance Gene <i>mtrF</i> Are <i>pilT</i> Responsive in <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2007, 189, 5399-5402.	1.0	11
86	The TolC-Like Protein of <i>Neisseria meningitidis</i> Is Required for Extracellular Production of the Repeats-in-Toxin Toxin FrpC but Not for Resistance to Antimicrobials Recognized by the Mtr Efflux Pump System. <i>Infection and Immunity</i> , 2007, 75, 6008-6012.	1.0	10
87	Differential Regulation of <i>ponA</i> and <i>pilMNOPQ</i> Expression by the MtrR Transcriptional Regulatory Protein in <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2007, 189, 4569-4577.	1.0	31
88	Regulation of the MtrC-MtrD-MtrE Efflux Pump System Modulates the <i>In Vivo</i> Fitness of <i>Neisseria gonorrhoeae</i> . <i>Journal of Infectious Diseases</i> , 2007, 196, 1804-1812.	1.9	116
89	Integration Host Factor is required for FarR repression of the <i>farAB</i> -encoded efflux pump of <i>Neisseria gonorrhoeae</i> . <i>Molecular Microbiology</i> , 2006, 60, 1381-1400.	1.2	16
90	Towards an Understanding of Chromosomally Mediated Penicillin Resistance in <i>Neisseria gonorrhoeae</i> : Evidence for a Porin-Efflux Pump Collaboration. <i>Journal of Bacteriology</i> , 2006, 188, 2297-2299.	1.0	29

#	ARTICLE	IF	CITATIONS
91	Antimicrobial peptides and endotoxin inhibit cytokine and nitric oxide release but amplify respiratory burst response in human and murine macrophages. <i>Cellular Microbiology</i> , 2005, 7, 1251-1262.	1.1	111
92	Resistance of <i>Neisseria meningitidis</i> to the Toxic Effects of Heme Iron and Other Hydrophobic Agents Requires Expression of ght. <i>Journal of Bacteriology</i> , 2005, 187, 5214-5223.	1.0	11
93	Characterization of the Multiple Transferable Resistance Repressor, MtrR, from <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2005, 187, 5008-5012.	1.0	43
94	Regulation of mtrF Expression in <i>Neisseria gonorrhoeae</i> and Its Role in High-Level Antimicrobial Resistance. <i>Journal of Bacteriology</i> , 2005, 187, 3713-3720.	1.0	49
95	CspA Regulates Pigment Production in <i>Staphylococcus aureus</i> through a SigB-Dependent Mechanism. <i>Journal of Bacteriology</i> , 2005, 187, 8181-8184.	1.0	49
96	Characterization of the MacA-MacB efflux system in <i>Neisseria gonorrhoeae</i> . <i>Journal of Antimicrobial Chemotherapy</i> , 2005, 56, 856-860.	1.3	100
97	Cationic Antimicrobial Peptide Resistance in <i>Neisseria meningitidis</i> . <i>Journal of Bacteriology</i> , 2005, 187, 5387-5396.	1.0	209
98	A Mutant Form of the <i>Neisseria gonorrhoeae</i> Pilus Secretin Protein PilQ Allows Increased Entry of Heme and Antimicrobial Compounds. <i>Journal of Bacteriology</i> , 2004, 186, 730-739.	1.0	49
99	The Down-Regulation of Cathepsin G in THP-1 Monocytes after Infection with <i>Mycobacterium tuberculosis</i> Is Associated with Increased Intracellular Survival of Bacilli. <i>Infection and Immunity</i> , 2004, 72, 5712-5721.	1.0	51
100	Modulation of the mtrCDE-encoded efflux pump gene complex of <i>Neisseria meningitidis</i> due to a <i>Correa</i> element insertion sequence. <i>Molecular Microbiology</i> , 2004, 54, 731-741.	1.2	68
101	Degradation of Human Antimicrobial Peptide LL-37 by <i>Staphylococcus aureus</i> -Derived Proteinases. <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 4673-4679.	1.4	454
102	Divergence and transcriptional analysis of the division cell wall (dcw) gene cluster in <i>Neisseria</i> spp.. <i>Molecular Microbiology</i> , 2003, 47, 431-442.	1.2	35
103	The increased bactericidal activity of a fatty acid-modified synthetic antimicrobial peptide of human cathepsin G correlates with its enhanced capacity to interact with model membranes. <i>International Journal of Antimicrobial Agents</i> , 2003, 21, 13-19.	1.1	54
104	Susceptibility of <i>Treponema pallidum</i> to host-derived antimicrobial peptides. <i>Peptides</i> , 2003, 24, 1741-1746.	1.2	29
105	Identification of a cell envelope protein (MtrF) involved in hydrophobic antimicrobial resistance in <i>Neisseria gonorrhoeae</i> . <i>Journal of Antimicrobial Chemotherapy</i> , 2003, 51, 27-37.	1.3	47
106	The Major Cold Shock Gene, cspA, Is Involved in the Susceptibility of <i>Staphylococcus aureus</i> to an Antimicrobial Peptide of Human Cathepsin G. <i>Infection and Immunity</i> , 2003, 71, 4304-4312.	1.0	55
107	A Gonococcal Efflux Pump System Enhances Bacterial Survival in a Female Mouse Model of Genital Tract Infection. <i>Infection and Immunity</i> , 2003, 71, 5576-5582.	1.0	186
108	The NorM Efflux Pump of <i>Neisseria gonorrhoeae</i> and <i>Neisseria meningitidis</i> Recognizes Antimicrobial Cationic Compounds. <i>Journal of Bacteriology</i> , 2003, 185, 1101-1106.	1.0	111

#	ARTICLE	IF	CITATIONS
109	Overexpression of the MtrC-MtrD-MtrE Efflux Pump Due to an mtrR Mutation Is Required for Chromosomally Mediated Penicillin Resistance in <i>Neisseria gonorrhoeae</i> . <i>Journal of Bacteriology</i> , 2002, 184, 5619-5624.	1.0	166
110	Inducible, but Not Constitutive, Resistance of Gonococci to Hydrophobic Agents Due to the MtrC-MtrD-MtrE Efflux Pump Requires TonB-ExbB-ExbD Proteins. <i>Antimicrobial Agents and Chemotherapy</i> , 2002, 46, 561-565.	1.4	27
111	Tailoring an Antibacterial Peptide of Human Lysosomal Cathepsin G to Enhance its Broad-Spectrum Action Against Antibiotic-Resistant Bacterial Pathogens. <i>Current Pharmaceutical Design</i> , 2002, 8, 695-702.	0.9	42
112	Phase variable changes in genes lgtA and lgtC within the lgtABCDE operon of <i>Neisseria gonorrhoeae</i> can modulate gonococcal susceptibility to normal human serum. <i>Journal of Endotoxin Research</i> , 2002, 8, 47-58.	2.5	25
113	A Putatively Phase Variable Gene (dca) Required for Natural Competence in <i>Neisseria gonorrhoeae</i> but Not <i>Neisseria meningitidis</i> Is Located within the Division Cell Wall (dcw) Gene Cluster. <i>Journal of Bacteriology</i> , 2001, 183, 1233-1241.	1.0	27
114	Genetic organization and regulation of antimicrobial efflux systems possessed by <i>Neisseria gonorrhoeae</i> and <i>Neisseria meningitidis</i> . <i>Journal of Molecular Microbiology and Biotechnology</i> , 2001, 3, 219-24.	1.0	35
115	Decreased Azithromycin Susceptibility of <i>Neisseria gonorrhoeae</i> Due to mtrR Mutations. <i>Antimicrobial Agents and Chemotherapy</i> , 1999, 43, 2468-2472.	1.4	145
116	Induction of the mtrCDE-encoded efflux pump system of <i>Neisseria gonorrhoeae</i> requires MtrA, an AraC-like protein. <i>Molecular Microbiology</i> , 1999, 33, 651-658.	1.2	93
117	The farAB-encoded efflux pump mediates resistance of gonococci to long-chained antibacterial fatty acids. <i>Molecular Microbiology</i> , 1999, 33, 839-845.	1.2	148
118	Use of Cefazolin Microspheres to Treat Localized Methicillin-Resistant <i>Staphylococcus aureus</i> Infections in Rats. <i>Journal of Surgical Research</i> , 1999, 86, 97-102.	0.8	26
119	Loss-of-function mutations in the mtr efflux system of <i>Neisseria gonorrhoeae</i> . <i>Microbiology (United Kingdom)</i> , 1997, 143, 2117-2125.	0.7	103
120	Modulation of <i>Neisseria gonorrhoeae</i> susceptibility to vertebrate antibacterial peptides due to a member of the resistance/nodulation/division efflux pump family. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 1829-1833.	3.3	353
121	The MtrD protein of <i>Neisseria gonorrhoeae</i> is a member of the resistance/nodulation/division protein family constituting part of an efflux system. <i>Microbiology (United Kingdom)</i> , 1997, 143, 2117-2125.	0.7	103
122	Generation of antiserum to specific epitopes. <i>Molecular Biotechnology</i> , 1996, 6, 231-240.	1.3	0
123	Membrane glycerophospholipid biosynthesis in <i>Neisseria meningitidis</i> and <i>Neisseria gonorrhoeae</i> : identification, characterization, and mutagenesis of a lysophosphatidic acid acyltransferase. <i>Molecular Microbiology</i> , 1995, 18, 401-412.	1.2	24
124	Importance of lipooligosaccharide structure in determining gonococcal resistance to hydrophobic antimicrobial agents resulting from the mtr efflux system. <i>Molecular Microbiology</i> , 1995, 16, 1001-1009.	1.2	47
125	Resistance of <i>Neisseria gonorrhoeae</i> to antimicrobial hydrophobic agents is modulated by the mtrRCDE efflux system. <i>Microbiology (United Kingdom)</i> , 1995, 141, 611-622.	0.7	355
126	Cleavage of gonococcal pilin and alteration of common pilin epitopes by human neutrophil lysosomal elastase. <i>FEMS Microbiology Letters</i> , 1988, 52, 47-51.	0.7	3

#	ARTICLE	IF	CITATIONS
127	Neutrophil Killing of Bacteria by Oxygen-Independent Mechanisms: A Historical Summary. <i>Clinical Infectious Diseases</i> , 1985, 7, 398-403.	2.9	71