Beatriz Vazquez

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/2062238/beatriz-vazquez-publications-by-year.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

80 2,043 41 27 h-index g-index citations papers 82 2,320 4.5 4.51 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
80	Antimicrobial Activity of Five Apitoxins from on Two Common Foodborne Pathogens. <i>Antibiotics</i> , 2020 , 9,	4.9	2
79	Antimicrobial activity of apitoxin from Apis mellifera in Salmonella enterica strains isolated from poultry and its effects on motility, biofilm formation and gene expression. <i>Microbial Pathogenesis</i> , 2019 , 137, 103771	3.8	6
78	Determination of Florfenicol, Thiamfenicol and Chloramfenicol at Trace Levels in Animal Feed by HPLC?MS/MS. <i>Antibiotics</i> , 2019 , 8,	4.9	7
77	Consumption of pharmaceuticals by dairy cows via watering throught: Uncontroled intake. <i>Agriculture, Ecosystems and Environment</i> , 2019 , 280, 95-101	5.7	5
76	Tracing recombinant bovine somatotropin ab(use) through transcriptomics: the potential of bovine somatic cells in a multi-dose longitudinal study. <i>Scientific Reports</i> , 2019 , 9, 4788	4.9	6
75	Transcriptomics: A powerful tool to evaluate the behavior of foodborne pathogens in the food production chain. <i>Food Research International</i> , 2019 , 125, 108543	7	18
74	Short Chain Fatty Acids Commonly Produced by Gut Microbiota Influence Motility, Biofilm Formation, and Gene Expression. <i>Antibiotics</i> , 2019 , 8,	4.9	23
73	Influence of milk, chicken residues and oxygen levels on biofilm formation on stainless steel, gene expression and small RNAs in Salmonella enterica. <i>Food Control</i> , 2018 , 90, 1-9	6.2	17
7 ²	Constant iodine intake through the diet could improve hypothyroidism treatment: a case report. Journal of Physiology and Biochemistry, 2018 , 74, 189-193	5	
71	Salmonella and Campylobacter biofilm formation: a comparative assessment from farm to fork. Journal of the Science of Food and Agriculture, 2018, 98, 4014-4032	4.3	43
70	A comprehensive review of non-enterica subspecies of Salmonella enterica. <i>Microbiological Research</i> , 2018 , 206, 60-73	5.3	93
69	Tracing Recombinant Bovine Somatotropin Ab(Use) Through Gene Expression in Blood, Hair Follicles, and Milk Somatic Cells: A Matrix Comparison. <i>Molecules</i> , 2018 , 23,	4.8	2
68	Food preservatives influence biofilm formation, gene expression and small RNAs in Salmonella enterica. LWT - Food Science and Technology, 2018, 97, 1-8	5.4	9
67	Confirmation of Eagonist residues in bovine retina and liver using HPLC-MS/MS and evaluation of matrix-dependent problems. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment,</i> 2017 , 34, 598-607	3.2	3
66	Identification and quantification of 12 pharmaceuticals in water collected from milking parlors: Food safety implications. <i>Journal of Dairy Science</i> , 2017 , 100, 3373-3383	4	4
65	Phytanic acid consumption and human health, risks, benefits and future trends: A review. <i>Food Chemistry</i> , 2017 , 221, 237-247	8.5	28
64	Effects of bovine pregnancy on the fatty acid composition of milk: the significance for humans needs. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2017 , 34, 608-616	3.2	3

(2010-2017)

63	Effect of Food Residues in Biofilm Formation on Stainless Steel and Polystyrene Surfaces by Salmonella enterica Strains Isolated from Poultry Houses. <i>Foods</i> , 2017 , 6,	4.9	13	
62	Biofilm Formation and Morphotypes of Salmonella enterica subsp.arizonae Differs from Those of Other Salmonella enterica Subspecies in Isolates from Poultry Houses. <i>Journal of Food Protection</i> , 2016 , 79, 1127-34	2.5	18	
61	A Confirmatory Method Based on HPLC-MS/MS for the Detection and Quantification of Residue of Tetracyclines in Nonmedicated Feed. <i>Journal of Analytical Methods in Chemistry</i> , 2016 , 2016, 1202954	2	17	
60	An Evaluation of Alternatives to Nitrites and Sulfites to Inhibit the Growth of Salmonella enterica and Listeria monocytogenes in Meat Products. <i>Foods</i> , 2016 , 5,	4.9	19	
59	Prevalence, molecular characterization and antimicrobial resistance of Salmonella serovars isolated from northwestern Spanish broiler flocks (2011-2015). <i>Poultry Science</i> , 2016 , 95, 2097-105	3.9	32	
58	Biofilm formation, phenotypic production of cellulose and gene expression in Salmonella enterica decrease under anaerobic conditions. <i>International Journal of Food Microbiology</i> , 2016 , 238, 63-67	5.8	25	
57	Assessment of Tetracyclines Residues and Tetracycline Resistant Bacteria in Conventional and Organic Baby Foods. <i>Foods</i> , 2015 , 4, 306-317	4.9	4	
56	Detection of veterinary drug residues in surface waters collected nearby farming areas in Galicia, North of Spain. <i>Environmental Science and Pollution Research</i> , 2014 , 21, 2367-2377	5.1	40	
55	Quantitative detection of tetracycline-resistant microorganisms in conventional and organic beef, pork and chicken meat. <i>CYTA - Journal of Food</i> , 2014 , 12, 383-388	2.3	6	
54	Monitoring the presence of 13 active compounds in surface water collected from rural areas in northwestern Spain. <i>International Journal of Environmental Research and Public Health</i> , 2014 , 11, 5251-	72 ^{4.6}	21	
53	A simple and rapid method for the identification and quantification of malachite green and its metabolite in hake by HPLCMS/MS. <i>Food Control</i> , 2013 , 31, 102-107	6.2	40	
52	Determination of the Presence of Three Antimicrobials in Surface Water Collected from Urban and Rural Areas. <i>Antibiotics</i> , 2013 , 2, 46-57	4.9	12	
51	Direct quantification and distribution of tetracycline-resistant genes in meat samples by real-time polymerase chain reaction. <i>Journal of Food Science</i> , 2012 , 77, M372-6	3.4	5	
50	Development of a Hamburger Patty with Healthier Lipid Formulation and Study of its Nutritional, Sensory, and Stability Properties. <i>Food and Bioprocess Technology</i> , 2012 , 5, 200-208	5.1	43	
49	Detection and quantitative analysis of 21 veterinary drugs in river water using high-pressure liquid chromatography coupled to tandem mass spectrometry. <i>Environmental Science and Pollution Research</i> , 2012 , 19, 3235-49	5.1	61	
48	Real-time polymerase chain reaction for the quantitative detection of tetA and tetB bacterial tetracycline resistance genes in food. <i>International Journal of Food Microbiology</i> , 2011 , 146, 284-9	5.8	32	
47	The effects of industrial pre-frying and domestic cooking methods on the nutritional compositions and fatty acid profiles of two different frozen breaded foods. <i>LWT - Food Science and Technology</i> , 2010 , 43, 1271-1276	5.4	23	
46	Determination of the hormonal growth promoter 17alpha-methyltestosterone in food-producing animals: bovine hair analysis by HPLC-MS/MS. <i>Meat Science</i> , 2010 , 84, 196-201	6.4	17	

45	Sensitive Determination of Decoquinate in Milk by High-Performance Liquid Chromatographic Coupled to Laser-Induced Fluorescence Detection. <i>Food Analytical Methods</i> , 2010 , 3, 138-143	3.4	6
44	Determination of naturally occurring progestogens in bovine milk as their oxime derivatives using high performance liquid chromatography-electrospray ionization-tandem mass spectrometry. <i>Journal of the Science of Food and Agriculture</i> , 2010 , 90, 1621-7	4.3	9
43	Quantitative LC-MS/MS method for the sensitive and simultaneous determination of natural hormones in bovine serum. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009 , 877, 2457-64	3.2	39
42	Use of acrylic acid in the synthesis of molecularly imprinted polymers for the analysis of cyproheptadine. <i>Materials Science and Engineering C</i> , 2009 , 29, 398-404	8.3	16
41	Development and validation of an LC-MS/MS confirmatory method for residue analysis of cyproheptadine in urine of food-producing animals. <i>Journal of Agricultural and Food Chemistry</i> , 2009 , 57, 2595-8	5.7	9
40	Influence of farming methods on microbiological contamination and prevalence of resistance to antimicrobial drugs in isolates from beef. <i>Meat Science</i> , 2009 , 82, 284-8	6.4	23
39	Microbiological quality and antimicrobial resistance of Escherichia coli and Staphylococcus aureus isolated from conventional and organic Arz - Ulloal Lheese. CYTA - Journal of Food, 2009, 7, 103-110	2.3	12
38	Antimicrobial resistance in Enterobacteriaceae strains isolated from organic chicken, conventional chicken and conventional turkey meat: A comparative survey. <i>Food Control</i> , 2008 , 19, 412-416	6.2	55
37	Development of a rapid and confirmatory procedure to detect 17beta-estradiol 3-benzoate treatments in bovine hair. <i>Journal of Agricultural and Food Chemistry</i> , 2008 , 56, 11607-11	5.7	9
36	Evolution of resistance in poultry intestinal Escherichia coli during three commonly used antimicrobial therapeutic treatments in poultry. <i>Poultry Science</i> , 2008 , 87, 1643-8	3.9	29
35	Comparison of antimicrobial resistance in Escherichia coli, Staphylococcus aureus, and Listeria monocytogenes strains isolated from organic and conventional poultry meat. <i>Journal of Food Protection</i> , 2008 , 71, 2537-42	2.5	41
34	Antimicrobial resistance in Escherichia coli strains isolated from organic and conventional pork meat: a comparative survey. <i>European Food Research and Technology</i> , 2008 , 226, 371-375	3.4	18
33	Liquid chromatography-electrospray ionization-mass spectrometry method in multiple reaction monitoring mode to determine 17alpha-ethynylestradiol residues in cattle hair without previous digestion. <i>Journal of Agricultural and Food Chemistry</i> , 2007 , 55, 9325-9	5.7	8
32	Antimicrobial resistance in Enterococcus spp. strains isolated from organic chicken, conventional chicken, and turkey meat: a comparative survey. <i>Journal of Food Protection</i> , 2007 , 70, 1021-4	2.5	43
31	Study of a room temperature phosphorescence phenomenon to allow the detection of aflatoxigenic strains in culture media. <i>International Journal of Food Microbiology</i> , 2007 , 115, 149-58	5.8	7
30	Cryogenic grinding pre-treatment improves extraction efficiency of fluoroquinolones for HPLC-MS/MS determination in animal tissue. <i>Analytical and Bioanalytical Chemistry</i> , 2007 , 387, 1933-7	4.4	18
29	Effect of cooking on enrofloxacin residues in chicken tissue. <i>Food Additives and Contaminants</i> , 2006 , 23, 988-93		29
28	Study of enrofloxacin depletion in the eggs of laying hens using diphasic dialysis extraction/purification and determinative HPLC-MS analysis. <i>Journal of Agricultural and Food Chemistry</i> , 2005 , 53, 2849-52	5.7	46

(1999-2005)

27	Study of interferences by several metabolites from Aspergillus spp. in the detection of aflatoxigenic strains in media added with cyclodextrin. <i>Food Control</i> , 2005 , 16, 445-450	6.2	14
26	Evaluation of Chromocult enterococci agar for the isolation and selective enumeration of Enterococcus spp. in broilers. <i>Letters in Applied Microbiology</i> , 2005 , 41, 153-6	2.9	16
25	Detection of synthetic corticosteroids in bovine urine by chemiluminescence high-performance liquid chromatography. <i>Luminescence</i> , 2005 , 20, 197-204	2.5	16
24	Shelf life extension of beef retail cuts subjected to an advanced vacuum skin packaging system. <i>European Food Research and Technology</i> , 2004 , 218, 118-122	3.4	30
23	A survey of the fungal contamination and presence of ochratoxin A and zearalenone on Spanish feed and raw materials. <i>Journal of the Science of Food and Agriculture</i> , 2004 , 84, 832-840	4.3	22
22	Microbiological and physicochemical properties of fresh retail cuts of beef packaged under an advanced vacuum skin system and stored at 4 degrees C. <i>Journal of Food Protection</i> , 2003 , 66, 2085-92	2.5	18
21	Application of a modified culture medium for the simultaneous counting of molds and yeasts and detection of aflatoxigenic strains of Aspergillus flavus and Aspergillus parasiticus. <i>Journal of Food Protection</i> , 2003 , 66, 311-8	2.5	5
20	Development of a method for direct visual determination of aflatoxin production by colonies of the Aspergillus flavus group. <i>International Journal of Food Microbiology</i> , 2003 , 83, 219-25	5.8	27
19	Use of the diphasic dialysis as a new extraction procedure in the determination of enrofloxacin and ciprofloxacin in egg. <i>Analytica Chimica Acta</i> , 2003 , 480, 123-130	6.6	23
18	Development of a diphasic dialysis method for the extraction/purification of residues of ethinylestradiol in hair of cattle, and determination by gas chromatography-tandem mass spectrometry. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life	3.2	12
17	Application of the luminol chemiluminescence reaction for the determination of nine corticosteroids. <i>Analytica Chimica Acta</i> , 2002 , 468, 43-52	6.6	19
16	Gas chromatography-tandem mass spectrometry determination of 17alpha-ethinylestradiol residue in the hair of cattle. Application to treated animals. <i>Journal of Agricultural and Food Chemistry</i> , 2002 , 50, 436-40	5.7	26
15	Inhibitory effects of eugenol and thymol on Penicillium citrinum strains in culture media and cheese. <i>International Journal of Food Microbiology</i> , 2001 , 67, 157-63	5.8	109
14	New additive for culture media for rapid identification of aflatoxin-producing Aspergillus strains. <i>Applied and Environmental Microbiology</i> , 2001 , 67, 4858-62	4.8	64
13	Application of the assay of aflatoxins by liquid chromatography with fluorescence detection in food analysis. <i>Journal of Chromatography A</i> , 2000 , 882, 1-10	4.5	144
12	Determination of quinolones in animal tissues and eggs by high-performance liquid chromatography with photodiode-array detection. <i>Journal of Chromatography A</i> , 2000 , 871, 31-6	4.5	92
11	Chemiluminescence detection of nine corticosteroids in liver. <i>Analyst, The</i> , 2000 , 125, 2071-4	5	17
10	Determination of clenbuterol residues in bovine hair by using diphasic dialysis and gas chromatography-mass spectrometry. <i>Biomedical Applications</i> , 1999 , 726, 133-9		34

9	heptakis-di-O-methyl-Ecyclodextrin as post-column HPLC reagent. <i>Analytical Communications</i> , 1999 , 36, 5-7		15
8	Determination of dexamethasone in bovine liver by chemiluminescence high-performance liquid chromatography. <i>Journal of Agricultural and Food Chemistry</i> , 1999 , 47, 4275-9	5.7	16
7	Interaction between cyclodextrins and aflatoxins Q1, M1 and P1. Fluorescence and chromatographic studies. <i>Journal of Chromatography A</i> , 1998 , 815, 21-9	4.5	33
6	GROWTH OF ESCHERICHIA COLI O157:H7 IN THE PRESENCE OF PSEUDOMONAS FLUORESCENS IN SKIMMED MILK AT 7 OR 25C. <i>Journal of Food Safety</i> , 1997 , 16, 273-285	2	4
5	Determination of residues of the beta-agonist clenbuterol in liver of medicated farm animals by gas chromatography-mass spectrometry using diphasic dialysis as an extraction procedure. <i>Biomedical Applications</i> , 1997 , 693, 321-6		26
4	Rapid semi-quantitative fluorimetric determination of citrinin in fungal cultures isolated from cheese and cheese factories. <i>Letters in Applied Microbiology</i> , 1997 , 24, 397-400	2.9	14
3	Simple and sensitive high-performance liquid chromatographyfluorescence method for the determination of citrinin application to the analysis of fungal cultures and cheese extracts. <i>Journal of Chromatography A</i> , 1996 , 723, 69-75	4.5	44
2	Postcolumn excitation of aflatoxins using cyclodextrins in liquid chromatography for food analysis. Journal of Chromatography A, 1996 , 721, 69-74	4.5	31
1	Simultaneous high-performance liquid chromatographic determination of ochratoxin A and citrinin in cheese by time-resolved luminescence using terbium. <i>Journal of Chromatography A</i> 1996 , 727, 185-19	4 ·5	51