

Inger Mechlenburg

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/206877/publications.pdf>

Version: 2024-02-01

123
papers

1,842
citations

304368

22
h-index

360668

35
g-index

129
all docs

129
docs citations

129
times ranked

1760
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence of knee contractures is high in children with cerebral palsy in Denmark. <i>Physiotherapy Theory and Practice</i> , 2023, 39, 200-207.	0.6	1
2	Is the Femoral-Epiphyseal Acetabular Roof (FEAR) index associated with hip pain in patients with hip dysplasia?. <i>Acta Radiologica</i> , 2023, 64, 666-674.	0.5	2
3	Exercise booster sessions as a mean to maintain the effect of an exercise-intervention - A systematic review. <i>Physical Therapy Reviews</i> , 2022, 27, 103-113.	0.3	3
4	Socioeconomic status and use of analgesic drugs before and after primary hip arthroplasty: a population-based cohort study of 103,209 patients during 1996â€“2018. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2022, 93, 171-178.	1.2	8
5	Study protocol: Prosthesis versus Active (ProAct) exercise program in patients with glenohumeral osteoarthritis â€” a multicenter, randomized controlled trial. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2022, 93, 303-316.	1.2	1
6	Effectiveness of a bandage to prevent re-dislocation after total hip arthroplasty in patients with a previous hip dislocation. A randomized controlled trial with 12-week follow-up. <i>Clinical Rehabilitation</i> , 2022, , 026921552210814.	1.0	1
7	Early offsetâ€“increasing migration predicts later revision for humeral head resurfacing implants. A randomized controlled radiostereometry trial with 10â€“year clinical followâ€“up. <i>Journal of Orthopaedic Research</i> , 2022, , .	1.2	0
8	Five-Year Follow-up After Hip Arthroscopic Surgery in the Horsens-Aarhus Femoroacetabular Impingement (HAFAI) Cohort. <i>Orthopaedic Journal of Sports Medicine</i> , 2022, 10, 232596712210756.	0.8	5
9	The impact of full-thickness rotator cuff tear on shoulder function and quality of life in patients who sustain a proximal humerus fractureâ€”a prospective cohort study. <i>JSES International</i> , 2022, 6, 268-274.	0.7	0
10	Risk Stratification for Postoperative Pulmonary Complications following Major Cardiothoracic and Abdominal Surgery â€” development of the PPC Risk Prediction Score for Physiotherapists Clinical Decision-making. <i>Physiotherapy Theory and Practice</i> , 2022, , 1-12.	0.6	2
11	Do changes in outcomes following primary and revision hip replacement differ and relate to markers of socioeconomic status? A 1-year population-based cohort study. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2022, 93, 397-404.	1.2	4
12	Exercise With Low-Loads and Concurrent Partial Blood Flow Restriction Combined With Patient Education in Females Suffering From Gluteal Tendinopathy: A Feasibility Study. <i>Frontiers in Sports and Active Living</i> , 2022, 4, 881054.	0.9	6
13	Long-term results after weight loss intervention in knee arthroplasty patients with obesity.. <i>Danish Medical Journal</i> , 2022, 69, .	0.5	0
14	Feasibility and acceptability of a six-month exercise and patient education intervention for patients with HIP dysplasia: A mixed Methods study. <i>Musculoskeletal Science and Practice</i> , 2022, , 102615.	0.6	2
15	Monitoring Physical Behavior in Rehabilitation Using a Machine Learningâ€“Based Algorithm for Thigh-Mounted Accelerometers: Development and Validation Study. <i>JMIR Bioinformatics and Biotechnology</i> , 2022, 3, e38512.	0.4	1
16	Does the use of i-FACTOR bone graft affect bone healing in those undergoing periacetabular osteotomy (PAO) for developmental dysplasia of the hip (DDH)? A retrospective study. <i>Journal of Hip Preservation Surgery</i> , 2022, 9, 165-171.	0.6	1
17	Hip muscle and joint contact forces before, 6 and 12 months after minimally invasive periacetabular osteotomy. <i>HIP International</i> , 2021, 31, 676-682.	0.9	5
18	Pain is frequent in children with cerebral palsy and negatively affects physical activity and participation. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 301-306.	0.7	17

#	ARTICLE	IF	CITATIONS
19	Hip contractures were associated with low gross motor function in children with cerebral palsy. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 1562-1568.	0.7	4
20	Weight Loss Intervention Before Total Knee Replacement: A 12-Month Randomized Controlled Trial. <i>Scandinavian Journal of Surgery</i> , 2021, 110, 3-12.	1.3	26
21	Exercise as Medicine During the Course of Hip Osteoarthritis. <i>Exercise and Sport Sciences Reviews</i> , 2021, 49, 77-87.	1.6	4
22	Developing and validating an accelerometer-based algorithm with machine learning to classify physical activity after acquired brain injury. <i>Brain Injury</i> , 2021, 35, 460-467.	0.6	7
23	Ankle contractures are frequent among children with cerebral palsy and associated with lower gross motor function and degree of spasticity. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 2171-2178.	0.7	3
24	Does Preoperative Pain Catastrophizing Influence Objectively Measured Physical Activity Before and After Total Knee Arthroplasty: A Prospective Cohort Study. <i>Journal for the Measurement of Physical Behaviour</i> , 2021, 4, 53-59.	0.5	0
25	The Benefits and Harms of Early Mobilization and Supervised Exercise Therapy after Non-surgically Treated Proximal Humerus or Distal Radius fracture: A systematic Review and Meta-analysis. <i>Current Reviews in Musculoskeletal Medicine</i> , 2021, 14, 107-129.	1.3	12
26	Does Daily Physical Activity Differ Between Patients with Femoroacetabular Impingement Syndrome and Patients with Hip Dysplasia? A Cross-Sectional Study in 157 Patients and 60 Healthy Volunteers. <i>International Journal of Sports Physical Therapy</i> , 2021, 16, 1084-1092.	0.5	6
27	Total hip arthroplasty versus progressive resistance training in patients with severe hip osteoarthritis: protocol for a multicentre, parallel-group, randomised controlled superiority trial. <i>BMJ Open</i> , 2021, 11, e051392.	0.8	3
28	Partial remission of Type 2 diabetes and changes in quality of life after gastric bypass. <i>Danish Medical Journal</i> , 2021, 68, .	0.5	0
29	Effects of Low-Load Blood-Flow Restricted Resistance Training on Functional Capacity and Patient-Reported Outcome in a Young Male Suffering From Reactive Arthritis. <i>Frontiers in Sports and Active Living</i> , 2021, 3, 798902.	0.9	6
30	Despite patient-reported outcomes improve, patients with femoroacetabular impingement syndrome do not increase their objectively measured sport and physical activity level 1 year after hip arthroscopic surgery. Results from the HAFAL cohort. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2020, 28, 1639-1647.	2.3	17
31	Efficacy of preoperative progressive resistance training in patients undergoing total knee arthroplasty: 12-month follow-up data from a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2020, 34, 82-90.	1.0	15
32	Fractures after stroke – A Danish register-based study of 106 001 patients. <i>Acta Neurologica Scandinavica</i> , 2020, 141, 47-55.	1.0	8
33	No effect of cognitive behavioral patient education for patients with pain catastrophizing before total knee arthroplasty: a randomized controlled trial. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2020, 91, 98-103.	1.2	42
34	Efficacy of low-load blood flow restricted resistance EXercise in patients with Knee osteoarthritis scheduled for total knee replacement (EXKnee): protocol for a multicentre randomised controlled trial. <i>BMJ Open</i> , 2020, 10, e034376.	0.8	14
35	Is progressive resistance training feasible in patients with symptomatic external snapping hip?. <i>Physiotherapy Theory and Practice</i> , 2020, , 1-13.	0.6	1
36	Low arthroplasty survival after treatment for proximal humerus fracture sequelae: 3,245 shoulder replacements from the Nordic Arthroplasty Register Association. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2020, , 1-6.	1.2	8

#	ARTICLE	IF	CITATIONS
37	Does pain and hip function improve 2 years after reverse periacetabular osteotomy? A follow-up study of 74 patients. <i>Journal of Hip Preservation Surgery</i> , 2020, 7, 130-139.	0.6	3
38	Patients undergoing shoulder arthroplasty for failed nonoperative treatment of proximal humerus fracture have low implant survival and low patient-reported outcomes: 837 cases from the Danish Shoulder Arthroplasty Registry. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2020, 91, 319-325.	1.2	16
39	14-year hip survivorship after periacetabular osteotomy: a follow-up study on 1,385 hips. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2020, 91, 299-305.	1.2	24
40	Daily activity and functional performance in people with chronic disease: A cross-sectional study. <i>Cogent Medicine</i> , 2020, 7, .	0.7	4
41	Accelerometer-Assessed Prolonged Sitting During Work and Leisure Time and Associations With Age, Body Mass Index, and Health: A Cross-Sectional Study. <i>Journal for the Measurement of Physical Behaviour</i> , 2020, 3, 211-218.	0.5	0
42	Effectiveness of supervised resistance training for patients with hip osteoarthritis - a systematic review. <i>Danish Medical Journal</i> , 2020, 67, .	0.5	1
43	Effects of supervised exercise compared to non-supervised exercise early after total hip replacement on patient-reported function, pain, health-related quality of life and performance-based function – a systematic review and meta-analysis of randomized controlled trials. <i>Clinical Rehabilitation</i> , 2019, 33, 13-23.	1.0	24
44	Association between periacetabular osteotomy and hip dysplasia among relatives: a cross-sectional study. <i>HIP International</i> , 2019, 29, 424-429.	0.9	4
45	Isokinetic dynamometry and gait analysis reveal different hip joint status in patients with hip dysplasia. <i>HIP International</i> , 2019, 29, 215-221.	0.9	9
46	High relative reliability and responsiveness of the forgotten joint score-12 in patients with femoroacetabular impingement undergoing hip arthroscopic treatment. A prospective survey-based study. <i>Journal of Hip Preservation Surgery</i> , 2019, 6, 149-156.	0.6	14
47	Operative versus non-operative treatment for 2-part proximal humerus fracture: A multicenter randomized controlled trial. <i>PLoS Medicine</i> , 2019, 16, e1002855.	3.9	57
48	Improved Patient-Reported Quality of Life and Hip Function After Cementless 1-Stage Revision of Chronic Periprosthetic Hip Joint Infection. <i>Journal of Arthroplasty</i> , 2019, 34, 2763-2769.e1.	1.5	4
49	The correlation between activity level, serum-ion concentrations and pseudotumours in patients with metal-on-metal hip articulations and metal-on-polyethylene total hip articulations. <i>Journal of Orthopaedic Translation</i> , 2019, 18, 74-83.	1.9	7
50	Association between pain catastrophizing, physical function and pain at first visit in the outpatient knee clinic. <i>Knee</i> , 2019, 26, 1286-1291.	0.8	6
51	The association between pain catastrophizing, physical function and pain in a cohort of patients undergoing knee arthroplasty. <i>BMC Musculoskeletal Disorders</i> , 2019, 20, 421.	0.8	52
52	Superior fixation and less periprosthetic stress-shielding of tibial components with a finned stem versus an I-beam block stem: a randomized RSA and DXA study with minimum 5 years™ follow-up. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2019, 90, 165-171.	1.2	10
53	Are changes in pain associated with changes in quality of life and hip function 2 years after periacetabular osteotomy? A follow-up study of 321 patients. <i>Journal of Hip Preservation Surgery</i> , 2019, 6, 69-76.	0.6	23
54	Reference values and variation of acetabular angles measured by computed tomography in 170 asymptomatic hips. <i>Acta Radiologica</i> , 2019, 60, 895-901.	0.5	4

#	ARTICLE	IF	CITATIONS
55	High frequency of labral pathology in symptomatic borderline dysplasia: a prospective magnetic resonance arthrography study of 99 patients. <i>Journal of Hip Preservation Surgery</i> , 2019, 6, 60-68.	0.6	14
56	Use of accelerometer-based activity monitoring in orthopaedics: benefits, impact and practical considerations. <i>EFORT Open Reviews</i> , 2019, 4, 678-685.	1.8	14
57	Efficacy of periacetabular osteotomy followed by progressive resistance training compared to progressive resistance training as non-surgical treatment in patients with hip dysplasia (PreserveHip) – a protocol for a randomised controlled trial. <i>BMJ Open</i> , 2019, 9, e032782.	0.8	7
58	Validation of an inertial measurement unit to determine countermovement jump height. <i>Asia-Pacific Journal of Sports Medicine, Arthroscopy, Rehabilitation and Technology</i> , 2019, 16, 8-13.	0.4	13
59	Is hip muscle strength normalised in patients with femoroacetabular impingement syndrome one year after surgery?. <i>Journal of Science and Medicine in Sport</i> , 2019, 22, 413-419.	0.6	19
60	Patient-reported outcome and muscle-tendon pain after periacetabular osteotomy are related: 1-year follow-up in 82 patients with hip dysplasia. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2019, 90, 40-45.	1.2	17
61	Is high body mass index a potential risk factor for poor outcome after hip arthroplasty? A cohort study of 98 patients 1 year after surgery. <i>Acta Orthopaedica Belgica</i> , 2019, 85, 91-99.	0.1	0
62	Asymmetry and pelvic movements 6 months after total hip replacement Secondary analyses from a randomized controlled trial. <i>Acta Orthopaedica Belgica</i> , 2019, 85, 338-345.	0.1	0
63	Muscle-tendon-related pain in 100 patients with hip dysplasia: prevalence and associations with self-reported hip disability and muscle strength. <i>Journal of Hip Preservation Surgery</i> , 2018, 5, 39-46.	0.6	31
64	Higher Prevalence of Mixed or Solid Pseudotumors in Metal-on-Polyethylene Total Hip Arthroplasty Compared With Metal-on-Metal Total Hip Arthroplasty and Resurfacing Hip Arthroplasty. <i>Journal of Arthroplasty</i> , 2018, 33, 2279-2286.	1.5	25
65	The Association Between Weight Loss and Quality of Life 1 and 5 Years After Laparoscopic Roux-en-Y Gastric Bypass in Danish Bariatric Patients. <i>Obesity Surgery</i> , 2018, 28, 1622-1628.	1.1	2
66	Reliability of the Danish version of the short questionnaire to assess health-enhancing physical activity (SQUASH). <i>Physiotherapy Theory and Practice</i> , 2018, 34, 637-642.	0.6	2
67	Muscle-tendon-related abnormalities detected by ultrasonography are common in symptomatic hip dysplasia. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2018, 138, 1059-1067.	1.3	14
68	Physical Activity Is Associated With the Level of Chromium but Not With Changes in Pseudotumor Size in Patients With Metal-on-Metal Hip Arthroplasty. <i>Journal of Arthroplasty</i> , 2018, 33, 2932-2939.	1.5	9
69	Comparison of Early High-Intensity and Low-Intensity Rehabilitation After Total Knee Arthroplasty: Comment on the Article by Bade et al. <i>Arthritis Care and Research</i> , 2018, 70, 1717-1717.	1.5	1
70	Impaired hip muscle strength in patients with symptomatic femoroacetabular impingement. <i>Journal of Science and Medicine in Sport</i> , 2018, 21, 334.	0.6	1
71	No Exacerbation of Knee Joint Pain and Effusion Following Preoperative Progressive Resistance Training in Patients Scheduled for Total Knee Arthroplasty: Secondary Analyses From a Randomized Controlled Trial. <i>PM and R</i> , 2018, 10, 687-692.	0.9	10
72	What level of pain reduction can be expected up to two years after periacetabular osteotomy? A prospective cohort study of 146 patients. <i>Journal of Hip Preservation Surgery</i> , 2018, 5, 274-281.	0.6	8

#	ARTICLE	IF	CITATIONS
73	Patient-reported quality of life and hip function after 2-stage revision of chronic periprosthetic hip joint infection: a cross-sectional study. <i>HIP International</i> , 2018, 28, 407-414.	0.9	28
74	Does the physical activity profile change in patients with hip dysplasia from before to 1 year after periacetabular osteotomy?. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2018, 89, 622-627.	1.2	16
75	Are patients with knee osteoarthritis and patients with knee joint replacement as physically active as healthy persons?. <i>Journal of Orthopaedic Translation</i> , 2018, 14, 8-15.	1.9	30
76	Progressive resistance training in patients with hip dysplasia: A feasibility study. <i>Journal of Rehabilitation Medicine</i> , 2018, 50, 751-758.	0.8	18
77	Intraobserver and interobserver reliability of recategorized Neer classification in differentiating 2-part surgical neck fractures from multi-fragmented proximal humeral fractures in 116 patients. <i>Journal of Shoulder and Elbow Surgery</i> , 2018, 27, 1756-1761.	1.2	11
78	Reliability of computer-assisted periacetabular osteotomy using a minimally invasive approach. <i>International Journal of Computer Assisted Radiology and Surgery</i> , 2018, 13, 2021-2028.	1.7	8
79	Leg power, pelvic movement and physical activity after periacetabular osteotomy. A prospective cohort study. <i>Acta Orthopaedica Belgica</i> , 2018, 84, 163-171.	0.1	4
80	Description of load progression and pain response during progressive resistance training early after total hip arthroplasty: secondary analyses from a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2017, 31, 11-22.	1.0	15
81	The efficacy of early initiated, supervised, progressive resistance training compared to unsupervised, home-based exercise after unicompartamental knee arthroplasty: a single-blinded randomized controlled trial. <i>Clinical Rehabilitation</i> , 2017, 31, 61-70.	1.0	24
82	Pain, activities of daily living and sport function at different time points after hip arthroscopy in patients with femoroacetabular impingement: a systematic review with meta-analysis. <i>British Journal of Sports Medicine</i> , 2017, 51, 572-579.	3.1	77
83	Impaired hip muscle strength in patients with femoroacetabular impingement syndrome. <i>Journal of Science and Medicine in Sport</i> , 2017, 20, 1062-1067.	0.6	58
84	Effectiveness of a physiotherapist delivered cognitive-behavioral patient education for patients who undergoes operation for total knee arthroplasty: a protocol of a randomized controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 116.	0.8	22
85	The Use of Ultrasound to Exclude Extremity Fractures in Adults. <i>JBJS Open Access</i> , 2017, 2, e0007.	0.8	3
86	The Association between Gender and Familial Prevalence of Hip Dysplasia in Danish Patients. <i>HIP International</i> , 2017, 27, 299-304.	0.9	8
87	Efficacy of Preoperative Progressive Resistance Training on Postoperative Outcomes in Patients Undergoing Total Knee Arthroplasty. <i>Arthritis Care and Research</i> , 2016, 68, 1239-1251.	1.5	57
88	Does a titanium sleeve reduce the frequency of pseudotumors in metal-on-metal total hip arthroplasty at 5-7 years follow-up?. <i>Orthopaedics and Traumatology: Surgery and Research</i> , 2016, 102, 1035-1041.	0.9	6
89	The Association Between Preoperative Symptoms of Obesity in Knee and Hip Joints and the Change in Quality of Life After Laparoscopic Roux-en-Y Gastric Bypass. <i>Obesity Surgery</i> , 2016, 26, 950-956.	1.1	12
90	No association between pseudotumors, high serum metal-ion levels and metal hypersensitivity in large-head metal-on-metal total hip arthroplasty at 5-7-year follow-up. <i>Skeletal Radiology</i> , 2016, 45, 115-125.	1.2	24

#	ARTICLE	IF	CITATIONS
91	Functional performance is associated with both knee extensor and flexor muscle strength in patients scheduled for total knee arthroplasty: A cross-sectional study. <i>Journal of Rehabilitation Medicine</i> , 2015, 47, 454-459.	0.8	22
92	A study of the inter-rater reliability of a test battery for use in patients after total hip replacement. <i>Clinical Rehabilitation</i> , 2015, 29, 165-174.	1.0	16
93	Cartilage Thickness and Cyst Volume Are Unchanged 10 Years After Periacetabular Osteotomy in Patients Without Hip Symptoms. <i>Clinical Orthopaedics and Related Research</i> , 2015, 473, 2644-2649.	0.7	11
94	Pelvic movement strategies and leg extension power in patients with end-stage medial compartment knee osteoarthritis: a cross-sectional study. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2015, 135, 1217-1226.	1.3	10
95	Walking patterns and hip contact forces in patients with hip dysplasia. <i>Gait and Posture</i> , 2015, 42, 529-533.	0.6	40
96	The Horsens-Aarhus Femoro Acetabular Impingement (HAFAl) cohort: outcome of arthroscopic treatment for femoroacetabular impingement. Protocol for a prospective cohort study. <i>BMJ Open</i> , 2015, 5, e008952.	0.8	10
97	No correlations between radiological angles and self-assessed quality of life in patients with hip dysplasia at 13 years of follow-up after periacetabular osteotomy. <i>Acta Radiologica</i> , 2015, 56, 196-203.	0.5	7
98	Progressive resistance training before and after total hip and knee arthroplasty: a systematic review. <i>Clinical Rehabilitation</i> , 2015, 29, 14-29.	1.0	50
99	Joint kinematics and kinetics during walking and running in 32 patients with hip dysplasia 1 year after periacetabular osteotomy. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2014, 85, 592-599.	1.2	27
100	No association between serum metal ions and implant fixation in large-head metal-on-metal total hip arthroplasty. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2014, 85, 355-362.	1.2	10
101	Effect of early supervised progressive resistance training compared to unsupervised home-based exercise after fast-track total hip replacement applied to patients with preoperative functional limitations. A single-blinded randomised controlled trial. <i>Osteoarthritis and Cartilage</i> , 2014, 22, 2051-2058.	0.6	53
102	Evaluation of periprosthetic bone mineral density and postoperative migration of humeral head resurfacing implants: two-year results of a randomized controlled clinical trial. <i>Journal of Shoulder and Elbow Surgery</i> , 2014, 23, 1427-1436.	1.2	14
103	Block-step asymmetry 5 years after large-head metal-on-metal total hip arthroplasty is related to lower muscle mass and leg power on the implant side. <i>Clinical Biomechanics</i> , 2014, 29, 684-690.	0.5	13
104	Blood perfusion and bone formation before and after minimally invasive periacetabular osteotomy analysed by Positron Emission Tomography combined with Computed Tomography. <i>International Orthopaedics</i> , 2013, 37, 789-794.	0.9	3
105	The Copeland resurfacing humeral head implant does not restore humeral head anatomy. A retrospective study. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2013, 133, 615-619.	1.3	45
106	Changes in walking and running in patients with hip dysplasia. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2013, 84, 265-270.	1.2	28
107	Automated measurement of diagnostic angles for hip dysplasia. , 2013, , .		6
108	Overweight preoperatively impairs clinical outcome after knee arthroplasty. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2013, 84, 392-397.	1.2	42

#	ARTICLE	IF	CITATIONS
109	Cyst Volume in the Acetabulum and Femoral Head Decreases after Periacetabular Osteotomy. <i>HIP International</i> , 2012, 22, 313-318.	0.9	7
110	Precision of novel radiological methods in relation to resurfacing humeral head implants: assessment by radiostereometric analysis, DXA, and geometrical analysis. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2012, 132, 1521-1530.	1.3	17
111	Cartilage Thickness in the Hip Measured by MRI and Stereology Before and After Periacetabular Osteotomy. <i>Clinical Orthopaedics and Related Research</i> , 2010, 468, 1884-1890.	0.7	19
112	Total Hip Replacement in the Congenitally Dislocated Hip Using the Paavilainen Technique. , 2010, , 61-65.		0
113	Radiation exposure to the orthopaedic surgeon during periacetabular osteotomy. <i>International Orthopaedics</i> , 2009, 33, 1747-1751.	0.9	34
114	What is the role of clinical tests and ultrasound in acetabular labral tear diagnostics?. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2009, 80, 314-318.	1.2	94
115	Total hip replacement in the congenitally dislocated hip using the Paavilainen technique. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2009, 80, 259-262.	1.2	20
116	No change detected by DEXA in bone mineral density after periacetabular osteotomy. <i>Acta Orthopaedica Belgica</i> , 2009, 75, 761-6.	0.1	3
117	Evaluation of Bernese periacetabular osteotomy. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2008, 79, 1-43.	1.2	13
118	Safe fixation with two acetabular screws after Ganz periacetabular osteotomy. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2007, 78, 344-349.	1.2	20
119	Cartilage thickness in the hip joint measured by MRI and stereology – a methodological study. <i>Osteoarthritis and Cartilage</i> , 2007, 15, 366-371.	0.6	35
120	Prospective bone density changes after periacetabular osteotomy: a methodological study. <i>International Orthopaedics</i> , 2005, 29, 281-286.	0.9	3
121	Changes in load-bearing area after Ganz periacetabular osteotomy. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2005, 76, 141-142.	1.2	2
122	Changes in load-bearing area after Ganz periacetabular osteotomy evaluated by multislice CT scanning and stereology. <i>Acta Orthopaedica</i> , 2004, 75, 147-153.	1.4	55
123	Effects of Resistance Training Prior to Total Hip or Knee Replacement on Post-operative Recovery in Functional Performance: A Systematic Review and Meta-Analysis. <i>Frontiers in Sports and Active Living</i> , 0, 4, .	0.9	5