

Tae-Hwa Chun

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2075705/tae-hwa-chun-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

43
papers

3,098
citations

28
h-index

43
g-index

43
ext. papers

3,379
ext. citations

7.7
avg, IF

4.56
L-index

#	Paper	IF	Citations
43	HIF2A-LOX Pathway Promotes Fibrotic Tissue Remodeling in Thyroid-Associated Orbitopathy. <i>Endocrinology</i> , 2019 , 160, 20-35	4.8	36
42	Fibro-Adipogenic Remodeling of the Diaphragm in Obesity-Associated Respiratory Dysfunction. <i>Diabetes</i> , 2019 , 68, 45-56	0.9	23
41	Transcription factor 21 (TCF21) promotes proinflammatory interleukin 6 expression and extracellular matrix remodeling in visceral adipose stem cells. <i>Journal of Biological Chemistry</i> , 2018 , 293, 6603-6610	5.4	12
40	Designing 3-D Adipospheres for Quantitative Metabolic Study. <i>Methods in Molecular Biology</i> , 2017 , 1566, 177-183	1.4	4
39	Membrane-Tethered Metalloproteinase Expressed by Vascular Smooth Muscle Cells Limits the Progression of Proliferative Atherosclerotic Lesions. <i>Journal of the American Heart Association</i> , 2017 , 6,	6	10
38	Immunomagnetic Separation of Fat Depot-specific Sca1 ^{high} Adipose-derived Stem Cells (ASCs). <i>Journal of Visualized Experiments</i> , 2016 ,	1.6	1
37	Adipose extracellular matrix remodelling in obesity and insulin resistance. <i>Biochemical Pharmacology</i> , 2016 , 119, 8-16	6	117
36	Thrombospondin 1 as a novel biological marker of obesity and metabolic syndrome. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 1490-9	12.7	44
35	The exocyst complex regulates free fatty acid uptake by adipocytes. <i>PLoS ONE</i> , 2015 , 10, e0120289	3.7	7
34	3-D adipocyte differentiation and peri-adipocyte collagen turnover. <i>Methods in Enzymology</i> , 2014 , 538, 15-34	1.7	13
33	Fat depot-specific gene signature and ECM remodeling of Sca1 ^(high) adipose-derived stem cells. <i>Matrix Biology</i> , 2014 , 36, 28-38	11.4	21
32	On being the right size: scaling effects in designing a human-on-a-chip. <i>Integrative Biology (United Kingdom)</i> , 2013 , 5, 1149-61	3.7	107
31	Thrombospondin 1 mediates high-fat diet-induced muscle fibrosis and insulin resistance in male mice. <i>Endocrinology</i> , 2013 , 154, 4548-59	4.8	50
30	Peri-adipocyte ECM remodeling in obesity and adipose tissue fibrosis. <i>Adipocyte</i> , 2012 , 1, 89-95	3.2	62
29	Adipogenic histone mark regulation by matrix metalloproteinase 14 in collagen-rich microenvironments. <i>Molecular Endocrinology</i> , 2011 , 25, 745-53		17
28	Genetic link between obesity and MMP14-dependent adipogenic collagen turnover. <i>Diabetes</i> , 2010 , 59, 2484-94	0.9	68
27	A pericellular collagenase directs the 3-dimensional development of white adipose tissue. <i>Cell</i> , 2006 , 125, 577-91	56.2	305

26	Angiotensin II suppresses growth arrest specific homeobox (Gax) expression via redox-sensitive mitogen-activated protein kinase (MAPK). <i>Regulatory Peptides</i> , 2005 , 127, 159-67		10
25	Therapeutic potential of thiazolidinediones in activation of peroxisome proliferator-activated receptor gamma for monocyte recruitment and endothelial regeneration. <i>European Journal of Pharmacology</i> , 2005 , 508, 255-65	5.3	36
24	An MT1-MMP-PDGF receptor-beta axis regulates mural cell investment of the microvasculature. <i>Genes and Development</i> , 2005 , 19, 979-91	12.6	91
23	MT1-matrix metalloproteinase directs arterial wall invasion and neointima formation by vascular smooth muscle cells. <i>Journal of Experimental Medicine</i> , 2005 , 202, 663-71	16.6	103
22	MT1-MMP-dependent neovessel formation within the confines of the three-dimensional extracellular matrix. <i>Journal of Cell Biology</i> , 2004 , 167, 757-67	7.3	281
21	Significance and therapeutic potential of the natriuretic peptides/cGMP/cGMP-dependent protein kinase pathway in vascular regeneration. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 3404-9	11.5	139
20	Modification of GATA-2 transcriptional activity in endothelial cells by the SUMO E3 ligase PIASy. <i>Circulation Research</i> , 2003 , 92, 1201-8	15.7	53
19	C-type natriuretic peptide induces redifferentiation of vascular smooth muscle cells with accelerated reendothelialization. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2001 , 21, 930-6	9.4	67
18	Oxidized LDL regulates vascular endothelial growth factor expression in human macrophages and endothelial cells through activation of peroxisome proliferator-activated receptor-gamma. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2001 , 21, 560-6	9.4	137
17	Thiazolidinediones, peroxisome proliferator-activated receptor gamma agonists, regulate endothelial cell growth and secretion of vasoactive peptides. <i>Atherosclerosis</i> , 2001 , 158, 113-9	3.1	90
16	Coordinate regulation of endothelin and adrenomedullin secretion by oxidative stress in endothelial cells. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2001 , 281, H1364-71	5.2	28
15	Oxidative stress augments secretion of endothelium-derived relaxing peptides, C-type natriuretic peptide and adrenomedullin. <i>Journal of Hypertension</i> , 2000 , 18, 575-80	1.9	63
14	Altered gene expression of uncoupling protein-2 and -3 in stroke-prone spontaneously hypertensive rats. <i>Journal of Hypertension</i> , 2000 , 18, 1233-8	1.9	13
13	Inhibition of rho-associated kinase results in suppression of neointimal formation of balloon-injured arteries. <i>Circulation</i> , 2000 , 101, 2030-3	16.7	172
12	Down regulation of peroxisome proliferator-activated receptor gamma expression by inflammatory cytokines and its reversal by thiazolidinediones. <i>Diabetologia</i> , 1999 , 42, 702-10	10.3	114
11	Vascular endothelial growth factor (VEGF) expression in human coronary atherosclerotic lesions: possible pathophysiological significance of VEGF in progression of atherosclerosis. <i>Circulation</i> , 1998 , 98, 2108-16	16.7	389
10	Regulation of endothelial production of C-type natriuretic peptide by interaction between endothelial cells and macrophages. <i>Endocrinology</i> , 1998 , 139, 1920-6	4.8	34
9	Effects of intravenously administered C-type natriuretic peptide in humans: comparison with atrial natriuretic peptide. <i>Hypertension Research</i> , 1998 , 21, 7-13	4.7	40

8	Physiologic shear stress suppresses endothelin-converting enzyme-1 expression in vascular endothelial cells. <i>Journal of Cardiovascular Pharmacology</i> , 1998 , 31 Suppl 1, S42-5	3.1	29
7	Opposite regulation of Gax homeobox expression by angiotensin II and C-type natriuretic peptide. <i>Hypertension</i> , 1997 , 29, 381-7	8.5	32
6	Adenovirus-mediated gene transfer of C-type natriuretic peptide causes G1 growth inhibition of cultured vascular smooth muscle cells. <i>Biochemical and Biophysical Research Communications</i> , 1997 , 239, 889-94	3.4	24
5	Significance of vascular natriuretic peptide system in vascular remodeling in humans and its application to gene therapy. <i>Annals of the New York Academy of Sciences</i> , 1997 , 811, 533-41	6.5	16
4	Shear stress augments expression of C-type natriuretic peptide and adrenomedullin. <i>Hypertension</i> , 1997 , 29, 1296-302	8.5	148
3	Cellular and Molecular Aspects of C-Type Natriuretic Peptide (CNP) 1997 , 107-122		1
2	cDNA cloning and gene expression of human type Ialpha cGMP-dependent protein kinase. <i>Hypertension</i> , 1996 , 27, 552-7	8.5	57
1	Vascular endothelial growth factor suppresses C-type natriuretic peptide secretion. <i>Hypertension</i> , 1996 , 27, 811-5	8.5	34