Helle Kieler

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/2078306/helle-kieler-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

57	1,490	15	38
papers	citations	h-index	g-index
60	1,863 ext. citations	4.7	4.39
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
57	Temporal trends in lung cancer survival: a population-based study. Acta Oncolgica, 2021, 1-7	3.2	1
56	Influence of statin-potency on the risk of kidney disease - A nationwide cohort study using laboratory data. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 210-219	2.6	О
55	Antibiotic use prior to a lung cancer diagnosis: a population-based study. <i>Cancer Causes and Control</i> , 2021 , 32, 597-607	2.8	2
54	Association of Childhood Growth Hormone Treatment With Long-term Cardiovascular Morbidity. <i>JAMA Pediatrics</i> , 2021 , 175, e205199	8.3	11
53	Use of sildenafil and other phosphodiesterase type 5 inhibitors among pregnant women in Scandinavia. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2021 , 100, 2111-2118	3.8	O
52	Prevalence trends and individual patterns of antiepileptic drug use in pregnancy 2006-2016: A study in the five Nordic countries, United States, and Australia. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 913-922	2.6	8
51	Anti-TNF treatment during pregnancy and birth outcomes: A population-based study from Denmark, Finland, and Sweden. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 316-327	2.6	20
50	Cancer risk in patients with primary immune thrombocytopenia - A Swedish nationwide register study. <i>Cancer Epidemiology</i> , 2020 , 69, 101806	2.8	2
49	Metformin versus insulin use for treatment of gestational diabetes and delivery by caesarean section: A nationwide Swedish cohort study. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2020 , 254, 271-276	2.4	3
48	Paediatric infections in the first 3 years of life after maternal anti-TNF treatment during pregnancy. <i>Alimentary Pharmacology and Therapeutics</i> , 2020 , 52, 843-854	6.1	10
47	Opioid maintenance treatment of pregnant women in the Scandinavian countries <i>NAD Nordic Studies on Alcohol and Drugs</i> , 2020 , 37, 298-312	1.1	O
46	Assessing Safety of Direct Thrombin Inhibitors, Direct Factor Xa Inhibitors and Vitamin K Antagonists in Patients with Atrial Fibrillation: A Nation-Wide Propensity Score Matched Cohort from Sweden. <i>Clinical Epidemiology</i> , 2020 , 12, 1029-1038	5.9	1
45	In utero opioid exposure and risk of infections in childhood: A multinational Nordic cohort study. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 1596-1604	2.6	1
44	Medication persistence to antihypertensive drug treatment - a cross-sectional study of attitudes towards hypertension and medication in persistent and non-persistent patients. <i>Blood Pressure</i> , 2019 , 28, 309-316	1.7	4
43	Adult height is associated with risk of cancer and mortality in 5.5 million Swedish women and men. <i>Journal of Epidemiology and Community Health</i> , 2019 , 73, 730-736	5.1	5
42	Monotherapy vs. combination therapy for post mania maintenance treatment: A population based cohort study. <i>European Neuropsychopharmacology</i> , 2019 , 29, 691-700	1.2	6
41	Impact of risk factors on the occurrence of arterial thrombosis and venous thromboembolism in adults with primary immune thrombocytopenia - Results from two nationwide cohorts. <i>Thrombosis Research</i> , 2019 , 178, 124-131	8.2	11

40	Exposure to antimuscarinic medications for treatment of overactive bladder and risk of lung cancer and colon cancer. <i>Clinical Epidemiology</i> , 2019 , 11, 133-143	5.9	3
39	Clinical characteristics and survival in non-small cell lung cancer patients by smoking history: a population-based cohort study. <i>Acta Oncolgica</i> , 2019 , 58, 1618-1627	3.2	8
38	Adherence to Swedish national pregnancy dating guidelines and management of discrepancies between pregnancy dating methods: a survey study. <i>Reproductive Health</i> , 2019 , 16, 95	3.5	3
37	Patient characteristics when starting treatment and patterns of treatment in adults with chronic immune thrombocytopenia. <i>Blood Coagulation and Fibrinolysis</i> , 2019 , 30, 350-356	1	3
36	Antidiabetic medication use during pregnancy: an international utilization study. <i>BMJ Open Diabetes Research and Care</i> , 2019 , 7, e000759	4.5	18
35	Systematic misclassification of gestational age by ultrasound biometry: implications for clinical practice and research methodology in the Nordic countries. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2018 , 97, 440-444	3.8	9
34	Long-term quality of life and postthrombotic syndrome in women after an episode of venous thromboembolism. <i>Phlebology</i> , 2018 , 33, 234-241	2	2
33	Effect of Maternal Psoriasis on Pregnancy and Birth Outcomes: A Population-based Cohort Study from Denmark and Sweden. <i>Acta Dermato-Venereologica</i> , 2018 , 98, 728-734	2.2	45
32	Discrepancy between pregnancy dating methods affects obstetric and neonatal outcomes: a population-based register cohort study. <i>Scientific Reports</i> , 2018 , 8, 6936	4.9	4
31	Association Between Methylphenidate and Amphetamine Use in Pregnancy and Risk of Congenital Malformations: A Cohort Study From the International Pregnancy Safety Study Consortium. <i>JAMA Psychiatry</i> , 2018 , 75, 167-175	14.5	60
30	Trends in attention-deficit hyperactivity disorder medication use: a retrospective observational study using population-based databases. <i>Lancet Psychiatry,the</i> , 2018 , 5, 824-835	23.3	105
29	Neuraminidase inhibitors during pregnancy and risk of adverse neonatal outcomes and congenital malformations: population based European register study. <i>BMJ, The</i> , 2017 , 356, j629	5.9	14
28	Reducing the rehospitalization risk after a manic episode: A population based cohort study of lithium, valproate, olanzapine, quetiapine and aripiprazole in monotherapy and combinations. <i>Journal of Affective Disorders</i> , 2017 , 217, 16-23	6.6	10
27	Utilization of Stimulants and Atomoxetine for Attention-Deficit/Hyperactivity Disorder among 5.4 Million Children Using Population-Based Longitudinal Data. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2017 , 120, 373-379	3.1	15
26	International trends in antipsychotic use: A study in 16 countries, 2005-2014. <i>European Neuropsychopharmacology</i> , 2017 , 27, 1064-1076	1.2	112
25	Pharmacological agents preceding a diagnosis of immune thrombocytopenia in adult patients developing the chronic form: A Swedish national register study. <i>Thrombosis Research</i> , 2017 , 160, 27-31	8.2	1
24	Risk of breast cancer in risperidone users: A nationwide cohort study. <i>Schizophrenia Research</i> , 2017 , 182, 98-103	3.6	11
23	Maternal and fetal characteristics affect discrepancies between pregnancy-dating methods: a population-based cross-sectional register study. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2017 , 96, 86-95	3.8	9

22	Influence of baseline low-density lipoprotein cholesterol values on statin therapy persistence. <i>European Journal of Clinical Pharmacology</i> , 2016 , 72, 349-57	2.8	11
21	Maternal and infant characteristics: differences and similarities between the Nordic countries and the US. <i>Clinical Epidemiology</i> , 2016 , 8, 285-94	5.9	3
20	Effects of ultrasound pregnancy dating on neonatal morbidity in late preterm and early term male infants: a register-based cohort study. <i>BMC Pregnancy and Childbirth</i> , 2016 , 16, 335	3.2	9
19	Cardiovascular disease and mortality after a first episode of venous thromboembolism in young and middle-aged women. <i>Thrombosis Research</i> , 2016 , 138, 80-85	8.2	4
18	Use of drugs for ADHD among adults-a multinational study among 15.8 million adults in the Nordic countries. <i>European Journal of Clinical Pharmacology</i> , 2016 , 72, 1507-1514	2.8	25
17	Predictors for initiation of pharmacological prophylaxis in patients with newly diagnosed bipolar disorderA nationwide cohort study. <i>Journal of Affective Disorders</i> , 2015 , 172, 204-10	6.6	5
16	Children with hemodynamically significant congenital heart disease can be identified through population-based registers. <i>Clinical Epidemiology</i> , 2015 , 7, 119-27	5.9	9
15	Use of SSRI and SNRI Antidepressants during Pregnancy: A Population-Based Study from Denmark, Iceland, Norway and Sweden. <i>PLoS ONE</i> , 2015 , 10, e0144474	3.7	58
14	Selective serotonin reuptake inhibitors and venlafaxine in early pregnancy and risk of birth defects: population based cohort study and sibling design. <i>BMJ, The,</i> 2015 , 350, h1798	5.9	119
13	Use of palivizumab is underestimated in the Swedish Prescribed Drug Register - implications for register-based drug studies. <i>Clinical Epidemiology</i> , 2015 , 7, 45-51	5.9	8
12	Post-Thrombotic Syndrome and Quality-of-Life after Venous Thromboembolism in Young and Middle-Aged Women. <i>Blood</i> , 2015 , 126, 2309-2309	2.2	1
11	Influence of coronary artery disease-associated genetic variants on risk of venous thromboembolism. <i>Thrombosis Research</i> , 2014 , 134, 426-32	8.2	15
10	Antibiotics in fetal and early life and subsequent childhood asthma: nationwide population based study with sibling analysis. <i>BMJ, The</i> , 2014 , 349, g6979	5.9	89
9	Birth outcomes in women with inflammatory bowel disease: effects of disease activity and drug exposure. <i>Inflammatory Bowel Diseases</i> , 2014 , 20, 1091-8	4.5	128
8	Risk factors for recurrent venous thromboembolism in young and middle-aged women. <i>Thrombosis Research</i> , 2014 , 133, 762-7	8.2	9
7	Selective serotonin reuptake inhibitors during pregnancy and risk of persistent pulmonary hypertension in the newborn: population based cohort study from the five Nordic countries. <i>BMJ, The</i> , 2012 , 344, d8012	5.9	197
6	Risks of adverse pregnancy and birth outcomes in women treated or not treated with mood stabilisers for bipolar disorder: population based cohort study. <i>BMJ, The</i> , 2012 , 345, e7085	5.9	146
5	Obesity Is a Risk Factor for Recurrent Venous Thrombosis in Young and Middle-Aged Women. Results From Tehs-Follow up, a Nested Cohort Study On 1394 Women with VTE <i>Blood</i> , 2012 , 120, 225	50- 2 250)

LIST OF PUBLICATIONS

4	Drug use during pregnancy in Sweden - assessed by the Prescribed Drug Register and the Medical Birth Register. <i>Clinical Epidemiology</i> , 2011 , 3, 43-50	5.9	103
3	The Nordic health registers - an important source when evaluating the safety of antidepressants during pregnancy. <i>Clinical Epidemiology</i> , 2010 , 2, 205-7	5.9	9
2	Emergency cesarean delivery in induction of labor: an evaluation of risk factors 2005, 84, 456		11
1	Assessing urinary albumin excretion in pre-eclamptic women: which sample to use?. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2003 , 110, 12-17	3.7	13