## Ehsan Ghonchepour

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2082617/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Multicomponent reactionâ $\in$ derived covalent inhibitor space. Science Advances, 2021, 7, .	10.3	24
2	Efficient heterogenization of palladium by citric acid on the magnetite nanoparticles surface (Nano-Fe3O4@CA-Pd), and its catalytic application in C-C coupling reactions. Journal of Organometallic Chemistry, 2019, 883, 1-10.	1.8	18
3	Glycoconjugates via Phosphorus Ylides. European Journal of Organic Chemistry, 2019, 2019, 3632-3635.	2.4	1
4	Bentonite clay as an efficient substrate for the synthesis of the super stable and recoverable magnetic nanocomposite of palladium (Fe3O4/Bentonite-Pd). Polyhedron, 2019, 162, 192-200.	2.2	36
5	Synthesis of recoverable palladium composite as an efficient catalyst for the reduction of nitroarene compounds and Suzuki cross-coupling reactions using sepiolite clay and magnetic nanoparticles (Fe3O4@sepiolite-Pd2+). Comptes Rendus Chimie, 2019, 22, 84-95.	0.5	24
6	Structure and Reactivity of Glycosyl Isocyanides. European Journal of Organic Chemistry, 2019, 2019, 50-55.	2.4	2
7	Three-component reaction for an efficient synthesis of 5-hydroxy-1-phenyl-1H-pyrazoles containing a stable phosphorus ylide moiety. Phosphorus, Sulfur and Silicon and the Related Elements, 2018, 193, 459-463.	1.6	9
8	Transition metal-free and base-mediated transformation arylation of unactivated benzene with aryl halides in presence of N,Nâ€2-bis(salicylidene)ethylenediamine as organocatalyst. Catalysis Communications, 2018, 107, 87-91.	3.3	6
9	Methyl Red as Organocatalyst for Arylation of Unactivated Benzene Derivatives with Aryl Halides. ChemistrySelect, 2018, 3, 11517-11521.	1.5	2
10	Synthesis of 1H-1,3-benzimidazoles, benzothiazoles and 3H-imidazo[4,5-c]pyridine using DMF in the presence of HMDS as a reagent under the transition-metal-free condition. Chemical Papers, 2018, 72, 2973-2978.	2.2	17
11	Preparation and characterization of copper chloride supported on citric acidâ€modified magnetite nanoparticles (Cu <sup>2+</sup> â€CA@Fe <sub>3</sub> O <sub>4</sub> ) and evaluation of its catalytic activity in the reduction of nitroarene compounds. Applied Organometallic Chemistry, 2017, 31, e3822.	3.5	13
12	Encapsulation of Pd(II) into superparamagnetic nanoparticles grafted with EDTA and their catalytic activity towards reduction of nitroarenes and Suzuki–Miyaura coupling. Applied Organometallic Chemistry, 2015, 29, 187-194.	3.5	26
13	FeSO4·7H2O-catalyzed oxidative amidation of methylarenes. Tetrahedron Letters, 2015, 56, 2674-2677.	1.4	16
14	Tributyltin grafted onto the surface of 3-aminopropyl functionalized γ-Fe <sub>2</sub> O <sub>3</sub> nanoparticles: a magnetically-recoverable catalyst for trimethylsilylation of alcohols and phenols. RSC Advances, 2014, 4, 34428.	3.6	5

2