Daisuke Nishi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2083425/publications.pdf

Version: 2024-02-01

185998 223531 2,884 125 28 46 citations h-index g-index papers 134 134 134 4016 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Nutritional medicine as mainstream in psychiatry. Lancet Psychiatry, the, 2015, 2, 271-274.	3.7	375
2	Estimating the risk of PTSD in recent trauma survivors: results of the International Consortium to Predict PTSD (ICPP). World Psychiatry, 2019, 18, 77-87.	4.8	126
3	Reliability and validity of the Japanese version of the Resilience Scale and its short version. BMC Research Notes, 2010, 3, 310.	0.6	118
4	Posttraumatic growth, posttraumatic stress disorder and resilience of motor vehicle accident survivors. BioPsychoSocial Medicine, 2010, 4, 7.	0.9	103
5	<scp>I Iscp>nternational <scp>S Scp>ociety for <scp>N Iscp>escp>P Iscp>P Scp>R Psychiatry Psychiatry</scp></scp></scp>	4.8	81
6	Peritraumatic Distress Inventory as a predictor of postâ€traumatic stress disorder after a severe motor vehicle accident. Psychiatry and Clinical Neurosciences, 2010, 64, 149-156.	1.0	73
7	Peritraumatic Distress, Watching Television, and Posttraumatic Stress Symptoms among Rescue Workers after the Great East Japan Earthquake. PLoS ONE, 2012, 7, e35248.	1.1	67
8	Concern over radiation exposure and psychological distress among rescue workers following the Great East Japan Earthquake Concern over radiation exposure and distress. BMC Public Health, 2012, 12, 249.	1.2	60
9	Prevalence of mental disorders and mental health service use in Japan. Psychiatry and Clinical Neurosciences, 2019, 73, 458-465.	1.0	58
10	Associations between Disaster Exposures, Peritraumatic Distress, and Posttraumatic Stress Responses in Fukushima Nuclear Plant Workers following the 2011 Nuclear Accident: The Fukushima NEWS Project Study. PLoS ONE, 2014, 9, e87516.	1.1	51
11	Clinician guidelines for the treatment of psychiatric disorders with nutraceuticals and phytoceuticals: The World Federation of Societies of Biological Psychiatry (WFSBP) and Canadian Network for Mood and Anxiety Treatments (CANMAT) Taskforce. World Journal of Biological Psychiatry, 2022, 23, 424-455.	1.3	49
12	Epidemiology of overdose episodes from the period prior to hospitalization for drug poisoning until discharge in Japan: An exploratory descriptive study using a nationwide claims database. Journal of Epidemiology, 2017, 27, 373-380.	1.1	48
13	Resilience, postâ€traumatic growth, and work engagement among health care professionals after the Great East Japan Earthquake: A 4â€year prospective followâ€up study. Journal of Occupational Health, 2016, 58, 347-353.	1.0	47
14	Optimal cutâ€off score of the Edinburgh Postnatal Depression Scale for major depressive episode during pregnancy in Japan. Psychiatry and Clinical Neurosciences, 2017, 71, 836-842.	1.0	45
15	Incidence and prediction of psychiatric morbidity after a motor vehicle accident in Japan: The Tachikawa Cohort of Motor Vehicle Accident Study. Critical Care Medicine, 2008, 36, 74-80.	0.4	44
16	Omega-3 Fatty Acids for Secondary Prevention of Posttraumatic Stress Disorder After Accidental Injury. Journal of Clinical Psychopharmacology, 2010, 30, 217-219.	0.7	43
17	Trends in the prevalence of psychological distress and the use of mental health services from 2007 to 2016 in Japan. Journal of Affective Disorders, 2018, 239, 208-213.	2.0	40
18	Reliability and validity of the Japanese version of the Peritraumatic Distress Inventory. General Hospital Psychiatry, 2009, 31, 75-79.	1.2	38

#	Article	IF	Citations
19	Potential Role of Brain-Derived Neurotrophic Factor in Omega–3 Fatty Acid Supplementation to Prevent Posttraumatic Distress after Accidental Injury: An Open-Label Pilot Study. Psychotherapy and Psychosomatics, 2011, 80, 310-312.	4.0	36
20	Fish Oil for Attenuating Posttraumatic Stress Symptoms among Rescue Workers after the Great East Japan Earthquake: A Randomized Controlled Trial. Psychotherapy and Psychosomatics, 2012, 81, 315-317.	4.0	36
21	Culturally sensitive and universal measure of resilience for <scp>J</scp> apanese populations: <scp>T</scp> achikawa <scp>R</scp> esilience <scp>S</scp> cale in comparison with <scp>R</scp> esilience <scp>S</scp> cale 14â€item version. Psychiatry and Clinical Neurosciences, 2013, 67, 174-181.	1.0	36
22	Longitudinal Changes in Serum Brain-Derived Neurotrophic Factor in Accident Survivors with Posttraumatic Stress Disorder. Neuropsychobiology, 2013, 68, 44-50.	0.9	36
23	Comparison of major depression diagnostic classification probability using the SCID, CIDI, and MINI diagnostic interviews among women in pregnancy or postpartum: An individual participant data metaâ€analysis. International Journal of Methods in Psychiatric Research, 2019, 28, e1803.	1.1	34
24	Docosahexaenoic Acid for Selective Prevention of Posttraumatic Stress Disorder Among Severely Injured Patients. Journal of Clinical Psychiatry, 2015, 76, e1015-e1022.	1.1	34
25	The impact of the COVID-19 pandemic on the mental health of healthcare workers: study protocol for the COVID-19 HEalth caRe wOrkErS (HEROES) study. Social Psychiatry and Psychiatric Epidemiology, 2022, 57, 633-645.	1.6	33
26	Association between regular physical exercise and depressive symptoms mediated through social support and resilience in Japanese company workers: a cross-sectional study. BMC Public Health, 2016, 16, 553.	1.2	32
27	Association between working hours, work engagement, and work productivity in employees: A crossâ€sectional study of the Japanese Study of Health, Occupation, and Psychosocial Factors Relates Equity. Journal of Occupational Health, 2019, 61, 182-188.	1.0	32
28	The Tachikawa cohort of motor vehicle accident study investigating psychological distress: design, methods and cohort profiles. Social Psychiatry and Psychiatric Epidemiology, 2009, 44, 333-340.	1.6	30
29	Secondary traumatic stress and associated factors among Japanese nurses working in hospitals. International Journal of Nursing Practice, 2012, 18, 155-163.	0.8	30
30	Construct validity and test-retest reliability of the World Mental Health Japan version of the World Health Organization Health and Work Performance Questionnaire Short Version: a preliminary study. Industrial Health, 2020, 58, 375-387.	0.4	30
31	Serum pro-BDNF/BDNF as a treatment biomarker for response to docosahexaenoic acid in traumatized people vulnerable to developing psychological distress: a randomized controlled trial. Translational Psychiatry, 2015, 5, e596-e596.	2.4	29
32	Psychological growth after childbirth: an exploratory prospective study. Journal of Psychosomatic Obstetrics and Gynaecology, 2017, 38, 87-93.	1.1	28
33	Effects of omega-3 polyunsaturated fatty acids on psychophysiological symptoms of post-traumatic stress disorder in accident survivors: A randomized, double-blind, placebo-controlled trial. Journal of Affective Disorders, 2017, 224, 27-31.	2.0	27
34	Fish consumption and resilience to depression in Japanese company workers: a cross-sectional study. Lipids in Health and Disease, 2015, 14, 51.	1.2	26
35	Glutamatergic system abnormalities in posttraumatic stress disorder. Psychopharmacology, 2015, 232, 4261-4268.	1.5	26
36	Antenatal psychological intervention for universal prevention of antenatal and postnatal depression: A systematic review and meta-analysis. Journal of Affective Disorders, 2020, 273, 231-239.	2.0	26

#	Article	IF	Citations
37	Serum Oxytocin, Posttraumatic Coping and C-Reactive Protein in Motor Vehicle Accident Survivors by Gender. Neuropsychobiology, 2015, 71, 196-201.	0.9	24
38	The effect of social restrictions, loss of social support, and loss of maternal autonomy on postpartum depression in 1 to 12-months postpartum women during the COVID-19 pandemic. Journal of Affective Disorders, 2022, 307, 206-214.	2.0	23
39	Limited effect of omega-3 fatty acids on the quality of life in survivors of traumatic injury: A randomized, placebo-controlled trial. Prostaglandins Leukotrienes and Essential Fatty Acids, 2017, 127, 1-5.	1.0	22
40	Personal lifestyle as a resource for work engagement. Journal of Occupational Health, 2017, 59, 17-23.	1.0	22
41	Plasma estradiol levels and antidepressant effects of omega-3 fatty acids in pregnant women. Brain, Behavior, and Immunity, 2020, 85, 29-34.	2.0	22
42	Perinatal mental health and <scp>COVID</scp> â€19 in Japan. Psychiatry and Clinical Neurosciences, 2020, 74, 502-503.	1.0	22
43	Attenuating posttraumatic distress with omega-3 polyunsaturated fatty acids among disaster medical assistance team members after the Great East Japan Earthquake: The APOP randomized controlled trial. BMC Psychiatry, 2011, 11, 132.	1.1	19
44	Care difficulties and burden during COVIDâ€19 pandemic lockdowns among caregivers of people with schizophrenia: A crossâ€sectional study. Neuropsychopharmacology Reports, 2021, 41, 242-247.	1.1	19
45	Potential impact of propofol immediately after motor vehicle accident on later symptoms of posttraumatic stress disorder at 6-month follow up: a retrospective cohort study. Critical Care, 2012, 16, R196.	2.5	18
46	The longitudinal mental health impact of Fukushima nuclear disaster exposures and public criticism among power plant workers: the Fukushima NEWS Project study. Psychological Medicine, 2016, 46, 3117-3125.	2.7	18
47	The Efficacy of Omega-3 Fatty Acids for Depressive Symptoms among Pregnant Women in Japan and Taiwan: A Randomized, Double-Blind, Placebo-Controlled Trial (SYNCHRO; NCT01948596). Psychotherapy and Psychosomatics, 2019, 88, 122-124.	4.0	18
48	Postâ€traumatic stress symptoms among medical rescue workers exposed to <scp>COVID</scp> â€19 in Japan. Psychiatry and Clinical Neurosciences, 2020, 74, 503-505.	1.0	18
49	Prevalence and related factors of common mental disorders during pregnancy in Japan: a cross-sectional study. BioPsychoSocial Medicine, 2016, 10, 17.	0.9	17
50	Incidence and Prediction of Post-Traumatic Stress Disorder at 6 Months After Motor Vehicle Accidents in Japan. Psychosomatics, 2013, 54, 263-271.	2.5	16
51	Internet-based cognitive–behavioural therapy for prevention of depression during pregnancy and in the post partum (iPDP): a protocol for a large-scale randomised controlled trial. BMJ Open, 2020, 10, e036482.	0.8	16
52	Patterns and correlates of patientâ€reported helpfulness of treatment for common mental and substance use disorders in the <scp>WHO</scp> World Mental Health Surveys. World Psychiatry, 2022, 21, 272-286.	4.8	16
53	Serum Levels of Polyunsaturated Fatty Acids and the Risk of Posttraumatic Stress Disorder. Psychotherapy and Psychosomatics, 2013, 82, 408-410.	4.0	15
54	Exposure to psychotropic medications prior to overdose: a case-control study. Psychopharmacology, 2015, 232, 3101-3109.	1.5	15

#	Article	IF	Citations
55	Risk of recurrent overdose associated with prescribing patterns of psychotropic medications after nonfatal overdose. Neuropsychiatric Disease and Treatment, 2017, Volume 13, 653-665.	1.0	15
56	Secondary trauma and posttraumatic growth among mental health clinicians involved in disaster relief activities following the 2011 Tohoku earthquake and tsunami in Japan. Counselling Psychology Quarterly, 2020, 33, 427-447.	1.5	15
57	Psychological distress with and without a history of depression: Results from the World Mental Health Japan 2nd Survey (WMHJ2). Journal of Affective Disorders, 2020, 265, 545-551.	2.0	15
58	Association of response rate and prevalence estimates of common mental disorders across 129 areas in a nationally representative survey of adults in Japan. Social Psychiatry and Psychiatric Epidemiology, 2020, 55, 1373-1382.	1.6	15
59	Implementation Outcome Scales for Digital Mental Health (iOSDMH): Scale Development and Cross-sectional Study. JMIR Formative Research, 2021, 5, e24332.	0.7	15
60	Tachikawa project for prevention of posttraumatic stress disorder with polyunsaturated fatty acid (TPOP): study protocol for a randomized controlled trial. BMC Psychiatry, 2013, 13, 8.	1.1	14
61	Association between personal values in adolescence and mental health and well-being in adulthood: a cross-cultural study of working populations in Japan and the United States. Annals of General Psychiatry, 2020, 19, 7.	1.2	14
62	Post-Traumatic Stress Symptoms and Burnout Among Medical Rescue Workers 4 Years After the Great East Japan Earthquake: A Longitudinal Study. Disaster Medicine and Public Health Preparedness, 2016, 10, 848-853.	0.7	14
63	Mental health service requirements in a Japanese medical centre emergency department. Emergency Medicine Journal, 2006, 23, 468-469.	0.4	13
64	Are patients after severe injury who drop out of a longitudinal study at high risk of mental disorder?. Comprehensive Psychiatry, 2008, 49, 393-398.	1.5	13
65	Omegaâ€3 fatty acid supplementation for expectant mothers with depressive symptoms in Japan and Taiwan: An openâ€label trial. Psychiatry and Clinical Neurosciences, 2016, 70, 253-254.	1.0	13
66	Change in blood levels of eicosapentaenoic acid and posttraumatic stress symptom: A secondary analysis of data from a placebo-controlled trial of omega3 supplements. Journal of Affective Disorders, 2016, 205, 289-291.	2.0	13
67	Personal values in adolescence and suicidality: a cross-sectional study based on a retrospective recall. BMC Psychiatry, 2019, 19, 214.	1.1	13
68	Serum neuropeptide Y in accident survivors with depression or posttraumatic stress disorder. Neuroscience Research, 2014, 83, 8-12.	1.0	12
69	Personal values in adolescence and psychological distress in adults: A cross-sectional study based on a retrospective recall. PLoS ONE, 2019, 14, e0225454.	1.1	12
70	Association between energy drink consumption and substance use in adolescence: A systematic review of prospective cohort studies. Drug and Alcohol Dependence, 2021, 219, 108470.	1.6	12
71	Temporary employment and suicidal ideation in COVID-19 pandemic in Japan: A cross-sectional nationwide survey. Journal of Occupational Health, 2022, 64, e12319.	1.0	12
72	Cross-cultural validity of the WHO-5 Well-Being Index and Euthymia Scale: A clinimetric analysis. Journal of Affective Disorders, 2022, 311, 276-283.	2.0	12

#	Article	IF	CITATIONS
73	Spatial analysis for regional behavior of patients with mental disorders in Japan. Psychiatry and Clinical Neurosciences, 2017, 71, 254-261.	1.0	11
74	Japanese version of the 42-item psychological well-being scale (PWBS-42): a validation study. BMC Psychology, 2020, 8, 75.	0.9	11
75	Psychoâ€educational interventions focused on maternal or infant sleep for pregnant women to prevent the onset of antenatal and postnatal depression: A systematic review. Neuropsychopharmacology Reports, 2021, 41, 2-13.	1.1	11
76	The synchronized trial on expectant mothers with depressive symptoms by omega-3 PUFAs (SYNCHRO): Study protocol for a randomized controlled trial. BMC Psychiatry, 2016, 16, 321.	1.1	10
77	Urbanization and Internet addiction in a nationally representative sample of adult community residents in Japan: A cross-sectional, multilevel study. Psychiatry Research, 2019, 273, 699-705.	1.7	10
78	Perceived helpfulness of treatment for generalized anxiety disorder: a World Mental Health Surveys report. BMC Psychiatry, 2021, 21, 392.	1.1	10
79	Interactions Between Lithium and Non-Steroidal Antiinflammatory Drugs. Clinical Neuropharmacology, 2002, 25, 241-242.	0.2	9
80	Towards an Explanation of Inconsistent Rates of Posttraumatic Stress Disorder across Different Countries: Infant Mortality Rate as a Marker of Social Circumstances and Basic Population Health. Psychotherapy and Psychosomatics, 2010, 79, 56-57.	4.0	9
81	Hypnosis for induction of labour. The Cochrane Library, 2014, , CD010852.	1.5	9
82	Implementation of psychiatricâ€focused lifestyle medicine programs in <scp>A</scp> sia. Asia-Pacific Psychiatry, 2015, 7, 345-354.	1.2	9
83	Association between frequency of fried food consumption and resilience to depression in Japanese company workers: a cross-sectional study. Lipids in Health and Disease, 2016, 15, 156.	1.2	9
84	The association of personal importance of religion and religious service attendance with suicidal ideation by age group in the National Survey on Drug Use and Health. Psychiatry Research, 2017, 255, 321-327.	1.7	9
85	Impact of psychiatric morbidity on quality of life after motor vehicle accident at 1â€month follow up. Psychiatry and Clinical Neurosciences, 2009, 63, 235-237.	1.0	8
86	Psychological intervention for universal prevention of antenatal and postnatal depression among pregnant women: protocol for a systematic review and meta-analysis. Systematic Reviews, 2019, 8, 297.	2.5	8
87	The role of high-density lipoprotein cholesterol in risk for posttraumatic stress disorder: Taking a nutritional approach toward universal prevention. European Psychiatry, 2014, 29, 408-413.	0.1	7
88	Aromatic Selenoic, Selenothioic, and Diselenoic Acid Salts: Isolation, Characterization, and 77Se NMR Spectra, Together with Theoretical Elucidation. Bulletin of the Chemical Society of Japan, 2014, 87, 677-692.	2.0	7
89	Psychotic Experiences and Hikikomori in a Nationally Representative Sample of Adult Community Residents in Japan: A Cross-Sectional Study. Frontiers in Psychiatry, 2020, 11, 602678.	1.3	7
90	Longitudinal Change of Psychological Distress among Healthcare Professionals with and without Psychological First Aid Training Experience during the COVID-19 Pandemic. International Journal of Environmental Research and Public Health, 2021, 18, 12474.	1.2	7

#	Article	lF	Citations
91	Perceived helpfulness of treatment for social anxiety disorder: findings from the WHO World Mental Health Surveys. Social Psychiatry and Psychiatric Epidemiology, 2022, 57, 2079-2095.	1.6	7
92	The Effect of Omega-3 Fatty Acids on Psychophysiological Assessment for the Secondary Prevention of Posttraumatic Stress Disorder: An Open-Label Pilot Study. Global Journal of Health Science, 2011, 4, 3-9.	0.1	6
93	Serum polyunsaturated fatty acids and risk of psychiatric disorder after acute coronary syndrome: A prospective cohort study. Journal of Affective Disorders, 2017, 218, 306-312.	2.0	6
94	Psychological distress among people in Fukushima prefecture before and after the Great East Japan Earthquake using a nationâ€wide survey. Psychiatry and Clinical Neurosciences, 2018, 72, 878-878.	1.0	6
95	Effects of brief family psychoeducation for caregivers of people with schizophrenia in Japan provided by visiting nurses: protocol for a cluster randomised controlled trial. BMJ Open, 2020, 10, e034425.	0.8	6
96	Internet-based behavioural activation to improve depressive symptoms and prevent child abuse in postnatal women (SmartMama): a protocol for a pragmatic randomized controlled trial. BMC Pregnancy and Childbirth, 2021, 21, 314.	0.9	6
97	Sensitivity and concurrent validity of the Japanese version of the Euthymia scale: a clinimetric analysis. BMC Psychiatry, 2021, 21, 482.	1.1	6
98	Longitudinal change in depressive symptoms among healthcare professionals with and without COVID-19 vaccine hesitancy from October 2020 to June 2021 in Japan. Industrial Health, 2021, 60, 387-394.	0.4	5
99	Differences between Japan and Taiwan in the treatment of pregnant women with depressive symptoms by omega-3 fatty acids: An open-label pilot study. Nutritional Neuroscience, 2019, 22, 63-71.	1.5	4
100	Association between personal values in adolescence and impaired bonding relationship with children. BMC Psychology, 2020, 8, 98.	0.9	4
101	Time preference and personal value: a population-based cross-sectional study in Japan. BMC Psychology, 2020, 8, 85.	0.9	4
102	Internet-based acceptance and commitment therapy programme †Happiness Mom†for well-being: a protocol for a randomised controlled trial. BMJ Open, 2021, 11, e042167.	0.8	4
103	Associations Between Personal Values and Work Engagement. Journal of Occupational and Environmental Medicine, 2021, 63, e335-e340.	0.9	4
104	Personal values in adolescence and sense of coherence in adulthood: A crossâ€sectional study based on a retrospective recall. Neuropsychopharmacology Reports, 2020, 40, 262-267.	1.1	4
105	Peritraumatic distress after an earthquake: a bridge between neuroimaging and epidemiology. Molecular Psychiatry, 2013, 18, 743-744.	4.1	3
106	Mental Health Difficulties and Countermeasures during the Coronavirus Disease Pandemic in Japan: A Nationwide Questionnaire Survey of Mental Health and Psychiatric Institutions. International Journal of Environmental Research and Public Health, 2021, 18, 7318.	1.2	3
107	Previous disorders and depression outcomes in individuals with 12-month major depressive disorder in the World Mental Health surveys. Epidemiology and Psychiatric Sciences, 2021, 30, e70.	1.8	3
108	Perceived helpfulness of service sectors used for mental and substance use disorders: Findings from the WHO World Mental Health Surveys. International Journal of Mental Health Systems, 2022, 16, 6.	1.1	3

#	Article	IF	Citations
109	Uncovering heterogeneous associations of disasterâ€related traumatic experiences with subsequent mental health problems: A machine learning approach. Psychiatry and Clinical Neurosciences, 2022, 76, 97-105.	1.0	3
110	Approach to precise indoor characterization of multi-junction CPV cells using reference component cells. AIP Conference Proceedings, 2013 , , .	0.3	2
111	PTSD and the Attenuating Effects of Fish Oils: Results of Supplementation After the 2011 Great East Japan Earthquake. , 2013, , .		2
112	Generalizability of Findings from a Randomized Controlled Trial of Fish Oil Supplementation for Attenuating Posttraumatic Stress Symptoms among Rescue Workers in Japan. Psychotherapy and Psychosomatics, 2018, 87, 114-115.	4.0	2
113	Longitudinal association between adolescent work values and mental health and well-being in adulthood: a 23-year prospective cohort study. Scientific Reports, 2020, 10, 13547.	1.6	2
114	Adolescent Work Values and Drug Use in Adulthood: A Longitudinal Prospective Cohort Study. Substance Use and Misuse, 2021, 56, 1483-1492.	0.7	2
115	Association between Urban Upbringing and Compulsive Internet Use in Japan: A Cross-Sectional, Multilevel Study with Retrospective Recall. International Journal of Environmental Research and Public Health, 2021, 18, 9890.	1.2	2
116	The Effect of the Imacoco Care Psychoeducation Website on Improving Psychological Distress Among Workers During the COVID-19 Pandemic: Randomized Controlled Trial. JMIR Formative Research, 2022, 6, e33883.	0.7	2
117	Association between Work-Related Trauma Exposure and Posttraumatic Stress Symptoms among Child Welfare Workers in Japan: A Cross-Sectional Study. International Journal of Environmental Research and Public Health, 2021, 18, 3541.	1.2	1
118	Impact of peritraumatic distress on posttraumatic stress disorder symptoms at 6 months after acute coronary syndrome: a prospective cohort study. Högre Utbildning, 2021, 12, 1854511.	1.4	1
119	Erratum to "The Role of High-Density Lipoprotein Cholesterol in Risk For Posttraumatic Stress Disorder: Taking a Nutritional Approach Towards Universal Prevention―[Eur. Psych. 29 (2014) 408–413]. European Psychiatry, 2015, 30, 354-354.	0.1	0
120	Role of resilience for the association between trait hostility and depressive symptoms in Japanese company workers. Current Psychology, 2021, 40, 2301-2308.	1.7	0
121	Preconception menstrual cycle disorder and antenatal depression: a cross-sectional study with prerecorded information. Journal of Psychosomatic Obstetrics and Gynaecology, 2021, , 1-8.	1.1	0
122	Title is missing!. , 2019, 14, e0225454.		0
123	Title is missing!. , 2019, 14, e0225454.		0
124	Title is missing!. , 2019, 14, e0225454.		0
125	Title is missing!. , 2019, 14, e0225454.		0