

# Bang-Gee Hsu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2090602/publications.pdf>

Version: 2024-02-01

178  
papers

1,711  
citations

361045

20  
h-index

476904

29  
g-index

179  
all docs

179  
docs citations

179  
times ranked

1895  
citing authors

#	ARTICLE	IF	CITATIONS
1	Inverse association of serum osteocalcin and bone mineral density in renal transplant recipients. Tzu Chi Medical Journal, 2023, 35, 165.	0.4	1
2	Serum myostatin level is a positive predictor of endothelial function measured by digital thermal monitoring of vascular reactivity in kidney transplantation patients. Tzu Chi Medical Journal, 2022, 34, 62.	0.4	1
3	Positive correlation of serum angiotensin-like protein 3 levels with metabolic syndrome in patients with coronary artery disease. Tzu Chi Medical Journal, 2022, 34, 75.	0.4	2
4	A Retrospective Cohort Study of the Association between Serum Osteopontin Levels and Aortic Stiffness in Hypertensive Patients. International Journal of Environmental Research and Public Health, 2022, 19, 477.	1.2	1
5	The Correlation of Serum Myostatin Levels with Gait Speed in Kidney Transplantation Recipients. International Journal of Environmental Research and Public Health, 2022, 19, 465.	1.2	1
6	Serum Fibroblast Growth Factor 21 Level Is Associated with Aortic Stiffness in Patients on Maintenance Hemodialysis. International Journal of Hypertension, 2022, 2022, 1-8.	0.5	2
7	Dapagliflozin Ameliorates Lipopolysaccharide Related Acute Kidney Injury in Mice with Streptozotocin-induced Diabetes Mellitus. International Journal of Medical Sciences, 2022, 19, 729-739.	1.1	13
8	A Comparison of SARC-F, Calf Circumference, and Their Combination for Sarcopenia Screening among Patients Undergoing Peritoneal Dialysis. Nutrients, 2022, 14, 923.	1.7	10
9	A Novel Application of Serum Creatinine and Cystatin C to Predict Sarcopenia in Advanced CKD. Frontiers in Nutrition, 2022, 9, 828880.	1.6	7
10	Association between Serum Adipocyte Fatty Acid Binding Protein Level and Endothelial Dysfunction in Chronic Hemodialysis Patients. Life, 2022, 12, 316.	1.1	3
11	Age and Serum Adipocyte Fatty-Acid-Binding Protein Level Are Associated with Aortic Stiffness in Coronary Artery Bypass Graft Patients. Journal of Cardiovascular Development and Disease, 2022, 9, 105.	0.8	1
12	Serum Osteopontin Level Is Positively Associated with Aortic Stiffness in Patients with Peritoneal Dialysis. Life, 2022, 12, 397.	1.1	3
13	Negative Correlation of Serum Adiponectin Level with Aortic Stiffness in Elderly Diabetic Persons. International Journal of Environmental Research and Public Health, 2022, 19, 3062.	1.2	1
14	Serum Sclerostin Level Is Negatively Associated with Bone Mineral Density in Hemodialysis Patients. Medicina (Lithuania), 2022, 58, 385.	0.8	4
15	Assessment of uremic sarcopenia in dialysis patients: An update. Tzu Chi Medical Journal, 2022, 34, 182.	0.4	1
16	Serum P-Cresyl Sulfate Level Is an Independent Marker of Peripheral Arterial Stiffness as Assessed Using Brachial-Ankle Pulse Wave Velocity in Patients with Non-Dialysis Chronic Kidney Disease Stage 3 to 5. Toxins, 2022, 14, 287.	1.5	0
17	Correlation between Serum 25-Hydroxyvitamin D Level and Peripheral Arterial Stiffness in Chronic Kidney Disease Stage 3-5 Patients. Nutrients, 2022, 14, 2429.	1.7	5
18	Serum Trimethylamine N-Oxide Levels Correlate with Metabolic Syndrome in Coronary Artery Disease Patients. International Journal of Environmental Research and Public Health, 2022, 19, 8710.	1.2	3

#	ARTICLE	IF	CITATIONS
19	Urine phthalate metabolites are associated with urothelial cancer in chronic kidney disease patients. <i>Chemosphere</i> , 2021, 273, 127834.	4.2	6
20	Arterial stiffness: A brief review. <i>Tzu Chi Medical Journal</i> , 2021, 33, 115.	0.4	19
21	Association between serum indoxyl sulfate levels with carotid-femoral pulse wave velocity in patients with chronic kidney disease. <i>Renal Failure</i> , 2021, 43, 796-802.	0.8	4
22	Vascular calcification of chronic kidney disease: A brief review. <i>Tzu Chi Medical Journal</i> , 2021, 33, 34.	0.4	6
23	Positive Associations between Adipocyte Fatty Acid-Binding Protein Level and Central Arterial Stiffness in Peritoneal Dialysis Patients. <i>International Journal of Hypertension</i> , 2021, 2021, 1-8.	0.5	5
24	Serum Levels of Fibroblast Growth Factor 21 Are Positively Associated with Aortic Stiffness in Patients with Type 2 Diabetes Mellitus. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 3434.	1.2	3
25	Discovery of Novel Protein Biomarkers in Urine for Diagnosis of Urothelial Cancer Using iTRAQ Proteomics. <i>Journal of Proteome Research</i> , 2021, 20, 2953-2963.	1.8	18
26	Hypoadiponectinemia is associated with aortic stiffness in nondialysis diabetic patients with stage 3-5 chronic kidney disease. <i>Vascular</i> , 2021, , 170853812110076.	0.4	2
27	Serum Osteoprotegerin Level Is Negatively Associated with Bone Mineral Density in Patients Undergoing Maintenance Hemodialysis. <i>Medicina (Lithuania)</i> , 2021, 57, 762.	0.8	1
28	Serum Galectin-3 Level Is Positively Associated with Endothelial Dysfunction in Patients with Chronic Kidney Disease Stage 3 to 5. <i>Toxins</i> , 2021, 13, 532.	1.5	8
29	Positive correlation of serum indoxyl sulfate level with peripheral arterial disease in hemodialysis patients. <i>Vascular</i> , 2021, , 170853812110399.	0.4	0
30	Serum indices based on creatinine and cystatin C predict mortality in patients with non-dialysis chronic kidney disease. <i>Scientific Reports</i> , 2021, 11, 16863.	1.6	14
31	Serum Angiotensin-like Protein 3 Level Is Associated with Peripheral Arterial Stiffness in Patients with Coronary Artery Disease. <i>Medicina (Lithuania)</i> , 2021, 57, 1011.	0.8	4
32	The MicroRNA Prediction Models as Ancillary Diagnosis Biomarkers for Urothelial Carcinoma in Patients With Chronic Kidney Disease. <i>Frontiers in Medicine</i> , 2021, 8, 726214.	1.2	1
33	Effects of ketoanalogues on skeletal muscle mass in patients with advanced chronic kidney disease: real-world evidence. <i>Nutrition</i> , 2021, 91-92, 111384.	1.1	2
34	Association of serum osteopontin with first hospitalization and all-cause mortality after kidney transplantation. <i>Tzu Chi Medical Journal</i> , 2021, 34, 200-206.	0.4	0
35	Serum Irisin Level Is Positively Associated with Bone Mineral Density in Patients on Maintenance Hemodialysis. <i>International Journal of Endocrinology</i> , 2021, 2021, 1-6.	0.6	8
36	Linagliptin Protects against Endotoxin-Induced Acute Kidney Injury in Rats by Decreasing Inflammatory Cytokines and Reactive Oxygen Species. <i>International Journal of Molecular Sciences</i> , 2021, 22, 11190.	1.8	13

#	ARTICLE	IF	CITATIONS
37	Low serum 3-methyl histidine level is associated with aortic stiffness in maintenance hemodialysis patients. <i>Therapeutic Apheresis and Dialysis</i> , 2021, , .	0.4	1
38	Positive Association of Serum Adiponectin Level With the Vascular Reactivity Index in Kidney Transplant Patients. <i>Experimental and Clinical Transplantation</i> , 2021, , .	0.2	0
39	Association of endothelial dysfunction with sarcopenia and muscle function in a relatively young cohort of kidney transplant recipients. <i>PeerJ</i> , 2021, 9, e12521.	0.9	7
40	Positive Correlation of Serum Resistin Level with Peripheral Artery Disease in Patients with Chronic Kidney Disease Stage 3 to 5. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 12746.	1.2	2
41	Positive Association of Serum Adipocyte Fatty Acid Binding Protein Level With Peripheral Artery Disease in Hemodialysis Patients. <i>Therapeutic Apheresis and Dialysis</i> , 2020, 24, 300-306.	0.4	3
42	Serum creatinine to cystatin C ratio predicts skeletal muscle mass and strength in patients with non-dialysis chronic kidney disease. <i>Clinical Nutrition</i> , 2020, 39, 2435-2441.	2.3	67
43	Low serum leptin levels are associated with malnutrition status according to malnutrition-inflammation score in patients undergoing chronic hemodialysis. <i>Hemodialysis International</i> , 2020, 24, 221-227.	0.4	10
44	Impact of sarcopenia and its diagnostic criteria on hospitalization and mortality in chronic hemodialysis patients: A 3-year longitudinal study. <i>Journal of the Formosan Medical Association</i> , 2020, 119, 1219-1229.	0.8	29
45	Serum P-Cresyl Sulfate Is a Predictor of Central Arterial Stiffness in Patients on Maintenance Hemodialysis. <i>Toxins</i> , 2020, 12, 10.	1.5	20
46	Association of Low Serum L-Carnitine Levels with Aortic Stiffness in Patients with Non-Dialysis Chronic Kidney Disease. <i>Nutrients</i> , 2020, 12, 2918.	1.7	2
47	Low Serum 3-Methylhistidine Levels Are Associated With First Hospitalization in Kidney Transplantation Recipients. <i>Transplantation Proceedings</i> , 2020, 52, 3214-3220.	0.3	3
48	Association of Serum Indoxyl Sulfate Levels with Skeletal Muscle Mass and Strength in Chronic Hemodialysis Patients: A 2-year Longitudinal Analysis. <i>Calcified Tissue International</i> , 2020, 107, 257-265.	1.5	9
49	Serum Malondialdehyde-Modified Low-Density Lipoprotein Is a Risk Factor for Central Arterial Stiffness in Maintenance Hemodialysis Patients. <i>Nutrients</i> , 2020, 12, 2160.	1.7	11
50	P0785 POSITIVE CORRELATION OF SERUM RESISTIN LEVEL WITH PERIPHERAL ARTERY OCCLUSIVE DISEASE IN PATIENTS WITH CHRONIC KIDNEY DISEASE STAGE 3 TO 5. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.4	0
51	Serum indoxyl sulfate as a potential biomarker of aortic arterial stiffness in coronary artery disease. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020, 30, 2320-2327.	1.1	7
52	Serum Osteocalcin Level is Negatively Associated with Vascular Reactivity Index by Digital Thermal Monitoring in Kidney Transplant Recipients. <i>Medicina (Lithuania)</i> , 2020, 56, 400.	0.8	5
53	P0768 NOVEL URINARY PROTEIN BIOMARKERS FOR UROTHELIAL CANCER IN PATIENTS WITH CHRONIC KIDNEY DISEASE. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.4	0
54	Associations between High Serum Adipocyte Fatty Acid Binding Protein and First Hospitalization in Kidney Transplantation Patients: A 5-Year Follow-up Study. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 7567.	1.2	1

#	ARTICLE	IF	CITATIONS
55	Association of SARC-F Questionnaire and Mortality in Prevalent Hemodialysis Patients. <i>Diagnostics</i> , 2020, 10, 890.	1.3	9
56	Serum leptin levels are positively associated with aortic stiffness in patients with chronic kidney disease stage 3-5. <i>Adipocyte</i> , 2020, 9, 206-211.	1.3	7
57	Increased Serum Leptin Level Predicts Bone Mineral Density in Hemodialysis Patients. <i>International Journal of Endocrinology</i> , 2020, 2020, 1-7.	0.6	4
58	Lower Serum Fibroblast Growth Factor 21 Levels are Associated with Normal Lumbar Spine Bone Mineral Density in Hemodialysis Patients. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 1938.	1.2	5
59	Serum Sclerostin But Not DKK-1 Correlated with Central Arterial Stiffness in End Stage Renal Disease Patients. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 1230.	1.2	9
60	Positive correlation of serum fibroblast growth factor 23 with peripheral arterial stiffness in kidney transplantation patients. <i>Clinica Chimica Acta</i> , 2020, 505, 9-14.	0.5	4
61	Serum osteoprotegerin level is positively associated with peripheral artery disease in patients with peritoneal dialysis. <i>Renal Failure</i> , 2020, 42, 131-136.	0.8	8
62	High-Serum Angiotensin-Like Protein 3 Levels Associated with Cardiovascular Outcome in Patients with Coronary Artery Disease. <i>International Journal of Hypertension</i> , 2020, 2020, 1-7.	0.5	10
63	Negative correlation of serum adiponectin level with peripheral artery occlusive disease in hemodialysis patients. <i>Tzu Chi Medical Journal</i> , 2020, 32, 70.	0.4	2
64	High Serum Fibroblast Growth Factor 23 Level Is Associated With Metabolic Syndrome in Kidney Transplantation Patients. <i>Transplantation Proceedings</i> , 2020, 52, 3168-3172.	0.3	3
65	Low serum adiponectin level is associated with central arterial stiffness in patients undergoing peritoneal dialysis. <i>Tzu Chi Medical Journal</i> , 2020, 32, 272.	0.4	0
66	Association of Low Serum Adiponectin Levels with Aortic Arterial Stiffness in Patients with Type 2 Diabetes. <i>Journal of Clinical Medicine</i> , 2019, 8, 887.	1.0	11
67	Out-of-pocket costs and productivity losses in haemodialysis and peritoneal dialysis from a patient interview survey in Taiwan. <i>BMJ Open</i> , 2019, 9, e023062.	0.8	13
68	Association between Serum Indoxyl Sulfate Levels and Endothelial Function in Non-Dialysis Chronic Kidney Disease. <i>Toxins</i> , 2019, 11, 589.	1.5	21
69	SP084Hypoalbuminemia is positively associated with peripheral arterial stiffness in patients with chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.4	0
70	SP124Serum leptin level is positively associated with aortic arterial stiffness in patients with chronic kidney disease stage 3 to 5. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.4	0
71	SP737Serum osteocalcin level is negatively associated with vascular reactivity index by digital thermal monitoring in kidney transplant patient. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.4	0
72	FP637Serum p-cresyl sulfate level is positively associated with central arterial stiffness in patients on hemodialysis. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.4	0

#	ARTICLE	IF	CITATIONS
73	FP724Serum fibroblast growth factor 21 levels are positively associated with skeletal muscle index in chronic hemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.4	0
74	Acute Kidney Injury and Septic Shockâ€”Defined by Updated Sepsis-3 Criteria in Critically Ill Patients. <i>Journal of Clinical Medicine</i> , 2019, 8, 1731.	1.0	6
75	SP134Elevated serum leptin levels are associated with low muscle strength and muscle quality in chronic hemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.4	0
76	&lt;p&gt;Metabolic Syndrome With Aortic Arterial Stiffness And First Hospitalization Or Mortality In Coronary Artery Disease Patients&lt;/p&gt;. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2019, Volume 12, 2065-2073.	1.1	7
77	Association of Low Serum L-Carnitine Levels with Peripheral Arterial Stiffness in Patients Who Undergo Kidney Transplantation. <i>Nutrients</i> , 2019, 11, 2000.	1.7	9
78	Serum Adipocyte Fatty Acid-Binding Protein Level is Negatively Associated with Vascular Reactivity Index Measured by Digital Thermal Monitoring in Kidney Transplant Patients. <i>Metabolites</i> , 2019, 9, 159.	1.3	8
79	SP748Inverse association of serum osteocalcin and bone mineral density in renal transplant recipients. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.4	0
80	Serum Fibroblast Growth Factor 21 Levels Are Positively Associated with Metabolic Syndrome in Patients with Type 2 Diabetes. <i>International Journal of Endocrinology</i> , 2019, 2019, 1-8.	0.6	25
81	Serum osteoprotegerin is an independent marker of central arterial stiffness as assessed using carotidâ€”femoral pulse wave velocity in hemodialysis patients: a cross sectional study. <i>BMC Nephrology</i> , 2019, 20, 184.	0.8	9
82	Development and validation of a nomogram for urothelial cancer in patients with chronic kidney disease. <i>Scientific Reports</i> , 2019, 9, 3473.	1.6	4
83	High serum adipocyte fatty acid binding protein concentration linked with increased aortic arterial stiffness in patients with type 2 diabetes. <i>Clinica Chimica Acta</i> , 2019, 495, 35-39.	0.5	12
84	Angiotensin II receptor blockade is associated with preserved muscle strength in chronic hemodialysis patients. <i>BMC Nephrology</i> , 2019, 20, 54.	0.8	19
85	Correlation between sclerostin and Dickkopf-1 with aortic arterial stiffness in patients with type 2 diabetes: A prospective, cross-sectional study. <i>Diabetes and Vascular Disease Research</i> , 2019, 16, 281-288.	0.9	8
86	Osteocalcin and carotidâ€”femoral pulse wave velocity in patients on peritoneal dialysis. <i>Tzu Chi Medical Journal</i> , 2019, 31, 23.	0.4	6
87	Reply from authors for Serum leptin levels and peripheral arterial stiffness in patients with type 2 diabetes. <i>Tzu Chi Medical Journal</i> , 2019, 31, 295.	0.4	0
88	The serum sclerostin level is positively associated with the aortic augmentation index in patients on peritoneal dialysis. <i>Tzu Chi Medical Journal</i> , 2019, 31, 102.	0.4	1
89	Negative Correlation of Serum Adiponectin Levels With Carotidâ€”Femoral Pulse Wave Velocity in Patients Treated With Hemodialysis. <i>Biological Research for Nursing</i> , 2018, 20, 462-468.	1.0	6
90	Serum cystatin C is independently associated with aortic arterial stiffness in patients with type 2 diabetes. <i>Clinica Chimica Acta</i> , 2018, 480, 114-118.	0.5	9

#	ARTICLE	IF	CITATIONS
91	A positive association between interleukin-1 receptor antagonist and insulin resistance in postmenopausal women. <i>Gynecological Endocrinology</i> , 2018, 34, 574-578.	0.7	0
92	Serum adipocyte fatty acid-binding protein level is associated with arterial stiffness quantified with cardio-ankle vascular index in kidney transplant patients. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 188-195.	0.7	7
93	FP627ASERUM SCLEROSTIN LEVEL IS POSITIVELY ASSOCIATED BONE MINERAL DENSITY IN HEMODIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i253-i253.	0.4	0
94	Association between hyperleptinemia and cardiovascular outcomes in patients with coronary artery disease. <i>Therapeutics and Clinical Risk Management</i> , 2018, Volume 14, 1855-1862.	0.9	15
95	FP112SERUM OSTEOPONTIN LEVEL IS INDEPENDENTLY ASSOCIATED WITH CENTRAL ARTERIAL STIFFNESS IN PATIENTS ON HEMODIALYSIS. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i15-i15.	0.4	0
96	FP121SERUM ADIPOCYTE FATTY ACID BINDING PROTEIN LEVEL IS POSITIVELY ASSOCIATED WITH AORTIC ARTERIAL STIFFNESS IN CHRONIC KIDNEY DISEASE PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i18-i18.	0.4	0
97	SP705LOW SERUM 3-METHYL HISTIDINE LEVEL IS ASSOCIATED WITH FIRST HOSPITALIZATION IN KIDNEY TRANSPLANTATION PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i584-i584.	0.4	0
98	Serum levels of sclerostin as a potential biomarker in central arterial stiffness among hypertensive patients. <i>BMC Cardiovascular Disorders</i> , 2018, 18, 214.	0.7	14
99	High Levels of Serum Adipocyte Fatty Acid-binding Protein Predict Cardiovascular Events in Coronary Artery Disease Patients. <i>International Journal of Medical Sciences</i> , 2018, 15, 1268-1274.	1.1	8
100	Inverse Association of N-terminal Pro-B-type Natriuretic Peptide Level With Metabolic Syndrome in Kidney Transplant Patients. <i>Transplantation Proceedings</i> , 2018, 50, 2496-2501.	0.3	6
101	FP488POSITIVE CORRELATES OF SERUM SCLEROSTIN LEVEL AND ASSOCIATION WITH CENTRAL ARTERIAL STIFFNESS IN PATIENTS WITH PERITONEAL DIALYSIS. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i202-i202.	0.4	1
102	Positive correlation of the serum angiotensin-like protein 3 levels with the aortic augmentation index in patients with coronary artery disease. <i>Therapeutics and Clinical Risk Management</i> , 2018, Volume 14, 231-236.	0.9	10
103	Decreased serum fatty acid binding protein 4 concentrations are associated with sarcopenia in chronic hemodialysis patients. <i>Clinica Chimica Acta</i> , 2018, 485, 113-118.	0.5	16
104	Hyperleptinemia positively associated with central arterial stiffness in hemodialysis patients. <i>PLoS ONE</i> , 2018, 13, e0190694.	1.1	10
105	High serum leptin levels are associated with central arterial stiffness in geriatric patients on hemodialysis. <i>Tzu Chi Medical Journal</i> , 2018, 30, 227.	0.4	3
106	Positive correlation of serum leptin levels and peripheral arterial stiffness in patients with type 2 diabetes. <i>Tzu Chi Medical Journal</i> , 2018, 30, 10.	0.4	4
107	High serum leptin level is associated with peripheral artery disease in adult peritoneal dialysis patients. <i>Tzu Chi Medical Journal</i> , 2018, 30, 85.	0.4	4
108	Hyperleptinemia is associated with the aortic augmentation index in kidney transplant recipients. <i>Tzu Chi Medical Journal</i> , 2018, 30, 152.	0.4	2

#	ARTICLE	IF	CITATIONS
109	Negative correlation between leptin serum levels and sarcopenia in hemodialysis patients. <i>International Journal of Clinical and Experimental Pathology</i> , 2018, 11, 1715-1723.	0.5	3
110	Serum sclerostin as an independent marker of peripheral artery disease in elderly persons. <i>International Journal of Clinical and Experimental Pathology</i> , 2018, 11, 2816-2821.	0.5	3
111	Fasting serum resistin level positively correlates with metabolic syndrome in patients on peritoneal dialysis. <i>International Journal of Clinical and Experimental Pathology</i> , 2018, 11, 4257-4263.	0.5	0
112	Economic costs of automated and continuous ambulatory peritoneal dialysis in Taiwan: a combined survey and retrospective cohort analysis. <i>BMJ Open</i> , 2017, 7, e015067.	0.8	5
113	High serum adipocyte fatty acid binding protein level as a potential biomarker of aortic arterial stiffness in hypertensive patients with metabolic syndrome. <i>Clinica Chimica Acta</i> , 2017, 473, 166-172.	0.5	23
114	Low serum adiponectin level is associated with metabolic syndrome and is an independent marker of peripheral arterial stiffness in hypertensive patients. <i>Diabetology and Metabolic Syndrome</i> , 2017, 9, 49.	1.2	26
115	High serum resistin levels are associated with peripheral artery disease in the hypertensive patients. <i>BMC Cardiovascular Disorders</i> , 2017, 17, 80.	0.7	16
116	Triceps Skinfold Thickness Is Associated With Lumbar Bone Mineral Density in Peritoneal Dialysis Patients. <i>Therapeutic Apheresis and Dialysis</i> , 2017, 21, 102-107.	0.4	8
117	Serum leptin is a predictor for central arterial stiffness in hypertensive patients. <i>Nephrology</i> , 2017, 22, 783-789.	0.7	15
118	Acute Alcohol Intoxication Exacerbates Rhabdomyolysis-Induced Acute Renal Failure in Rats. <i>International Journal of Medical Sciences</i> , 2017, 14, 680-689.	1.1	21
119	Serum resistin as an independent marker of aortic stiffness in patients with coronary artery disease. <i>PLoS ONE</i> , 2017, 12, e0183123.	1.1	15
120	Leptin is an independent marker of metabolic syndrome in elderly adults with type 2 diabetes. <i>Tzu Chi Medical Journal</i> , 2017, 29, 109.	0.4	4
121	Association of hyperleptinemia with peripheral arterial disease in hypertensive patients. <i>Tzu Chi Medical Journal</i> , 2017, 29, 148.	0.4	3
122	Serum leptin level positively correlates with metabolic syndrome among elderly Taiwanese. <i>Tzu Chi Medical Journal</i> , 2017, 29, 159.	0.4	10
123	Elevated serum osteoprotegerin may predict peripheral arterial disease after kidney transplantation: a single-center prospective cross-sectional study in Taiwan. <i>PeerJ</i> , 2017, 5, e3847.	0.9	7
124	Positive correlation of serum adipocyte fatty acid binding protein levels with metabolic syndrome in kidney transplantation patients. <i>International Journal of Clinical and Experimental Pathology</i> , 2017, 10, 8727-8734.	0.5	4
125	Hypoadiponectemia is associated with metabolic syndrome in patients with type 2 diabetes. <i>International Journal of Clinical and Experimental Pathology</i> , 2017, 10, 10515-10521.	0.5	3
126	High Serum Adipocyte Fatty Acid Binding Protein Is Associated with Metabolic Syndrome in Patients with Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-7.	1.0	18



#	ARTICLE	IF	CITATIONS
127	Serum Sclerostin as an Independent Marker of Peripheral Arterial Stiffness in Renal Transplantation Recipients. <i>Medicine (United States)</i> , 2016, 95, e3300.	0.4	33
128	Serum adiponectin is a negative predictor of central arterial stiffness in kidney transplant patients. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2016, 76, 264-269.	0.6	8
129	Serum osteoprotegerin levels associated with the aortic augmentation index in renal transplant recipients. <i>Tzu Chi Medical Journal</i> , 2016, 28, 20-23.	0.4	2
130	Inverse relationship of bone mineral density and serum level of N-terminal pro-B-type natriuretic peptide in peritoneal dialysis patients. <i>Tzu Chi Medical Journal</i> , 2016, 28, 68-72.	0.4	5
131	Calcitriol decreases pro-inflammatory cytokines and protects against severe hemorrhagic shock induced-organ damage in rats. <i>Cytokine</i> , 2016, 83, 262-268.	1.4	8
132	Association of serum leptin levels with central arterial stiffness in coronary artery disease patients. <i>BMC Cardiovascular Disorders</i> , 2016, 16, 80.	0.7	45
133	Relationship between resistin and mortality in maintenance hemodialysis patients. <i>Clinical Nephrology</i> , 2016, 86, 125-131.	0.4	6
134	High Serum Osteoprotegerin Is Associated with Arterial Stiffness in Kidney Transplant Patients. <i>Tohoku Journal of Experimental Medicine</i> , 2015, 236, 247-253.	0.5	10
135	Serum Osteoprotegerin is Associated with Arterial Stiffness Assessed According to the Cardio-ankle Vascular Index in Hypertensive Patients. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015, 22, 304-312.	0.9	26
136	FP862SERUM OSTEOPROTEGERIN LEVELS ASSOCIATED WITH AORTIC AUGMENTATION INDEX IN RENAL TRANSPLANT RECIPIENTS. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, iii365-iii365.	0.4	0
137	Hypoadiponectinemia correlates with arterial stiffness in kidney transplantation patients. <i>Clinical and Experimental Nephrology</i> , 2015, 19, 534-541.	0.7	8
138	Positive correlation of serum adipocyte fatty acid binding protein levels with carotid-femoral pulse wave velocity in geriatric population. <i>BMC Geriatrics</i> , 2015, 15, 88.	1.1	22
139	Serum Osteopontin Level Correlates with Carotid-Femoral Pulse Wave Velocity in Geriatric Persons. <i>BioMed Research International</i> , 2014, 2014, 1-7.	0.9	18
140	Association of Serum Osteoprotegerin Levels With Carotid-Femoral Pulse Wave Velocity in Hypertensive Patients. <i>Journal of Clinical Hypertension</i> , 2014, 16, 301-308.	1.0	28
141	N-terminal pro-B-type natriuretic peptide level inversely associates with metabolic syndrome in elderly persons. <i>Diabetology and Metabolic Syndrome</i> , 2014, 6, 15.	1.2	5
142	Serum atrial natriuretic peptide level inversely associates with metabolic syndrome in older adults. <i>Geriatrics and Gerontology International</i> , 2014, 14, 640-646.	0.7	8
143	Calcitriol decreases TGF- $\beta$ 1 and angiotensin II production and protects against chlorhexide digluconate-induced liver peritoneal fibrosis in rats. <i>Cytokine</i> , 2014, 65, 105-118.	1.4	20
144	Inverse association of long-acting natriuretic peptide with metabolic syndrome in congestive heart failure patients. <i>Diabetology and Metabolic Syndrome</i> , 2013, 5, 19.	1.2	9

#	ARTICLE	IF	CITATIONS
145	Low Serum Long-acting Natriuretic Peptide Level Correlates with Metabolic Syndrome in Hypertensive Patients: A Cross-sectional Study. <i>Archives of Medical Research</i> , 2013, 44, 215-220.	1.5	13
146	Heavy Ethanol Intoxication Increases Proinflammatory Cytokines and Aggravates Hemorrhagic Shock-Induced Organ Damage in Rats. <i>Mediators of Inflammation</i> , 2013, 2013, 1-9.	1.4	15
147	N-terminal Pro-B-Type Natriuretic Peptide is Associated with Arterial Stiffness Measured Using the Cardio-Ankle Vascular Index in Renal Transplant Recipients. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013, 20, 646-653.	0.9	15
148	Inverse association of serum long-acting natriuretic peptide and bone mineral density in renal transplant recipients. <i>Clinical Transplantation</i> , 2012, 26, E105-10.	0.8	11
149	Valsartan decreases TGF- $\beta$ 21 production and protects against chlorhexidine digluconate-induced liver peritoneal fibrosis in rats. <i>Cytokine</i> , 2011, 53, 223-230.	1.4	29
150	Hyperleptinaemia positively correlated with metabolic syndrome in hemodialysis patients. <i>European Journal of Internal Medicine</i> , 2011, 22, e105-e109.	1.0	15
151	Melatonin Ameliorates Hemorrhagic Shock-Induced Organ Damage in Rats. <i>Journal of Surgical Research</i> , 2011, 167, e315-e321.	0.8	28
152	Rosiglitazone protects against severe hemorrhagic shock-induced organ damage in rats. <i>Medical Science Monitor</i> , 2011, 17, BR282-BR289.	0.5	6
153	Beneficial Effects of Enalapril on Chlorhexidine Digluconate-Induced Liver Peritoneal Fibrosis in Rats. <i>Chinese Journal of Physiology</i> , 2011, 54, 225-234.	0.4	9
154	Fasting Adiponectin is Inversely Correlated with Metabolic Syndrome in Patients with Coronary Artery Disease. <i>Internal Medicine</i> , 2010, 49, 739-747.	0.3	12
155	Fasting Serum Level of Fatty-Acid-Binding Protein 4 Positively Correlates With Metabolic Syndrome in Patients With Coronary Artery Disease. <i>Circulation Journal</i> , 2010, 74, 327-331.	0.7	53
156	Effect of Intravenous N-acetylcysteine on Plasma Total Homocysteine and Inflammatory Cytokines During High Flux Hemodialysis. <i>Tzu Chi Medical Journal</i> , 2010, 22, 90-95.	0.4	6
157	C-reactive Protein Positively Correlates With Metabolic Syndrome in Kidney Transplantation Patients. <i>Tzu Chi Medical Journal</i> , 2010, 22, 131-136.	0.4	0
158	Fasting Serum Fatty Acid-binding Protein 4 Level Positively Correlates with Metabolic Syndrome in Hemodialysis Patients. <i>Archives of Medical Research</i> , 2010, 41, 536-540.	1.5	15
159	Erythropoietin protects severe haemorrhagic shock-induced organ damage in conscious rats. <i>Injury</i> , 2010, 41, 724-730.	0.7	22
160	Fasting Serum Leptin Level Correlates With Mid-Arm Fat Area in Peritoneal Dialysis Patients. <i>Therapeutic Apheresis and Dialysis</i> , 2010, 14, 583-588.	0.4	10
161	Aliskiren ameliorates chlorhexidine digluconate-induced peritoneal fibrosis in rats. <i>European Journal of Clinical Investigation</i> , 2010, 40, 301-309.	1.7	20
162	Hyperleptinemia positively correlated with metabolic syndrome in renal transplant recipients. <i>Clinical Transplantation</i> , 2010, 24, E124-9.	0.8	11

#	ARTICLE	IF	CITATIONS
163	Methemoglobinemia induced by indoxacarb intoxication. <i>Clinical Toxicology</i> , 2010, 48, 766-767.	0.8	19
164	Stability of Blood Biochemistry Levels in Animal Model Research: Effects of Storage Condition and Time. <i>Biological Research for Nursing</i> , 2010, 11, 395-400.	1.0	8
165	Fasting Serum Adiponectin Level Inversely Correlates with Metabolic Syndrome in Peritoneal Dialysis Patients. <i>Blood Purification</i> , 2010, 30, 1-7.	0.9	12
166	The Association of Serum Osteoprotegerin and Osteoporosis in Postmenopausal Hemodialysis Patients: A Pilot Study. <i>Journal of Women's Health</i> , 2010, 19, 785-790.	1.5	8
167	Low-dose erythropoietin aggravates endotoxin-induced organ damage in conscious rats. <i>Cytokine</i> , 2010, 49, 155-162.	1.4	25
168	Fatal Outcome After Ingestion of Star Fruit in a Severe Chronic Kidney Disease Patient. <i>Dialysis and Transplantation</i> , 2009, 38, 102-105.	0.2	4
169	Fluvastatin attenuates severe hemorrhagic shock-induced organ damage in rats. <i>Resuscitation</i> , 2009, 80, 372-378.	1.3	36
170	LOW DOSE PROPOFOL AMELIORATES HAEMORRHAGIC SHOCK INDUCED ORGAN DAMAGE IN CONSCIOUS RATS. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2008, 35, 766-774.	0.9	16
171	Fasting Serum Total Ghrelin Level Inversely Correlates with Metabolic Syndrome in Hemodialysis Patients. <i>Archives of Medical Research</i> , 2008, 39, 785-790.	1.5	21
172	Intravenous N-acetylcysteine with Saline Hydration Improves Renal Function and Ameliorates Plasma Total Homocysteine in Patients Undergoing Cardiac Angiography. <i>Renal Failure</i> , 2008, 30, 527-533.	0.8	14
173	Fasting Serum Ghrelin Level is Associated with Metabolic Syndrome in Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2008, 28, 196-200.	1.1	6
174	TT Virus Infection in Patients on Peritoneal Dialysis in Taiwan. <i>Renal Failure</i> , 2007, 29, 553-557.	0.8	1
175	Sen Virus Infection in Patients on Peritoneal Dialysis. <i>Peritoneal Dialysis International</i> , 2007, 27, 74-78.	1.1	0
176	Fluvastatin ameliorates endotoxin induced multiple organ failure in conscious rats. <i>Resuscitation</i> , 2007, 74, 166-174.	1.3	33
177	SEN virus infection in patients on peritoneal dialysis. <i>Peritoneal Dialysis International</i> , 2007, 27, 74-8.	1.1	1
178	Post-treatment with N-acetylcysteine ameliorates endotoxin shock-induced organ damage in conscious rats. <i>Life Sciences</i> , 2006, 79, 2010-2016.	2.0	87