

# Gard F T Svingen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2103663/publications.pdf>

Version: 2024-02-01

23  
papers

643  
citations

759233

12  
h-index

642732

23  
g-index

23  
all docs

23  
docs citations

23  
times ranked

1538  
citing authors

#	ARTICLE	IF	CITATIONS
1	$\beta$ -blocker use and risk of all-cause mortality in patients with coronary heart disease: effect modification by serum vitamin A. <i>European Journal of Preventive Cardiology</i> , 2022, 28, 1897-1902.	1.8	5
2	Trimethyllysine predicts all-cause and cardiovascular mortality in community-dwelling adults and patients with coronary heart disease. <i>European Heart Journal Open</i> , 2021, 1, .	2.3	4
3	Creatinine, total cysteine and uric acid are associated with serum retinol in patients with cardiovascular disease. <i>European Journal of Nutrition</i> , 2020, 59, 2383-2393.	3.9	10
4	Dietary choline is related to increased risk of acute myocardial infarction in patients with stable angina pectoris. <i>Biochimie</i> , 2020, 173, 68-75.	2.6	11
5	Lipid parameters and vitamin A modify cardiovascular risk prediction by plasma neopterin. <i>Heart</i> , 2020, 106, 1073-1079.	2.9	4
6	Transsulfuration metabolites and the association with incident atrial fibrillation – An observational cohort study among Norwegian patients with stable angina pectoris. <i>International Journal of Cardiology</i> , 2020, 317, 75-80.	1.7	5
7	Elevated plasma cystathionine is associated with increased risk of mortality among patients with suspected or established coronary heart disease. <i>American Journal of Clinical Nutrition</i> , 2019, 109, 1546-1554.	4.7	8
8	Plasma Concentrations and Dietary Intakes of Choline and Betaine in Association With Atrial Fibrillation Risk: Results From 3 Prospective Cohorts With Different Health Profiles. <i>Journal of the American Heart Association</i> , 2018, 7, .	3.7	31
9	Serum Carnitine Metabolites and Incident Type 2 Diabetes Mellitus in Patients With Suspected Stable Angina Pectoris. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 1033-1041.	3.6	27
10	Plasma methionine and risk of acute myocardial infarction: Effect modification by established risk factors. <i>Atherosclerosis</i> , 2018, 272, 175-181.	0.8	13
11	Plasma Cystathionine and Risk of Incident Stroke in Patients With Suspected Stable Angina Pectoris. <i>Journal of the American Heart Association</i> , 2018, 7, e008824.	3.7	14
12	Fibrinogen and Neopterin Is Associated with Future Myocardial Infarction and Total Mortality in Patients with Stable Coronary Artery Disease. <i>Thrombosis and Haemostasis</i> , 2018, 47, 778-790.	3.4	16
13	Plasma cystathionine and risk of acute myocardial infarction among patients with coronary heart disease: Results from two independent cohorts. <i>International Journal of Cardiology</i> , 2018, 266, 24-30.	1.7	15
14	Usefulness of Higher Levels of Cardiac Troponin T in Patients With Stable Angina Pectoris to Predict Risk of Acute Myocardial Infarction. <i>American Journal of Cardiology</i> , 2018, 122, 1142-1147.	1.6	11
15	Increased plasma trimethylamine- N -oxide is associated with incident atrial fibrillation. <i>International Journal of Cardiology</i> , 2018, 267, 100-106.	1.7	67
16	Plasma Homoarginine Concentrations According to Use of Hormonal Contraception. <i>Scientific Reports</i> , 2018, 8, 12217.	3.3	5
17	The kynurenine:tryptophan ratio as a predictor of incident type 2 diabetes mellitus in individuals with coronary artery disease. <i>Diabetologia</i> , 2017, 60, 1712-1721.	6.3	58
18	Relations between lipoprotein(a) concentrations, LPA genetic variants, and the risk of mortality in patients with established coronary heart disease: a molecular and genetic association study. <i>Lancet Diabetes and Endocrinology</i> , 2017, 5, 534-543.	11.4	84

#	ARTICLE	IF	CITATIONS
19	Cardiovascular disease risk associated with serum apolipoprotein B is modified by serum vitamin A. <i>Atherosclerosis</i> , 2017, 265, 325-330.	0.8	12
20	Neopterin as an Effect Modifier of the Cardiovascular Risk Predicted by Total Homocysteine: A Prospective Cohort Study. <i>Journal of the American Heart Association</i> , 2017, 6, .	3.7	12
21	Methylenetetrahydrofolate Dehydrogenase 1 Polymorphisms Modify the Associations of Plasma Glycine and Serine With Risk of Acute Myocardial Infarction in Patients With Stable Angina Pectoris in WENBIT (Western Norway B Vitamin Intervention Trial). <i>Circulation: Cardiovascular Genetics</i> , 2016, 9, 541-547.	5.1	6
22	Associations of Plasma Kynurenines With Risk of Acute Myocardial Infarction in Patients With Stable Angina Pectoris. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2015, 35, 455-462.	2.4	133
23	Plasma Dimethylglycine and Risk of Incident Acute Myocardial Infarction in Patients With Stable Angina Pectoris. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2013, 33, 2041-2048.	2.4	92