

Miguel Angel Rodriguez-Delgado

List of Publications by Year in Descending Order

Source:
<https://exaly.com/author-pdf/2113562/miguel-angel-rodriguez-delgado-publications-by-year.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

111 papers	5,063 citations	41 h-index	67 g-index
113 ext. papers	5,504 ext. citations	5.2 avg, IF	5.86 L-index

#	Paper	IF	Citations
111	Safety assessment of citrus and olive by-products using a sustainable methodology based on natural deep eutectic solvents.. <i>Journal of Chromatography A</i> , 2022 , 1669, 462922	4.5	1
110	Application of polyaniline-based magnetic-dispersive-solid-phase microextraction combined with liquid chromatography tandem mass spectrometry for the evaluation of plastic migrants in food matrices.. <i>Journal of Chromatography A</i> , 2022 , 1670, 462988	4.5	2
109	Analysis of alkylphenols, bisphenols and alkylphenol ethoxylates in microbial-fermented functional beverages and bottled water: Optimization of a dispersive liquid-liquid microextraction protocol based on natural hydrophobic deep eutectic solvents.. <i>Food Chemistry</i> , 2021 , 377, 131921	8.5	1
108	Recent Applications of Deep Eutectic Solvents in Environmental Analysis. <i>Applied Sciences (Switzerland)</i> , 2021 , 11, 4779	2.6	6
107	Deep eutectic solvents. The new generation of green solvents in analytical chemistry. <i>TrAC - Trends in Analytical Chemistry</i> , 2021 , 134, 116108	14.6	41
106	Sustainable polypyrrole-based magnetic-microextraction of phthalates from jellies and apple-based beverages prior to tandem mass spectrometry analysis. <i>Journal of Chromatography A</i> , 2021 , 1637, 461858	4.5	1
105	Development of a Green Alternative Vortex-Assisted Dispersive Liquid-Liquid Microextraction Based on Natural Hydrophobic Deep Eutectic Solvents for the Analysis of Phthalate Esters in Soft Drinks. <i>ACS Sustainable Chemistry and Engineering</i> , 2021 , 9, 2161-2170	8.3	7
104	Quality assessment of environmental water by a simple and fast non-ionic hydrophobic natural deep eutectic solvent-based extraction procedure combined with liquid chromatography tandem mass spectrometry for the determination of plastic migrants. <i>Analytical and Bioanalytical Chemistry</i> , 2021 , 413, 1967-1981	4.4	7
103	Novel applications of nanotechnology in food safety assessment 2021 , 461-505		1
102	Comparison of Pesticide Residue Levels in Red Wines from Canary Islands, Iberian Peninsula, and Cape Verde. <i>Foods</i> , 2020 , 9,	4.9	6
101	Deep eutectic solvents 2020 , 123-177		5
100	A simple, fast and easy methodology for the monitoring of plastic migrants in alcoholic and non-alcoholic beverages using the QuEChERS method prior to gas chromatography tandem mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2020 , 412, 1551-1561	4.4	4
99	Nano-liquid chromatography combined with a sustainable microextraction based on natural deep eutectic solvents for analysis of phthalate esters. <i>Electrophoresis</i> , 2020 , 41, 1768-1775	3.6	5
98	A green and simple procedure based on deep eutectic solvents for the extraction of phthalates from beverages. <i>Food Chemistry</i> , 2020 , 312, 125798	8.5	21
97	Recent Applications of Magnetic Nanoparticles in Food Analysis. <i>Processes</i> , 2020 , 8, 1140	2.9	3
96	Critical review and re-assessment of analyte protectants in gas chromatography. <i>Journal of Chromatography A</i> , 2020 , 1632, 461596	4.5	8
95	Carbon-based adsorbents 2020 , 83-127		4

94	Current trends in QuEChERS method. A versatile procedure for food, environmental and biological analysis. <i>TrAC - Trends in Analytical Chemistry</i> , 2019 , 116, 214-235	14.6	71
93	Nanomaterials as alternative dispersants for the multiresidue analysis of phthalates in soil samples using matrix solid phase dispersion prior to ultra-high performance liquid chromatography tandem mass spectrometry. <i>Chemosphere</i> , 2019 , 236, 124377	8.4	10
92	Use of Basolite F300 metal-organic framework for the dispersive solid-phase extraction of phthalic acid esters from water samples prior to LC-MS determination. <i>Talanta</i> , 2019 , 195, 236-244	6.2	30
91	Determination of phthalates in beverages using multiwalled carbon nanotubes dispersive solid-phase extraction before HPLC-MS. <i>Journal of Separation Science</i> , 2018 , 41, 2613-2622	3.4	21
90	Multiresidue analysis of oestrogenic compounds in cow, goat, sheep and human milk using core-shell polydopamine coated magnetic nanoparticles as extraction sorbent in micro-dispersive solid-phase extraction followed by ultra-high-performance liquid chromatography tandem mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2018 , 410, 2031-2042	4.4	23
89	Determination of phthalic acid esters in water samples by hollow fiber liquid-phase microextraction prior to gas chromatography tandem mass spectrometry. <i>Chemosphere</i> , 2018 , 201, 254-261	8.4	33
88	Reduced graphene oxide-coated magnetic-nanoparticles as sorbent for the determination of phthalates in environmental samples by micro-dispersive solid-phase extraction followed by ultra-high-performance liquid chromatography tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2018 , 1565, 36-47	4.5	33
87	Multiresidue determination of estrogens in different dairy products by ultra-high-performance liquid chromatography triple quadrupole mass spectrometry. <i>Journal of Chromatography A</i> , 2017 , 1496, 58-67	4.5	21
86	Determination of phthalic acid esters in water samples using core-shell poly(dopamine) magnetic nanoparticles and gas chromatography tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2017 , 1530, 35-44	4.5	31
85	Recent applications of nanomaterials in food safety. <i>TrAC - Trends in Analytical Chemistry</i> , 2017 , 96, 172-200	20.6	58
84	Multiclass analytical method for the determination of natural/synthetic steroid hormones, phytoestrogens, and mycoestrogens in milk and yogurt. <i>Analytical and Bioanalytical Chemistry</i> , 2017 , 409, 4467-4477	4.4	14
83	Core-shell poly(dopamine) magnetic nanoparticles for the extraction of estrogenic mycotoxins from milk and yogurt prior to LC-MS analysis. <i>Food Chemistry</i> , 2017 , 215, 362-8	8.5	42
82	Estrogenic Compounds in Yogurt 2017 , 451-472		
81	Determination of estrogenic compounds in milk and yogurt samples by hollow-fibre liquid-phase microextraction-gas chromatography-triple quadrupole mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2016 , 408, 7447-59	4.4	13
80	Nanomaterials as sorbents for food sample analysis. <i>TrAC - Trends in Analytical Chemistry</i> , 2016 , 85, 203-220	20.6	69
79	Capillary electrochromatography and nano-liquid chromatography coupled to nano-electrospray ionization interface for the separation and identification of estrogenic compounds. <i>Electrophoresis</i> , 2016 , 37, 356-62	3.6	13
78	Application of multiwalled carbon nanotubes as sorbents for the extraction of mycotoxins in water samples and infant milk formula prior to high performance liquid chromatography mass spectrometry analysis. <i>Electrophoresis</i> , 2016 , 37, 1359-66	3.6	17
77	Evolution and applications of the QuEChERS method. <i>TrAC - Trends in Analytical Chemistry</i> , 2015 , 71, 169-185	14.6	214

76	Core-shell polydopamine magnetic nanoparticles as sorbent in micro-dispersive solid-phase extraction for the determination of estrogenic compounds in water samples prior to high-performance liquid chromatography-mass spectrometry analysis. <i>Journal of Chromatography A</i> 2015 , 1397, 1-10	4.5	48
75	Evaluation of two molecularly imprinted polymers for the solid-phase extraction of natural, synthetic and mycoestrogens from environmental water samples before liquid chromatography with mass spectrometry. <i>Journal of Separation Science</i> , 2015 , 38, 2692-9	3.4	26
74	Evaluation of the combination of a dispersive liquid-liquid microextraction method with micellar electrokinetic chromatography coupled to mass spectrometry for the determination of estrogenic compounds in milk and yogurt. <i>Electrophoresis</i> , 2015 , 36, 615-25	3.6	36
73	Estrogenic compounds determination in water samples by dispersive liquid-liquid microextraction and micellar electrokinetic chromatography coupled to mass spectrometry. <i>Journal of Chromatography A</i> , 2014 , 1344, 109-21	4.5	40
72	Determination of estrogens in environmental water samples using 1,3-dipentylimidazolium hexafluorophosphate ionic liquid as extraction solvent in dispersive liquid-liquid microextraction. <i>Electrophoresis</i> , 2014 , 35, 2479-87	3.6	24
71	Analysis of oestrogenic compounds in dairy products by hollow-fibre liquid-phase microextraction coupled to liquid chromatography. <i>Food Chemistry</i> , 2014 , 149, 319-25	8.5	31
70	Use of ammonium formate in QuEChERS for high-throughput analysis of pesticides in food by fast, low-pressure gas chromatography and liquid chromatography tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2014 , 1358, 75-84	4.5	78
69	Determination of organophosphorus pesticides and metabolites in cereal-based baby foods and wheat flour by means of ultrasound-assisted extraction and hollow-fiber liquid-phase microextraction prior to gas chromatography with nitrogen phosphorus detection. <i>Journal of Chromatography A</i> , 2013 , 1272, 144-51	4.5	44
68	Comparison between magnetic and non magnetic multi-walled carbon nanotubes-dispersive solid-phase extraction combined with ultra-high performance liquid chromatography for the determination of sulfonamide antibiotics in water samples. <i>Talanta</i> , 2013 , 116, 695-703	6.2	91
67	Hollow-fiber liquid-phase microextraction for the determination of natural and synthetic estrogens in milk samples. <i>Journal of Chromatography A</i> , 2013 , 1313, 175-84	4.5	38
66	Dispersive liquid-liquid microextraction combined with ultra-high performance liquid chromatography for the simultaneous determination of 25 sulfonamide and quinolone antibiotics in water samples. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2013 , 75, 130-7	3.5	103
65	Analysis of pesticides residues in environmental water samples using multiwalled carbon nanotubes dispersive solid-phase extraction. <i>Journal of Separation Science</i> , 2013 , 36, 556-63	3.4	55
64	Chromatographic analysis of natural and synthetic estrogens in milk and dairy products. <i>TrAC - Trends in Analytical Chemistry</i> , 2013 , 44, 58-77	14.6	45
63	Carbon nanotubes applications in separation science: a review. <i>Analytica Chimica Acta</i> , 2012 , 734, 1-30	6.6	264
62	Dispersive liquid-liquid microextraction of pesticides and metabolites from soils using 1,3-dipentylimidazolium hexafluorophosphate ionic liquid as an alternative extraction solvent. <i>Electrophoresis</i> , 2012 , 33, 1449-57	3.6	19
61	Hollow-fiber liquid-phase microextraction for the determination of pesticides and metabolites in soils and water samples using HPLC and fluorescence detection. <i>Electrophoresis</i> , 2012 , 33, 2184-91	3.6	30
60	Determination of quinolone residues in infant and young children powdered milk combining solid-phase extraction and ultra-performance liquid chromatography-tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2011 , 1218, 7608-14	4.5	43
59	Ionic liquid-dispersive liquid-liquid microextraction for the simultaneous determination of pesticides and metabolites in soils using high-performance liquid chromatography and fluorescence detection. <i>Journal of Chromatography A</i> , 2011 , 1218, 4808-16	4.5	105

58	Oxidized multi-walled carbon nanotubes for the dispersive solid-phase extraction of quinolone antibiotics from water samples using capillary electrophoresis and large volume sample stacking with polarity switching. <i>Journal of Chromatography A</i> , 2011 , 1218, 5352-61	4.5	83
57	Insecticides extraction from banana leaves using a modified QuEChERS method. <i>Food Chemistry</i> , 2011 , 125, 1083-1090	8.5	31
56	Evaluation of a modified QuEChERS method for the extraction of pesticides from agricultural, ornamental and forestal soils. <i>Analytical and Bioanalytical Chemistry</i> , 2010 , 396, 2307-19	4.4	92
55	Dispersive liquid-liquid microextraction combined with nonaqueous capillary electrophoresis for the determination of fluoroquinolone antibiotics in waters. <i>Electrophoresis</i> , 2010 , 31, 3457-65	3.6	55
54	Dispersive liquid-liquid microextraction for determination of organic analytes. <i>TrAC - Trends in Analytical Chemistry</i> , 2010 , 29, 728-751	14.6	219
53	Carbon nanotubes: Solid-phase extraction. <i>Journal of Chromatography A</i> , 2010 , 1217, 2618-41	4.5	281
52	Recent food safety and food quality applications of CE-MS. <i>Electrophoresis</i> , 2009 , 30, 1624-46	3.6	30
51	Pesticide extraction from table grapes and plums using ionic liquid based dispersive liquid-liquid microextraction. <i>Analytical and Bioanalytical Chemistry</i> , 2009 , 395, 2387-95	4.4	57
50	Fluoroquinolone antibiotic determination in bovine, ovine and caprine milk using solid-phase extraction and high-performance liquid chromatography-fluorescence detection with ionic liquids as mobile phase additives. <i>Journal of Chromatography A</i> , 2009 , 1216, 7281-7	4.5	58
49	Ionic liquid based dispersive liquid-liquid microextraction for the extraction of pesticides from bananas. <i>Journal of Chromatography A</i> , 2009 , 1216, 7336-45	4.5	138
48	Analysis of pesticide residues in bananas harvested in the Canary Islands (Spain). <i>Food Chemistry</i> , 2009 , 113, 313-319	8.5	70
47	Evaluation of multi-walled carbon nanotubes as solid-phase extraction adsorbents of pesticides from agricultural, ornamental and forestal soils. <i>Analytica Chimica Acta</i> , 2009 , 647, 167-76	6.6	68
46	Ionic liquids as mobile phase additives for the high-performance liquid chromatographic analysis of fluoroquinolone antibiotics in water samples. <i>Analytical and Bioanalytical Chemistry</i> , 2008 , 392, 1439-46	4.4	50
45	Multiwalled carbon nanotubes as solid-phase extraction materials for the gas chromatographic determination of organophosphorus pesticides in waters. <i>Journal of Separation Science</i> , 2008 , 31, 3612-9	3.4	33
44	Simultaneous determination of seven pesticides in waters using multi-walled carbon nanotube SPE and NACE. <i>Electrophoresis</i> , 2008 , 29, 4412-21	3.6	40
43	Solid-phase microextraction and sample stacking micellar electrokinetic chromatography for the analysis of pesticide residues in red wines. <i>Food Chemistry</i> , 2008 , 111, 764-770	8.5	58
42	Multi-walled carbon nanotubes as efficient solid-phase extraction materials of organophosphorus pesticides from apple, grape, orange and pineapple fruit juices. <i>Journal of Chromatography A</i> , 2008 , 1211, 33-42	4.5	127
41	Pesticide analysis in tomatoes by solid-phase microextraction and micellar electrokinetic chromatography. <i>Journal of Chromatography A</i> , 2008 , 1185, 151-4	4.5	34

40	Determination of Abamectin Residues in Avocados by Microwave-Assisted Extraction and HPLC with Fluorescence Detection. <i>Chromatographia</i> , 2008 , 67, 69-75	2.1	20
39	MEKC combined with SPE and sample stacking for multiple analysis of pesticides in water samples at the ng/L level. <i>Electrophoresis</i> , 2007 , 28, 1805-14	3.6	28
38	Multiple pesticide analysis in wine by MEKC combined with solid-phase microextraction and sample stacking. <i>Electrophoresis</i> , 2007 , 28, 4072-81	3.6	31
37	Rapid analysis of triazolopyrimidine sulfoanilide herbicides in waters and soils by high-performance liquid chromatography with UV detection using a C18 monolithic column. <i>Journal of Separation Science</i> , 2007 , 30, 8-14	3.4	24
36	Pesticide analysis in rose wines by micellar electrokinetic chromatography. <i>Journal of Separation Science</i> , 2007 , 30, 3240-6	3.4	17
35	Determination of pesticides in wine using micellar electrokinetic chromatography with UV detection and sample stacking. <i>Journal of Chromatography A</i> , 2007 , 1150, 348-55	4.5	28
34	Sample treatments prior to capillary electrophoresis-mass spectrometry. <i>Journal of Chromatography A</i> , 2007 , 1153, 214-26	4.5	49
33	Analysis of abamectin residues in avocados by high-performance liquid chromatography with fluorescence detection. <i>Journal of Chromatography A</i> , 2007 , 1165, 52-7	4.5	23
32	Pesticides analysis by liquid chromatography and capillary electrophoresis. <i>Journal of Separation Science</i> , 2006 , 29, 2557-77	3.4	32
31	Optimization of the Microwave-Assisted Saponification and Extraction of Organic Pollutants from Marine Biota Using Experimental Design and Artificial Neural Networks. <i>Chromatographia</i> , 2006 , 63, 155-160	2.1	14
30	Determination of antioxidants in edible grain derivatives from the Canary Islands by capillary electrophoresis. <i>Food Chemistry</i> , 2005 , 91, 105-111	8.5	25
29	Determination of herbicides in mineral and stagnant waters at ng/L levels using capillary electrophoresis and UV detection combined with solid-phase extraction and sample stacking. <i>Journal of Chromatography A</i> , 2005 , 1070, 171-7	4.5	42
28	Analysis of triazolopyrimidine herbicides in soils using field-enhanced sample injection-coelectroosmotic capillary electrophoresis combined with solid-phase extraction. <i>Journal of Chromatography A</i> , 2005 , 1100, 236-42	4.5	30
27	Rapid Separation of Antioxidants in Food Samples by Coelectroosmotic CE. <i>Chromatographia</i> , 2005 , 62, 271-276	2.1	12
26	Combining solid-phase microextraction and on-line preconcentration-capillary electrophoresis for sensitive analysis of pesticides in foods. <i>Electrophoresis</i> , 2005 , 26, 980-9	3.6	53
25	Chiral analysis of pollutants and their metabolites by capillary electromigration methods. <i>Electrophoresis</i> , 2005 , 26, 3799-813	3.6	35
24	Analysis of pesticides in soy milk combining solid-phase extraction and capillary electrophoresis-mass spectrometry. <i>Journal of Separation Science</i> , 2005 , 28, 948-56	3.4	58
23	Highly sensitive analysis of multiple pesticides in foods combining solid-phase microextraction, capillary electrophoresis-mass spectrometry, and chemometrics. <i>Electrophoresis</i> , 2004 , 25, 2065-76	3.6	65

22	Optimization of a solid-phase microextraction procedure for the determination of herbicides by micellar electrokinetic chromatography. <i>Journal of Separation Science</i> , 2004 , 27, 660-6	3.4	21
21	Pesticide analysis by capillary electrophoresis. <i>Journal of Separation Science</i> , 2004 , 27, 947-63	3.4	71
20	Content of aliphatic hydrocarbons in limpets as a new way for classification of species using artificial neural networks. <i>Chemosphere</i> , 2004 , 54, 1059-69	8.4	18
19	Content of free phenolic compounds in bananas from Tenerife (Canary Islands) and Ecuador. <i>European Food Research and Technology</i> , 2003 , 217, 287-290	3.4	36
18	Fast analysis of proteins in wines by capillary gel electrophoresis. <i>European Food Research and Technology</i> , 2002 , 214, 536-540	3.4	11
17	Fast determination of retinol and alpha-tocopherol in plasma by LC. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2002 , 28, 991-7	3.5	12
16	Trans-resveratrol in wines from the Canary Islands (Spain). Analysis by high performance liquid chromatography. <i>Food Chemistry</i> , 2002 , 76, 371-375	8.5	45
15	Principal component analysis of the polyphenol content in young red wines. <i>Food Chemistry</i> , 2002 , 78, 523-532	8.5	129
14	Comparison of different coatings in solid-phase microextraction for the determination of organochlorine pesticides in ground water. <i>Journal of Chromatography A</i> , 2002 , 963, 95-105	4.5	31
13	Determination of esters in dry and sweet white wines by headspace solid-phase microextraction and gas chromatography. <i>Journal of Chromatography A</i> , 2002 , 963, 213-23	4.5	70
12	Ultrafast sodium dodecyl sulfate micellar electrokinetic chromatography with very acidic running buffers. <i>Analytical Chemistry</i> , 2002 , 74, 257-60	7.8	19
11	Separation of phenolic compounds by high-performance liquid chromatography with absorbance and fluorimetric detection. <i>Journal of Chromatography A</i> , 2001 , 912, 249-57	4.5	260
10	Optimisation of sample preparation for the determination of trans-resveratrol and other polyphenolic compounds in wines by high performance liquid chromatography. <i>Analytica Chimica Acta</i> , 2001 , 428, 245-253	6.6	98
9	Optimization of the separation of phenolic compounds by micellar electrokinetic capillary chromatography. <i>Journal of Chromatography A</i> , 2000 , 871, 427-38	4.5	57
8	Determination of polycyclic aromatic hydrocarbons in limpet samples. <i>Chromatographia</i> , 1999 , 50, 235-238		4
7	Comparative study of the zero-crossing, ratio spectra derivative and partial least-squares methods applied to the simultaneous determination of atrazine and its degradation product desethylatrazin-2-hydroxy in ground waters. <i>Talanta</i> , 1997 , 44, 673-83	6.2	23
6	Separation and quantitation of debrisoquine and 4-hydroxydebrisoquine in human urine by capillary electrophoresis and high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1997 , 778, 389-96	4.5	10
5	Comparative study of the ratio spectra derivative and partial least squares methods applied to the simultaneous determination of atrazine and ametryn in ground waters. <i>Analyst, The</i> , 1996 , 121, 459-463 ⁵		1

4	Solute-micelle association constants of some polynuclear aromatic hydrocarbons by micellar liquid chromatography with alcohol additives to mobile phase. <i>Chromatographia</i> , 1994 , 38, 342-348	2.1	22
3	Correlations between retention data of polycyclic aromatic hydrocarbons in micellar liquid chromatography and several molecular descriptors. <i>Fresenius Journal of Analytical Chemistry</i> , 1993 , 345, 748-752		13
2	4th order derivative spectrophotometric determination of quinine in soft drinks. <i>Mikrochimica Acta</i> , 1993 , 110, 263-268	5.8	8
1	Solute-micelle association constants and correlation of octanol-water coefficients with hydrophobicity for polycyclic aromatic hydrocarbons by micellar chromatography. <i>Chromatographia</i> , 1992 , 34, 627-635	2.1	38