

James S Wiley

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2117723/publications.pdf>

Version: 2024-02-01

113
papers

7,535
citations

53660

45
h-index

54797

84
g-index

113
all docs

113
docs citations

113
times ranked

8408
citing authors

#	ARTICLE	IF	CITATIONS
1	Analysis of immune-related loci identifies 48 new susceptibility variants for multiple sclerosis. <i>Nature Genetics</i> , 2013, 45, 1353-1360.	9.4	1,213
2	Genome-wide meta-analysis identifies novel multiple sclerosis susceptibility loci. <i>Annals of Neurology</i> , 2011, 70, 897-912.	2.8	314
3	A Glu-496 to Ala Polymorphism Leads to Loss of Function of the Human P2X7 Receptor. <i>Journal of Biological Chemistry</i> , 2001, 276, 11135-11142.	1.6	276
4	A Furosemide-Sensitive Cotransport of Sodium plus Potassium in the Human Red Cell. <i>Journal of Clinical Investigation</i> , 1974, 53, 745-755.	3.9	228
5	A Polymorphism in the P2X7 Gene Increases Susceptibility to Extrapulmonary Tuberculosis. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2007, 175, 360-366.	2.5	188
6	The isoquinoline derivative KN-62 a potent antagonist of the P2Z-receptor of human lymphocytes. <i>British Journal of Pharmacology</i> , 1997, 120, 1483-1490.	2.7	174
7	The human P2X7 receptor and its role in innate immunity. <i>Tissue Antigens</i> , 2011, 78, 321-332.	1.0	172
8	Strikingly homologous immunoglobulin gene rearrangements and poor outcome in VH3-21-using chronic lymphocytic leukemia patients independent of geographic origin and mutational status. <i>Blood</i> , 2006, 107, 2889-2894.	0.6	167
9	Two haplotypes of the P2X ₇ receptor containing the Ala348 to Thr polymorphism exhibit a gain-of-function effect and enhanced interleukin-1 β secretion. <i>FASEB Journal</i> , 2010, 24, 2916-2927.	0.2	155
10	An Ile-568 to Asn Polymorphism Prevents Normal Trafficking and Function of the Human P2X7 Receptor. <i>Journal of Biological Chemistry</i> , 2003, 278, 17108-17113.	1.6	154
11	P2X7 Receptor Cell Surface Expression and Cytolytic Pore Formation Are Regulated by a Distal C-terminal Region. <i>Journal of Biological Chemistry</i> , 2003, 278, 8853-8860.	1.6	153
12	A Thr357 to Ser Polymorphism in Homozygous and Compound Heterozygous Subjects Causes Absent or Reduced P2X7 Function and Impairs ATP-induced Mycobacterial Killing by Macrophages. <i>Journal of Biological Chemistry</i> , 2006, 281, 2079-2086.	1.6	152
13	Rapid ATP-induced release of matrix metalloproteinase 9 is mediated by the P2X7 receptor. <i>Blood</i> , 2006, 107, 4946-4953.	0.6	149
14	A loss-of-function polymorphic mutation in the cytolytic P2X7 receptor gene and chronic lymphocytic leukaemia: a molecular study. <i>Lancet</i> , The, 2002, 359, 1114-1119.	6.3	145
15	Glu496 to Ala Polymorphism in the P2X7 Receptor Impairs ATP-Induced IL-1 β Release from Human Monocytes. <i>Journal of Immunology</i> , 2004, 172, 3399-3405.	0.4	140
16	An Arg307 to Gln Polymorphism within the ATP-binding Site Causes Loss of Function of the Human P2X7 Receptor. <i>Journal of Biological Chemistry</i> , 2004, 279, 31287-31295.	1.6	125
17	P2X7 Receptor-Mediated Killing of an Intracellular Parasite, <i>Toxoplasma gondii</i> , by Human and Murine Macrophages. <i>Journal of Immunology</i> , 2010, 184, 7040-7046.	0.4	124
18	X-linked Pyridoxine-Responsive Sideroblastic Anemia Due to a Thr388-to-Ser Substitution in Erythroid 5-Aminolevulinic Synthase. <i>New England Journal of Medicine</i> , 1994, 330, 675-679.	13.9	122

#	ARTICLE	IF	CITATIONS
19	The Role of the P2X7 Receptor in Infectious Diseases. <i>PLoS Pathogens</i> , 2011, 7, e1002212.	2.1	121
20	A Loss-of-Function Polymorphism in the Human P2X7 Receptor Abolishes ATP-Mediated Killing of Mycobacteria. <i>Journal of Immunology</i> , 2003, 171, 5442-5446.	0.4	115
21	Genetics of the P2X7 receptor and human disease. <i>Purinergic Signalling</i> , 2009, 5, 257-262.	1.1	114
22	Inhibition of cation contransport by cholesterol enrichment of human red cell membranes. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1975, 413, 425-431.	1.4	101
23	Extracellular ATP causes loss of L-selectin from human lymphocytes via occupancy of P2Z purinoceptors. , 1996, 166, 637-642.		92
24	The P2X7-nonmuscle myosin membrane complex regulates phagocytosis of nonopsonized particles and bacteria by a pathway attenuated by extracellular ATP. <i>Blood</i> , 2010, 115, 1621-1631.	0.6	90
25	The ATP4â” receptor-operated ion channel of human lymphocytes: Inhibition of ion fluxes by amiloride analogs and by extracellular sodium ions. <i>Archives of Biochemistry and Biophysics</i> , 1992, 292, 411-418.	1.4	86
26	The P_{2Z}-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP. <i>British Journal of Pharmacology</i> , 1994, 112, 946-950.	2.7	82
27	P2X7 Is a Scavenger Receptor for Apoptotic Cells in the Absence of Its Ligand, Extracellular ATP. <i>Journal of Immunology</i> , 2011, 187, 2365-2375.	0.4	81
28	Extracellular ATP dissociates nonmuscle myosin from P2X₇-complex: this dissociation regulates P2X₇-pore formation. <i>American Journal of Physiology - Cell Physiology</i> , 2009, 297, C430-C439.	2.1	79
29	Detection of P2X purinergic receptors on human B lymphocytes. <i>Cell and Tissue Research</i> , 2001, 304, 231-236.	1.5	70
30	Purinoceptors are involved in the induction of an osmolyte permeability in malariaâ€infectected and oxidized human erythrocytes. <i>FASEB Journal</i> , 2006, 20, 133-135.	0.2	69
31	Phospholipase D activation by P2Z-purinoceptor agonists in human lymphocytes is dependent on bivalent cation influx. <i>Biochemical Journal</i> , 1996, 313, 529-535.	1.7	65
32	Extracellular ATP Increases Cation Fluxes in Human Erythrocytes by Activation of the P2X7 Receptor. <i>Journal of Biological Chemistry</i> , 2004, 279, 44749-44755.	1.6	65
33	A High-Density SNP Genomewide Linkage Scan for Chronic Lymphocytic Leukemiaâ€Susceptibility Loci. <i>American Journal of Human Genetics</i> , 2005, 77, 420-429.	2.6	65
34	Analysis of human leukaemias and lymphomas using extensive immunophenotypes from an antibody microarray. <i>British Journal of Haematology</i> , 2006, 135, 184-197.	1.2	65
35	Gene Dosage Determines the Negative Effects of Polymorphic Alleles of the P2X7Receptor on Adenosine Triphosphateâ€Mediated Killing of Mycobacteria by Human Macrophages. <i>Journal of Infectious Diseases</i> , 2005, 192, 149-155.	1.9	64
36	Single-nucleotide polymorphisms in the P2X7 receptor gene are associated with post-menopausal bone loss and vertebral fractures. <i>European Journal of Human Genetics</i> , 2012, 20, 675-681.	1.4	63

#	ARTICLE	IF	CITATIONS
37	Polymorphisms in the P2X7 receptor gene are associated with low lumbar spine bone mineral density and accelerated bone loss in post-menopausal women. <i>European Journal of Human Genetics</i> , 2012, 20, 559-564.	1.4	63
38	A rare functional haplotype of the <i>P2RX4</i> and <i>P2RX7</i> genes leads to loss of innate phagocytosis and confers increased risk of age-related macular degeneration. <i>FASEB Journal</i> , 2013, 27, 1479-1487.	0.2	61
39	ATP, a partial agonist for the P2Z receptor of human lymphocytes. <i>British Journal of Pharmacology</i> , 1997, 122, 911-917.	2.7	57
40	A 5' intronic splice site polymorphism leads to a null allele of the P2X7 gene in 1-2% of the Caucasian population. <i>FEBS Letters</i> , 2005, 579, 2675-2678.	1.3	55
41	A Polymorphism in the HLA-DPB1 Gene Is Associated with Susceptibility to Multiple Sclerosis. <i>PLoS ONE</i> , 2010, 5, e13454.	1.1	55
42	A rare P2X7 variant Arg307Gln with absent pore formation function protects against neuroinflammation in multiple sclerosis. <i>Human Molecular Genetics</i> , 2015, 24, 5644-5654.	1.4	53
43	A Loss-of-Function Polymorphism in the Human P2X4 Receptor Is Associated With Increased Pulse Pressure. <i>Hypertension</i> , 2011, 58, 1086-1092.	1.3	52
44	P2X7 as a scavenger receptor for innate phagocytosis in the brain. <i>British Journal of Pharmacology</i> , 2018, 175, 4195-4208.	2.7	50
45	Chelerythrine and other benzophenanthridine alkaloids block the human P2X7 receptor. <i>British Journal of Pharmacology</i> , 2004, 142, 1015-1019.	2.7	48
46	Extracellular adenosine 5'-triphosphate induces a loss of CD23 from human dendritic cells via activation of P2X7 receptors. <i>International Immunology</i> , 2002, 14, 1415-1421.	1.8	46
47	Point mutations confer loss of ATP-induced human P2X7 receptor function. <i>FEBS Letters</i> , 2002, 512, 43-46.	1.3	46
48	Purinergic receptors <i>P2RX4</i> and <i>P2RX7</i> in familial multiple sclerosis. <i>Human Mutation</i> , 2017, 38, 736-744.	1.1	46
49	Human Epidermal and Monocyte-Derived Langerhans Cells Express Functional P2X7 Receptors. <i>Journal of Investigative Dermatology</i> , 2005, 125, 482-490.	0.3	45
50	Shear stress modulates endothelial KLF2 through activation of P2X4. <i>Purinergic Signalling</i> , 2015, 11, 139-153.	1.1	41
51	P2X7 Receptors Mediate Innate Phagocytosis by Human Neural Precursor Cells and Neuroblasts. <i>Stem Cells</i> , 2015, 33, 526-541.	1.4	40
52	Innate phagocytosis by peripheral blood monocytes is altered in Alzheimer's disease. <i>Acta Neuropathologica</i> , 2016, 132, 377-389.	3.9	40
53	A new role for the P2X7 receptor: a scavenger receptor for bacteria and apoptotic cells in the absence of serum and extracellular ATP. <i>Purinergic Signalling</i> , 2012, 8, 579-586.	1.1	39
54	P2X7 receptor activation induces cell death and microparticle release in murine erythroleukemia cells. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2010, 1798, 1797-1804.	1.4	38

#	ARTICLE	IF	CITATIONS
55	Red cell survival studies in hereditary spherocytosis. <i>Journal of Clinical Investigation</i> , 1970, 49, 666-672.	3.9	37
56	Activation of the P2Z/P2X7 Receptor in Human Lymphocytes Produces a Delayed Permeability Lesion: Involvement of Phospholipase D. <i>Archives of Biochemistry and Biophysics</i> , 1999, 362, 197-202.	1.4	36
57	Dysregulation of the inflammatory response to the parasite, <i>Toxoplasma gondii</i> , in P2X7 receptor-deficient mice. <i>International Journal for Parasitology</i> , 2011, 41, 301-308.	1.3	35
58	Loss of Function of P2X7 Receptor Scavenger Activity in Aging Mice. <i>American Journal of Pathology</i> , 2017, 187, 1670-1685.	1.9	34
59	Transendothelial migration of lymphocytes in chronic lymphocytic leukaemia is impaired and involved down-regulation of both L-selectin and CD23. <i>British Journal of Haematology</i> , 1999, 105, 181-189.	1.2	33
60	Genomics of Alzheimer's disease implicates the innate and adaptive immune systems. <i>Cellular and Molecular Life Sciences</i> , 2021, 78, 7397-7426.	2.4	32
61	P2Y11 receptor expression by human lymphocytes: evidence for two cAMP-linked purinoceptors. <i>European Journal of Pharmacology</i> , 2001, 426, 157-163.	1.7	31
62	Association of the 1513C polymorphism in the P2X7 gene with familial forms of chronic lymphocytic leukaemia. <i>British Journal of Haematology</i> , 2004, 125, 815-817.	1.2	31
63	Canine erythrocytes express the P2X ₇ receptor: greatly increased function compared with human erythrocytes. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2007, 293, R2090-R2098.	0.9	31
64	Insight into the pathogenesis of chronic lymphocytic leukemia (CLL) through analysis of IgVH gene usage and mutation status in familial CLL. <i>Blood</i> , 2008, 111, 5691-5693.	0.6	30
65	Saturation of intracellular cytosine arabinoside triphosphate accumulation in human leukemic blast cells. <i>Leukemia Research</i> , 1990, 14, 475-479.	0.4	29
66	Calcium ions, drug action and the red cell membrane. , 1982, 18, 271-292.		28
67	Flow cytometric quantitation of nucleoside transporter sites on human leukemic cells. <i>Cytometry</i> , 1993, 14, 32-38.	1.8	28
68	A quantitative method for routine measurement of cell surface P2X7 receptor function in leucocyte subsets by two-colour time-resolved flow cytometry. <i>Journal of Immunological Methods</i> , 2007, 325, 67-77.	0.6	27
69	Murine epidermal Langerhans cells and keratinocytes express functional P2X ₇ receptors. <i>Experimental Dermatology</i> , 2010, 19, e151-7.	1.4	27
70	Lack of a Functioning P2X7 Receptor Leads to Increased Susceptibility to Toxoplasmic Ileitis. <i>PLoS ONE</i> , 2015, 10, e0129048.	1.1	27
71	Specific detection of non-functional human P2X7 receptors in HEK293 cells and B-lymphocytes. <i>FEBS Letters</i> , 2003, 538, 159-162.	1.3	26
72	Analysis of a large multi-generational family provides insight into the genetics of chronic lymphocytic leukemia. <i>British Journal of Haematology</i> , 2008, 142, 238-245.	1.2	26

#	ARTICLE	IF	CITATIONS
73	Hereditary spherocytosis of man. Altered binding of cytoskeletal components to the erythrocyte membrane. <i>Biochemical Journal</i> , 1982, 201, 259-266.	1.7	25
74	Nucleoside transporters, bcl-2 and apoptosis in CLL cells exposed to nucleoside analogues <i>in vitro</i> . <i>European Journal of Haematology</i> , 1996, 56, 213-220.	1.1	25
75	TGF- β 1 prevents up-regulation of the P2X7 receptor by IFN- β and LPS in leukemic THP-1 monocytes. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2010, 1798, 2058-2066.	1.4	25
76	The P2X7 receptor mediates the uptake of organic cations in canine erythrocytes and mononuclear leukocytes: comparison to equivalent human cell types. <i>Purinergic Signalling</i> , 2009, 5, 385-394.	1.1	24
77	A quantitative method for measuring innate phagocytosis by human monocytes using real-time flow cytometry. <i>Cytometry Part A: the Journal of the International Society for Analytical Cytology</i> , 2014, 85, 313-321.	1.1	24
78	P2X7 Receptor-mediated Scavenger Activity of Mononuclear Phagocytes toward Non-opsonized Particles and Apoptotic Cells Is Inhibited by Serum Glycoproteins but Remains Active in Cerebrospinal Fluid. <i>Journal of Biological Chemistry</i> , 2012, 287, 17318-17330.	1.6	23
79	P2X7 receptor activation causes phosphatidylserine exposure in human erythrocytes. <i>Biochemical and Biophysical Research Communications</i> , 2007, 355, 169-173.	1.0	22
80	Activation of the erythroid K-Cl cotransporter Kcc1 enhances sickle cell disease pathology in a humanized mouse model. <i>Blood</i> , 2015, 126, 2863-2870.	0.6	21
81	Impaired Transendothelial Migration of B-CLL Lymphocytes: a Defect Linked to Low L-Selectin Expression. <i>Leukemia and Lymphoma</i> , 2001, 42, 5-12.	0.6	20
82	The scavenger activity of the human P2X7 receptor differs from P2X7 pore function by insensitivity to antagonists, genetic variation and sodium concentration: Relevance to inflammatory brain diseases. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018, 1864, 1051-1059.	1.8	20
83	Rottlerin inhibits P2X7 receptor-stimulated phospholipase D activity in chronic lymphocytic leukaemia B-lymphocytes. <i>Immunology and Cell Biology</i> , 2007, 85, 68-72.	1.0	19
84	The P2X7 receptor of CLL lymphocytes-a molecule with a split personality. <i>Lancet, The</i> , 2002, 360, 1898-1899.	6.3	18
85	Epistasis with HLA DR3 implicates the P2X7 receptor in the pathogenesis of primary Sjögren's syndrome. <i>Arthritis Research and Therapy</i> , 2013, 15, R71.	1.6	17
86	Non-synonymous polymorphisms in the P2RX4 are related to bone mineral density and osteoporosis risk in a cohort of Dutch fracture patients. <i>Purinergic Signalling</i> , 2013, 9, 123-130.	1.1	15
87	Functional significance of P2RX7 polymorphisms associated with affective mood disorders. <i>Journal of Psychiatric Research</i> , 2010, 44, 1116-1117.	1.5	14
88	NORMAL FLUIDITY OF RED CELL MEMBRANES IN HEREDITARY SPHEROCYTOSIS. <i>British Journal of Haematology</i> , 1980, 46, 299-301.	1.2	13
89	Nucleoside transport and cytosine arabinoside (araC) metabolism in human T lymphoblasts resistant to araC, thymidine and 6-methylmectapurine riboside. <i>European Journal of Cancer & Clinical Oncology</i> , 1985, 21, 1077-1082.	0.9	13
90	Inhibition of the human P2X7 receptor by a novel protein tyrosine kinase antagonist. <i>Biochemical and Biophysical Research Communications</i> , 2008, 365, 515-520.	1.0	13

#	ARTICLE	IF	CITATIONS
91	Flow cytometric studies of nucleoside transport regulation in single chromaffin cells. FEBS Letters, 1998, 422, 368-372.	1.3	11
92	A single nucleotide polymorphism haplotype promotes exon 7 and 8 skipping and disrupts receptor function. FASEB Journal, 2020, 34, 3884-3901.	0.2	10
93	Deficits in Monocyte Function in Age Related Macular Degeneration: A Novel Systemic Change Associated With the Disease. Frontiers in Medicine, 2021, 8, 634177.	1.2	10
94	Transport of 2-deoxycoformycin in human leukemia and lymphoma cells. Biochemical Pharmacology, 1991, 42, 708-710.	2.0	9
95	Inherited red cell dehydration: a hemolytic syndrome in search of a name. Pathology, 1984, 16, 115-116.	0.3	8
96	Seeking the nucleoside transporter. Nature Medicine, 1997, 3, 25-26.	15.2	8
97	Treatment of acute promyelocytic leukaemia relapsing after allogeneic bone marrow transplantation with all-trans-retinoic acid: suppression of the leukaemic clone. British Journal of Haematology, 1991, 79, 331-334.	1.2	7
98	Genetic polymorphisms of the human P2X7 receptor and relationship to function. Drug Development Research, 2001, 53, 72-76.	1.4	7
99	CXCR4 but not CXCR3 expression correlates with lymphocyte counts in B-cell chronic lymphocytic leukemia. Annals of Hematology, 2004, 83, 326-327.	0.8	7
100	An unusual variant of hereditary spherocytosis. American Journal of Medicine, 1970, 48, 63-71.	0.6	6
101	Molecular Stability of the Philly ($\pm 2^{235}$ (C1) Tyr \hat{a} ' Phe). Hemoglobin, 1981, 5, 177-190.	0.4	6
102	Strikingly Homologous Immunoglobulin Gene Rearrangements and Poor Outcome in VH3-21-Utilizing Chronic Lymphocytic Leukemia Independent of Geographical Origin and Mutational Status.. Blood, 2005, 106, 175-175.	0.6	6
103	Immune thrombocytopenia association with oralgold treatment. Arthritis and Rheumatism, 1988, 31, 299-300.	6.7	5
104	Flow cytometry identifies an early stage of platelet apoptosis produced by agonists of the P2X1 and P2X7 receptors. Platelets, 2022, 33, 621-631.	1.1	5
105	Identification of Leukocyte Surface P2X7 as a Biomarker Associated with Alzheimer's Disease. International Journal of Molecular Sciences, 2022, 23, 7867.	1.8	5
106	Identification of the promoter region of the P2RX4 gene. Molecular Biology Reports, 2010, 37, 3369-3376.	1.0	4
107	Regulation of the Acute Sickness Response by the P2RX7 Receptor. Journal of Infectious Diseases, 2021, 224, 914-920.	1.9	4
108	Idiopathic thrombocytopenic purpura in adults. Medical Journal of Australia, 1999, 170, 196-197.	0.8	2

#	ARTICLE	IF	CITATIONS
109	Should we delete the digraphs (â€œaeâ€•and â€œoeâ€•) from Australian medical writing?. Medical Journal of Australia, 1986, 144, 667-668.	0.8	1
110	Heme Biosynthesis and Its Disorders. , 2018, , 497-513.e6.		1
111	Assays to Measure Purinoceptor Pore Dilation. Methods in Molecular Biology, 2020, 2041, 323-334.	0.4	1
112	From Budapest to Bar Harbor. Blood, 2005, 106, 1145-1146.	0.6	0
113	Editorial. Purinergic Signalling, 2009, 5, 127-128.	1.1	0