

# Clara Dees

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/211797/publications.pdf>

Version: 2024-02-01

48  
papers

3,959  
citations

136885

32  
h-index

214721

47  
g-index

48  
all docs

48  
docs citations

48  
times ranked

5584  
citing authors

#	ARTICLE	IF	CITATIONS
1	Cellular and molecular mechanisms in fibrosis. <i>Experimental Dermatology</i> , 2021, 30, 121-131.	1.4	39
2	X-linked inhibitor of apoptosis protein (XIAP) inhibition in systemic sclerosis (SSc). <i>Annals of the Rheumatic Diseases</i> , 2021, 80, 1048-1056.	0.5	3
3	Targeting of canonical WNT signaling ameliorates experimental sclerodermatous chronic graft-versus-host disease. <i>Blood</i> , 2021, 137, 2403-2416.	0.6	11
4	Engrailed 1 coordinates cytoskeletal reorganization to induce myofibroblast differentiation. <i>Journal of Experimental Medicine</i> , 2021, 218, .	4.2	16
5	Fibroblast growth factor receptor 3 activates a network of profibrotic signaling pathways to promote fibrosis in systemic sclerosis. <i>Science Translational Medicine</i> , 2020, 12, .	5.8	26
6	PGC-1 $\beta$ regulates autophagy to promote fibroblast activation and tissue fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2020, 79, 1227-1233.	0.5	19
7	Translational engagement of lysophosphatidic acid receptor 1 in skin fibrosis: from dermal fibroblasts of patients with scleroderma to tight skin 1 mouse. <i>British Journal of Pharmacology</i> , 2020, 177, 4296-4309.	2.7	19
8	TGF- $\beta$ -induced epigenetic deregulation of SOCS3 facilitates STAT3 signaling to promote fibrosis. <i>Journal of Clinical Investigation</i> , 2020, 130, 2347-2363.	3.9	76
9	PU.1 controls fibroblast polarization and tissue fibrosis. <i>Nature</i> , 2019, 566, 344-349.	13.7	121
10	Poly(ADP-ribose) polymerase-1 regulates fibroblast activation in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2018, 77, 744-751.	0.5	36
11	Protein kinases G are essential downstream mediators of the antifibrotic effects of sGC stimulators. <i>Annals of the Rheumatic Diseases</i> , 2018, 77, 459-459.	0.5	33
12	The histone demethylase Jumonji domain-containing protein 3 (JMJD3) regulates fibroblast activation in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2018, 77, 150-158.	0.5	51
13	The tyrosine phosphatase SHP2 controls TGF- $\beta$ -induced STAT3 signaling to regulate fibroblast activation and fibrosis. <i>Nature Communications</i> , 2018, 9, 3259.	5.8	89
14	The transcription factor GLI2 as a downstream mediator of transforming growth factor- $\beta$ -induced fibroblast activation in SSc. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 756-764.	0.5	53
15	Composition of TWIST1 dimers regulates fibroblast activation and tissue fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 244-251.	0.5	28
16	JAK1-dependent transphosphorylation of JAK2 limits the antifibrotic effects of selective JAK2 inhibitors on long-term treatment. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 1467-1475.	0.5	41
17	Nintedanib inhibits macrophage activation and ameliorates vascular and fibrotic manifestations in the Fra2 mouse model of systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 1941-1948.	0.5	149
18	Inhibition of Notch1 promotes hedgehog signalling in a HES1-dependent manner in chondrocytes and exacerbates experimental osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 2037-2044.	0.5	29

#	ARTICLE	IF	CITATIONS
19	Tribbles homologue 3 stimulates canonical TGF- $\beta$ 2 signalling to regulate fibroblast activation and tissue fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 609-616.	0.5	38
20	Activating transcription factor 3 regulates canonical TGF $\beta$ 2 signalling in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 586-592.	0.5	28
21	Type 2 innate lymphoid cell counts are increased in patients with systemic sclerosis and correlate with the extent of fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 623-626.	0.5	78
22	Inhibition of casein kinase II reduces TGF $\beta$ 2 induced fibroblast activation and ameliorates experimental fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 936-943.	0.5	45
23	Stimulation of the soluble guanylate cyclase (sGC) inhibits fibrosis by blocking non-canonical TGF $\beta$ 2 signalling. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 1408-1416.	0.5	92
24	Orphan nuclear receptor NR4A1 regulates transforming growth factor- $\beta$ 2 signaling and fibrosis. <i>Nature Medicine</i> , 2015, 21, 150-158.	15.2	267
25	From pathogenesis to therapy – Perspective on treatment strategies in fibrotic diseases. <i>Pharmacological Research</i> , 2015, 100, 93-100.	3.1	17
26	Stimulators of soluble guanylate cyclase (sGC) inhibit experimental skin fibrosis of different aetiologies. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 1621-1625.	0.5	60
27	Activation of liver X receptors inhibits experimental fibrosis by interfering with interleukin-6 release from macrophages. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 1317-1324.	0.5	28
28	S100A4 amplifies TGF- $\beta$ 2-induced fibroblast activation in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 1748-1755.	0.5	52
29	Vitamin D receptor regulates TGF- $\beta$ 2 signalling in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, e20-e20.	0.5	111
30	The Wnt antagonists DKK1 and SFRP1 are downregulated by promoter hypermethylation in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 1232-1239.	0.5	166
31	Morphogen pathways as molecular targets for the treatment of fibrosis in systemic sclerosis. <i>Archives of Dermatological Research</i> , 2013, 305, 1-8.	1.1	25
32	Blockade of canonical Wnt signalling ameliorates experimental dermal fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, 1255-1258.	0.5	98
33	Canonical Wnt signalling as a key regulator of fibrogenesis – implications for targeted therapies?. <i>Experimental Dermatology</i> , 2013, 22, 710-713.	1.4	49
34	Levels of target activation predict antifibrotic responses to tyrosine kinase inhibitors. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, 2039-2046.	0.5	20
35	A8.3 – Deficit of S100A4 Prevents Joint Destruction and Systemic Bone Loss in hTNF $\alpha$ Mouse Model. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A58.1-A58.	0.5	0
36	Inhibition of hedgehog signalling prevents experimental fibrosis and induces regression of established fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 785-789.	0.5	73

#	ARTICLE	IF	CITATIONS
37	Stimulation of soluble guanylate cyclase reduces experimental dermal fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 1019-1026.	0.5	74
38	Inhibition of hedgehog signaling for the treatment of murine sclerodermatous chronic graft-versus-host disease. <i>Blood</i> , 2012, 120, 2909-2917.	0.6	53
39	JAK $\epsilon$ as a novel mediator of the profibrotic effects of transforming growth factor $\beta$ 2 in systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2012, 64, 3006-3015.	6.7	115
40	Combined Inhibition of c-Abl and PDGF Receptors for Prevention and Treatment of Murine Sclerodermatous Chronic Graft-versus-Host Disease. <i>American Journal of Pathology</i> , 2012, 181, 1672-1680.	1.9	28
41	Activation of canonical Wnt signalling is required for TGF $\beta$ 2-mediated fibrosis. <i>Nature Communications</i> , 2012, 3, 735.	5.8	649
42	Hedgehog signaling controls fibroblast activation and tissue fibrosis in systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2012, 64, 2724-2733.	6.7	133
43	$\beta$ 2-catenin is a central mediator of pro-fibrotic Wnt signaling in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 761-767.	0.5	174
44	Inhibition of glycogen synthase kinase 3 $\alpha$ induces dermal fibrosis by activation of the canonical Wnt pathway. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, 2191-2198.	0.5	96
45	Notch signalling regulates fibroblast activation and collagen release in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, 1304-1310.	0.5	116
46	Inhibition of Notch signaling prevents experimental fibrosis and induces regression of established fibrosis. <i>Arthritis and Rheumatism</i> , 2011, 63, 1396-1404.	6.7	107
47	Platelet-derived serotonin links vascular disease and tissue fibrosis. <i>Journal of Experimental Medicine</i> , 2011, 208, 961-972.	4.2	222
48	The cannabinoid receptor CB2 exerts antifibrotic effects in experimental dermal fibrosis. <i>Arthritis and Rheumatism</i> , 2009, 60, 1129-1136.	6.7	106