## Iñaki Alvarez

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2147523/publications.pdf

Version: 2024-02-01

		430874	526287
30	747	18	27
papers	citations	h-index	g-index
			0.1.1
30	30	30	911
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Purification of HLA Immunopeptidomes from Human Thymus. Methods in Molecular Biology, 2022, 2420, 127-136.	0.9	О
2	Global Proteomic and Methylome Analysis in Human Induced Pluripotent Stem Cells Reveals Overexpression of a Human TLR3 Affecting Proper Innate Immune Response Signaling. Stem Cells, 2019, 37, 476-488.	3.2	7
3	<scp>PRBAM</scp> : a new tool to analyze the <scp>MHC</scp> class I and <scp>HLA</scp> â€ <scp>DR</scp> anchor motifs. Immunology, 2019, 156, 187-198.	4.4	1
4	Human Leukocyte Antigen (HLA)-DRB1*15:01 and HLA-DRB5*01:01 Present Complementary Peptide Repertoires. Frontiers in Immunology, 2017, 8, 984.	4.8	25
5	A Comparative Analysis of the Peptide Repertoires of HLA–DR Molecules Differentially Associated With Rheumatoid Arthritis. Arthritis and Rheumatology, 2016, 68, 2412-2421.	5.6	10
6	Comparative Analysis of the Endogenous Peptidomes Displayed by HLA-B*27 and Mamu-B*08: Two MHC Class I Alleles Associated with Elite Control of HIV/SIV Infection. Journal of Proteome Research, 2016, 15, 1059-1069.	3.7	16
7	Central T cell tolerance: Identification of tissue-restricted autoantigens in the thymus HLA-DR peptidome. Journal of Autoimmunity, 2015, 60, 12-19.	6.5	27
8	The Power and the Promise of Cell Reprogramming: Personalized Autologous Body Organ and Cell Transplantation. Journal of Clinical Medicine, 2014, 3, 373-387.	2.4	8
9	Peptides presented by HLA class I molecules in the human thymus. Journal of Proteomics, 2013, 94, 23-36.	2.4	14
10	The Repertoires of Peptides Presented by MHC-II in the Thymus and in Peripheral Tissue: A Clue for Autoimmunity?. Frontiers in Immunology, 2013, 4, 442.	4.8	22
11	Composition of the HLAâ€DRâ€associated human thymus peptidome. European Journal of Immunology, 2013, 43, 2273-2282.	2.9	38
12	The peptide-binding motif of HLA-DR8 shares important structural features with other type $1$ diabetes-associated alleles. Genes and Immunity, $2011$ , $12$ , $504-512$ .	4.1	19
13	Increased Apoptosis after Autoimmune Regulator Expression in Epithelial Cells Revealed by a Combined Quantitative Proteomics Approach. Journal of Proteome Research, 2010, 9, 2600-2609.	3.7	21
14	The rheumatoid arthritis–associated allele HLA–DR10 ( <i>DRB1*1001</i> ) shares part of its repertoire with HLA–DR1 ( <i>DRB1*0101</i> ) and HLA–DR4 ( <i>DRB*0401</i> ). Arthritis and Rheumatism, 2008, 58, 1630-1639.	6.7	34
15	Chapter 6 Peptides Presented In Vivo by HLA-DR in Thyroid Autoimmunity. Advances in Immunology, 2008, 99, 165-209.	2.2	13
16	Thyroglobulin Peptides Associate In Vivo to HLA-DR in Autoimmune Thyroid Glands. Journal of Immunology, 2008, 181, 795-807.	0.8	48
17	Analysis of the HLA class I associated peptide repertoire in a hepatocellular carcinoma cell line reveals tumor-specific peptides as putative targets for immunotherapy. Proteomics - Clinical Applications, 2007, 1, 286-298.	1.6	7
18	Infection with Salmonella typhimurium has no effect on the composition and cleavage specificity of the 20S proteasome in human lymphoid cells. Immunology, 2007, 122, 131-139.	4.4	4

#	Article	IF	CITATIONS
19	Dissection of the HLA-DR4 Peptide Repertoire in Endocrine Epithelial Cells: Strong Influence of Invariant Chain and HLA-DM Expression on the Nature of Ligands. Journal of Immunology, 2004, 173, 1085-1093.	0.8	46
20	HLA-B27: a registry of constitutive peptide ligands. Tissue Antigens, 2004, 63, 424-445.	1.0	91
21	Species-specific Differences in Proteasomal Processing and Tapasin-mediated Loading Influence Peptide Presentation by HLA-B27 in Murine Cells. Journal of Biological Chemistry, 2003, 278, 46461-46472.	3.4	24
22	Molecular Mimicry of an HLA-B27-derived Ligand of Arthritis-linked Subtypes with Chlamydial Proteins. Journal of Biological Chemistry, 2002, 277, 37573-37581.	3.4	74
23	Large sharing of T-cell epitopes and natural ligands between HLA-B27 subtypes (B*2702 and B*2705) associated with spondyloarthritis. Tissue Antigens, 2001, 58, 351-362.	1.0	15
24	Minimal alterations in the HLA-B27-bound peptide repertoire induced upon infection of lymphoid cells withSalmonella typhimurium. Arthritis and Rheumatism, 2001, 44, 1677-1688.	6.7	20
25	The Cys-67 Residue of HLA-B27 Influences Cell Surface Stability, Peptide Specificity, and T-cell Antigen Presentation. Journal of Biological Chemistry, 2001, 276, 48740-48747.	3.4	38
26	Identification of Novel HLA-B27 Ligands Derived from Polymorphic Regions of Its Own or Other Class I Molecules Based on Direct Generation by 20 S Proteasome. Journal of Biological Chemistry, 2001, 276, 32729-32737.	3.4	23
27	HLA-B27 and immunogenetics of spondyloarthropathies. Current Opinion in Rheumatology, 2000, 12, 248-253.	4.3	31
28	An N-Acetylated Natural Ligand of Human Histocompatibility Leukocyte Antigen (Hla)-B39. Journal of Experimental Medicine, 2000, 191, 2083-2092.	8.5	22
29	Limited Diversity of Peptides Related to an Alloreactive T Cell Epitope in the HLA-B27-Bound Peptide Repertoire Results from Restrictions at Multiple Steps Along the Processing-Loading Pathway. Journal of Immunology, 2000, 164, 329-337.	0.8	40
30	A molecular insight on the association of HLA-B27 with spondyloarthropathies. Current Rheumatology Reports, 1999, 1, 78-85.	4.7	9