

Eliza Maria Xavier Freire

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2148752/publications.pdf>

Version: 2024-02-01

82

papers

822

citations

623734

14

h-index

580821

25

g-index

83

all docs

83

docs citations

83

times ranked

983

citing authors

#	ARTICLE	IF	CITATIONS
1	A “hotspot” within a hotspot: the reptiles of the Estação Ecológica and Área de Proteção Ambiental de Murici, Atlantic Forest of northeastern Brazil. <i>Biota Neotropica</i> , 2022, 22, .	0.5	3
2	Association of <i>Strymon serapio</i> (Lycaenidae: Theclinae) and <i>Castnia invaria</i> (Castniidae: Castniini) with the Habitat-Forming Bromeliad <i>Encholirium spectabile</i> (Bromeliaceae: Pitcairnioideae) in the Brazilian Semiarid Caatinga. <i>Journal of the Lepidopterists' Society</i> , 2021, 75, .	0.2	1
3	The rupicolous bromeliad (<i>Encholirium spectabile</i>) as a keystone species for Brazilian semiarid biodiversity. <i>Ecology</i> , 2021, 102, e03357.	3.2	4
4	Cytogenetic analysis of nuclear abnormalities in the erythrocytes of gecko lizards (<i>Phyllopezus</i>) Tj ETQq0 O O rgBT /Overlock 10 Tf 50 63 radioactivity. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2021, 865, 503346.	1.7	2
5	Lizards and bromeliads in the Neotropics: literature review and relevance of this association to conservation. <i>Symbiosis</i> , 2021, 84, 119-130.	2.3	2
6	The Key Role of a Thorny Rupicolous Bromeliad for Semiarid Biodiversity. <i>Bulletin of the Ecological Society of America</i> , 2021, 102, e01890.	0.2	0
7	Ethnozoology of snakebite victims in a risk area in Northeast Brazil. <i>Toxicon</i> , 2021, 201, 155-163.	1.6	5
8	Reproductive behavior of Spixâ€™s Whiptails in the wild: understanding the costs and benefits of mate-guarding. <i>Acta Ethologica</i> , 2021, 24, 41-51.	0.9	0
9	Lizards as model organisms to evaluate environmental contamination and biomonitoring. <i>Environmental Monitoring and Assessment</i> , 2020, 192, 454.	2.7	12
10	New record of <i>Hydrodynastes gigas</i> (Dum´ril, Bibron & Dum´ril, 1854) (Serpentes, Dipsadinae) in northeastern Brazil. <i>Check List</i> , 2020, 16, 457-460.	0.4	2
11	Análise de correlação entre as regiões administrativas de Natal (RN) baseada em indicadores de vulnerabilidade socioambiental. <i>Revista Ibero-americana De Ciências Ambientais</i> , 2020, 11, 445-461.	0.1	0
12	Snakebite accidents in Rio Grande do Norte state, Brazil: Epidemiology, health management and influence of the environmental scenario. <i>Tropical Medicine and International Health</i> , 2019, 24, 432-441.	2.3	20
13	Behavioral adaptations in <i>Ameivula ocellifera</i> (Squamata: Teiidae) in response to thermal environmental changes. <i>Phylomedusa</i> , 2019, 18, 225-240.	0.2	1
14	Richness and composition of anuran assemblages from an Amazonian savanna. <i>ZooKeys</i> , 2019, 843, 149-169.	1.1	3
15	New Morphological Data and Geographic Distribution of the Worm Lizard <i>Amphisbaena littoralis</i> (Squamata: Amphisbaenidae) in Northeastern Brazil. <i>South American Journal of Herpetology</i> , 2019, 14, 81.	0.5	0
16	Vulnerabilidade socioambiental: uma análise dos indicadores na cidade de Natal/RN. <i>Revista Ibero-americana De Ciências Ambientais</i> , 2019, 10, 323-339.	0.1	0
17	Floral visitors and potential pollinators of a rupicolous bromeliad (Pitcairnioideae) in the Brazilian semiarid. <i>Neotropical Biology and Conservation</i> , 2018, 13, .	0.9	6
18	Vulnerabilidade ambiental em curtumes artesanais: um estudo sobre o município de Cabaceiras/PB, Brasil. <i>Boletim Do Observatório Ambiental Alberto Ribeiro Lamego</i> , 2018, 12, 260-274.	0.0	0

#	ARTICLE	IF	CITATIONS
19	Percepção ambiental de agricultores de região semiárida sobre os anfíbios anuros e biocontrole de insetos pragas em sistemas irrigados e não irrigados, às margens do rio São Francisco, Brasil. <i>Gaia Scientia</i> , 2017, 11, .	0.0	0
20	Análise de políticas socioambientais implicadas no caso de deslizamento de terra no bairro Mâe Luiza em Natal/RN. <i>Gaia Scientia</i> , 2017, 11, .	0.0	0
21	First record of <i>Pseudopaludicola boliviensis</i> Parker, 1927 (Anura, Leptodactylidae, Leiuperinae) in the Brazilian state of Amapá, with comments on its advertisement call and distribution. <i>Check List</i> , 2016, 12, 1991.	0.4	4
22	Cultural keystone species of fauna as a method for assessing conservation priorities in a Protected Area of the Brazilian semiarid. <i>Biota Neotropica</i> , 2016, 16, .	1.0	11
23	Fauna used by rural communities surrounding the protected area of Chapada do Araripe, Brazil. <i>Journal of Ethnobiology and Ethnomedicine</i> , 2016, 12, 41.	2.6	2
24	Diet of <i>Amphisbaena heathi</i> Schmidt, 1936 (Squamata: Amphisbaenidae) from Remnants of the Atlantic Forest, Northeastern Brazil, with Notes on Distribution, Diagnosis and Conservation. <i>South American Journal of Herpetology</i> , 2016, 11, 12-17.	0.5	3
25	ENVIRONMENTAL PERCEPTION OF RURAL COMMUNITIES AND ANALYSIS OF LANDSCAPE: SUBSIDIES FOR PRIORITY AREA FOR CONSERVATION PROPOSITION IN THE RIO GRANDE DO NORTE SEMIARID, BRAZIL. RA'E GA - O Espaço Geográfico Em Análise, 2015, 34, 69.	0.1	0
26	New record and update on the geographic distribution of <i>Enyalius bilineatus</i> Duméril & Bibron, 1837 in Brazil. <i>Check List</i> , 2015, 11, 1785.	0.4	1
27	On the natural history of the Caatinga Horned Frog, <i>Ceratophrys joazeirensis</i> (Anura: Tij ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 422	0.2	
28	Ovarian follicular cycle of <i>Tropidurus hispidus</i> and <i>Tropidurus semitaeniatus</i> (Squamata: Tij ETQq0 0 0 rgBT /Overlock 10 Tf 50 382 Td (0.5	
29	Geographic distribution model for <i>Mabuya agmosticha</i> (Squamata: Scincidae) in northeastern Brazil. <i>Zoologia</i> , 2015, 32, 71-76.	0.5	8
30	Diet and Foraging Behavior of <i>Ameivula ocellifera</i> (Squamata: Teiidae) in the Brazilian Semiarid Caatinga. <i>Journal of Herpetology</i> , 2015, 49, 579-585.	0.5	11
31	Conhecimento ecológico local sobre o veado, <i>Mazama gouazoubira</i> (g. Fischer, 1814), por moradores do entorno de uma área protegida do semiárido brasileiro. <i>Brazilian Journal of Environmental Sciences (Online)</i> , 2015, , 78-95.	0.4	1
32	Conhecimento ecológico local sobre anfíbios anuros por agricultores em sistemas agrícolas de região semiárida brasileira. <i>Brazilian Journal of Environmental Sciences (Online)</i> , 2015, , 198-211.	0.4	3
33	Distribution extension and geographic distribution map of <i>Elachistocleis helianneae</i> (Anura: Tij ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 422	0.4	
34	Percepção ambiental como instrumento de participação social na proposição de área prioritária no semiárido. <i>INTERthesis</i> , 2014, 11, 147.	0.1	3
35	New record of <i>Cercosaura ocellata</i> Wagler, 1830 (Squamata, Gymnophthalmidae) in northeastern Brazil, with a distribution map for the species in South America. <i>Check List</i> , 2014, 10, 1531.	0.4	2
36	Molecular phylogeny, species limits, and biogeography of the Brazilian endemic lizard genus <i>Enyalius</i> (Squamata: Leiosauridae): An example of the historical relationship between Atlantic Forests and Amazonia. <i>Molecular Phylogenetics and Evolution</i> , 2014, 81, 137-146.	2.7	42

#	ARTICLE	IF	CITATIONS
37	Percepção ambiental como instrumento de participação social na proposição de área prioritária no semiárido. INTERthesis, 2014, 11, 147.	0.1	0
38	Ecology and diversity of a lizard community in the semiarid region of Brazil. Biota Neotropica, 2013, 13, 199-209.	1.0	15
39	New record and geographic distribution map of <i>Alexandresaurus camaean Rodrigues, Pellegrino, Dixo, Verdade, Pavan, Argolo and Sites Jr., 2007</i> (Squamata: Gymnophthalmidae) in northeastern Brazil. Check List, 2013, 9, 783.	0.4	1
40	Caatinga Revisited: Ecology and Conservation of an Important Seasonal Dry Forest. Scientific World Journal, The, 2012, 2012, 1-18.	2.1	170
41	Feeding Habits and Predator-Prey Size Relationships in the Whiptail Lizard <i>Cnemidophorus ocellifer</i> (Teiidae) in the Semiarid Region of Brazil. South American Journal of Herpetology, 2012, 7, 149-156.	0.5	13
42	Population Density, Habitat Selection and Conservation of <i>Coleodactylus natalensis</i> (Squamata:) Tj ETQq0 0 0 rgBT /Overlock 10 10 Journal of Herpetology, 2012, 7, 181-190.	0.5	4
43	Environmental perception and use of fauna from a Private Natural Heritage Reserve (RPPN) in Brazilian semiarid. Acta Scientiarum - Biological Sciences, 2012, 34, .	0.3	3
44	Feeding ecology of the pygmy gecko <i>Coleodactylus natalensis</i> (Squamata: Sphaerodactylidae) in the Brazilian Atlantic Forest. Zoologia, 2012, 29, 293-299.	0.5	1
45	Reproductive and fat body cycles of <i>Tropidurus hispidus</i> and <i>Tropidurus semitaeniatus</i> (Squamata,) Tj ETQq1 1 0.784314 rgBT /Overlock 1.2 18 307-320.	1.2	18
46	Uso do habitat, peródos diários de atividade e ecologia tática de Ameiva ameiva (Squamata: Teiidae) em uma área de caatinga do nordeste do Brasil.. Phylomedusa, 2012, 10, 165.	0.2	3
47	First record of <i>Colobosaura modesta</i> (Reinhardt and Lütken, 1862) (Squamata: Gymnophthalmidae) to the Cariri region, state of Ceará, Brazil, with a map of its geographical distribution. Check List, 2012, 8, 970.	0.4	6
48	New records and geographic distribution map of <i>Typhlops amoipira</i> Rodrigues and Juncá, 2002 (Typhlopidae) in the Brazilian Rainforest. Check List, 2012, 8, 1347.	0.4	5
49	Thermal ecology and thermoregulatory behavior of <i>Coleodactylus natalensis</i> (Squamata:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 693-700.	0.5	6
50	A percepção ambiental de uma comunidade da caatinga sobre o turismo: visões e perspectivas para o planejamento turístico com vistas a sustentabilidade. Sociedade & Natureza, 2011, 23, 467-482.	0.0	6
51	Mating behavior and female accompaniment in the whiptail lizard <i>Cnemidophorus ocellifer</i> (Squamata,) Tj ETQq1 1 0.784314 rgBT /Ove	1.0	10
52	Trophic ecology and foraging behavior of <i>Tropidurus hispidus</i> and <i>Tropidurus semitaeniatus</i> (Squamata, Tropiduridae) in a caatinga area of northeastern Brazil. Iheringia - Serie Zoologia, 2011, 101, 225-232.	0.5	22
53	Communal nests of <i>Hemidactylus mabouia</i> (Moreau de Jonnés, 1818) (Squamata: Gekkonidae) in a remnant of Atlantic Forest in northeastern Brazil doi: 10.5007/2175-7925.2010v23n3p231. Biotemas, 2010, 23, .	0.1	1
54	Abordagem etnobotânica sobre plantas medicinais citadas por populações do entorno de uma unidade de conservação da caatinga do Rio Grande do Norte, Brasil. Revista Brasileira De Plantas Medicinais, 2010, 12, 427-435.	0.3	20

#	ARTICLE	IF	CITATIONS
55	Perception and use of fauna resources in communities surrounding a conservation unit in northeast Brazil. <i>Acta Scientiarum - Biological Sciences</i> , 2010, 32, .	0.3	4
56	Nematode infection in the lizard <i>< i>Bogertia lutzae</i></i> (Loveridge, 1941) from the Atlantic forest in north-eastern Brazil. <i>Journal of Helminthology</i> , 2010, 84, 199-201.	1.0	11
57	Body Mass and the Energy Budget of Gravid Hawksbill Turtles (<i>Eretmochelys imbricata</i>) during the Nesting Season. <i>Journal of Herpetology</i> , 2010, 44, 352-359.	0.5	26
58	Diet and Foraging Behavior of Two Species of <i>Tropidurus</i> (Squamata, Tropiduridae) in the Caatinga of Northeastern Brazil. <i>South American Journal of Herpetology</i> , 2010, 5, 35-44.	0.5	25
59	Ventral Colored Patches in <i>Tropidurus semitaeniatus</i> (Squamata, Tropiduridae): Sexual Dimorphism and Association with Reproductive Cycle. <i>Journal of Herpetology</i> , 2010, 44, 177-182.	0.5	15
60	Forced copulation attempt in the Blue-tailed Lizard, <i>Micrablepharus maximiliani</i> (Reinhardt & Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Neotropica, 2010, 10, 347-350.	1.0	6
61	Reptilia, Squamata, Leiosauridae, <i>Enyalius bibronii</i> Boulenger, 1885: distribution extension and geographic distribution map. <i>Check List</i> , 2010, 6, 652.	0.4	0
62	Reptilia, Squamata, Polychrotidae, <i>Anolis fuscoauratus</i> D'Orbigny, 1837: distribution extension for the state of Rio Grande do Norte, Brazil. <i>Check List</i> , 2010, 6, 681.	0.4	1
63	A new species of <i>Dendrophidion</i> (Serpentes: Colubridae) from the Atlantic Rain Forest of Northeastern Brazil. <i>Zootaxa</i> , 2010, 2719, 62.	0.5	10
64	The effects of seasonality on the foraging behavior of <i>Tropidurus hispidus</i> and <i>Tropidurus semitaeniatus</i> (Squamata: Tropiduridae) living in sympatry in the Caatinga of northeastern Brazil. <i>Zoologia</i> , 2009, 26, 581-585.	0.5	10
65	Conceitos, percepções e estratégias para conservação de uma estação ecológica da Caatinga nordestina por populações do seu entorno. <i>Sociedade & Natureza</i> , 2009, 21, 23-37.	0.0	8
66	Pentastomid, <i>Raillietiella mottae</i> Almeida, Freire and Lopes, 2008, infecting lizards in an area of caatinga, northeast, Brazil. <i>Brazilian Journal of Biology</i> , 2008, 68, 427-431.	0.9	7
67	Pentastomid infection in <i>Philodryas nattereri</i> Steindachner, 1870 and <i>Oxybelis aeneus</i> (Wagler, 1824) (Squamata: Colubridae) in a caatinga of northeastern Brazil. <i>Brazilian Journal of Biology</i> , 2008, 68, 193-197.	0.9	13
68	A new species of pentastomida infecting <i>Tropidurus hispidus</i> (Squamata: Tropiduridae) from Caatinga in Northeastern Brazil. <i>Brazilian Journal of Biology</i> , 2008, 68, 199-203.	0.9	12
69	Pentastomids infecting an invader lizard, <i>Hemidactylus mabouia</i> (Gekkonidae) in northeastern Brazil. <i>Brazilian Journal of Biology</i> , 2008, 68, 611-615.	0.9	14
70	Prevalence and intensity of pentastomid infection in two species of snakes from northeastern Brazil. <i>Brazilian Journal of Biology</i> , 2007, 67, 759-763.	0.9	12
71	The alien and native pentastomids fauna of an exotic lizard population from Brazilian Northeast. <i>Parasitology Research</i> , 2007, 101, 627-628.	1.6	13
72	OS LAGARTOS GIMNOFTALMÍDEOS (SQUAMATA: GYMNOPHTHALMIDAE) DO CARIRI PARAIBANO E DO SERIDÓ DO RIO GRANDE DO NORTE, NORDESTE DO BRASIL: CONSIDERAÇÕES ACERCA DA DISTRIBUIÇÃO GEOGRÁFICA E ECOLOGIA. <i>Oecologia Brasiliensis</i> , 2007, 11, 365-382.	0.5	10

#	ARTICLE	IF	CITATIONS
73	A new species of <i>Liophlops</i> (Serpentes: Anomalepididae) from the Atlantic Rain Forest of Northeastern Brazil. <i>Zootaxa</i> , 2007, 1393, 19.	0.5	2
74	Phylogenetic relationships of a new genus and species of microteiid lizard from the Atlantic forest of north-eastern Brazil (Squamata, Gymnophthalmidae). <i>Zoological Journal of the Linnean Society</i> , 2005, 144, 543-557.	2.3	42
75	TWO NEW SPECIES OF <i>PHYLLODYTES</i> (ANURA: HYLIDAE) FROM THE STATE OF ALAGOAS, NORTHEASTERN BRAZIL. <i>Herpetologica</i> , 2003, 59, 235-246.	0.4	44
76	Occurrence of the genus <i>Chiasmocleis</i> (Anura: Microhylidae) in the State of Alagoas, north-eastern Brazil, with a description of a new species. <i>Journal of Zoology</i> , 1999, 249, 123-126.	1.7	19
77	Occurrence of the genus <i>Chiasmocleis</i> (Anura: Microhylidae) in the State of Alagoas, north-eastern Brazil, with a description of a new species. <i>Journal of Zoology</i> , 1999, 249, 123-126.	1.7	0
78	Estudo ecológico e zoogeográfico sobre a fauna de lagartos (Sauria) das dunas de Natal, Rio Grande do Norte e da Restinga de Ponta de Campina, Cabedelo, Paraíba, Brasil. <i>Revista Brasileira De Zoologia</i> , 1996, 13, 903-921.	0.5	38
79	Obstacles to the effectiveness of Protected Areas: Mata da Pipa State Park, Tibau do Sul – RN – Brazil. <i>Desenvolvimento E Meio Ambiente</i> , 0, 46, .	0.0	1
80	Demystifying ophidism: bridging school and society to develop educational resources. <i>Ambiente & Sociedade</i> , 0, 24, .	0.5	2
81	Living among thorns: herpetofaunal community (Anura and Squamata) associated to the rupicolous bromeliad <i>Encholirium spectabile</i> (Pitcairnioideae) in the Brazilian semi-arid Caatinga. <i>Zoologia</i> , 0, 37, 1-12.	0.5	6
82	ENVIRONMENTAL DEGRADATION IN SEMIARID BRASILIAN: AN APPLICATION OF SUSTAINABILITY INDICATORS IN MUNICIPALITIES WITH CONSERVATION UNITS. <i>Holos</i> , 0, 1, 182-201.	0.0	1