
## Mitsutoshi Yoneyama

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2160871/publications.pdf Version: 2024-02-01



| #  | Article   | lF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Regulation of RIC-I-like receptor-mediated signaling: interaction between host and viral factors.<br>Cellular and Molecular Immunology, 2021, 18, 539-555.  | 4.8 | 179       |
| 2  | Identification of a new autoinhibitory domain of interferon-beta promoter stimulator-1 (IPS-1) for the<br>tight regulation of oligomerization-driven signal activation. Biochemical and Biophysical Research<br>Communications, 2019, 517, 662-669. | 1.0 | 3         |
| 3  | Virus Sensor RIG-I Represses RNA Interference by Interacting with TRBP through LGP2 in Mammalian<br>Cells. Genes, 2018, 9, 511.   | 1.0 | 16        |
| 4  | Regulation of antiviral innate immune signaling by stress-induced RNA granules. Journal of Biochemistry, 2016, 159, mvv122.   | 0.9 | 43        |
| 5  | Leader-Containing Uncapped Viral Transcript Activates RIG-I in Antiviral Stress Granules. PLoS<br>Pathogens, 2016, 12, e1005444.  | 2.1 | 68        |
| 6  | ID: 70. Cytokine, 2015, 76, 78.   | 1.4 | 0         |
| 7  | Viral RNA detection by RIG-I-like receptors. Current Opinion in Immunology, 2015, 32, 48-53.  | 2.4 | 371       |
| 8  | The ASK family kinases differentially mediate induction of type I interferon and apoptosis during the antiviral response. Science Signaling, 2015, 8, ra78.   | 1.6 | 29        |
| 9  | DHX36 Enhances RIG-I Signaling by Facilitating PKR-Mediated Antiviral Stress Granule Formation. PLoS Pathogens, 2014, 10, e1004012.   | 2.1 | 129       |
| 10 | A Novel Function of Human Pumilio Proteins in Cytoplasmic Sensing of Viral Infection. PLoS<br>Pathogens, 2014, 10, e1004417.  | 2.1 | 51        |
| 11 | Antiviral innate immunity and stress granule responses. Trends in Immunology, 2014, 35, 420-428.  | 2.9 | 192       |
| 12 | Dectin-2 Is a Direct Receptor for Mannose-Capped Lipoarabinomannan of Mycobacteria. Immunity, 2014,<br>41, 402-413.   | 6.6 | 243       |
| 13 | Encephalomyocarditis Virus Disrupts Stress Granules, the Critical Platform for Triggering Antiviral<br>Innate Immune Responses. Journal of Virology, 2013, 87, 9511-9522.   | 1.5 | 127       |
| 14 | Hepatitis C virus NS4B protein targets STING and abrogates RIG-I-mediated type I interferon-dependent innate immunity. Hepatology, 2013, 57, 46-58.   | 3.6 | 127       |
| 15 | Refeeding with a standard diet after a 48-h fast elicits an inflammatory response in the mouse liver.<br>Journal of Nutritional Biochemistry, 2013, 24, 1314-1323.  | 1.9 | 5         |
| 16 | 192. Cytokine, 2013, 63, 288.   | 1.4 | 0         |
| 17 | Virus-induced expression of retinoic acid inducible gene-I and melanoma differentiation-associated gene 5 in the cochlear sensory epithelium. Microbes and Infection, 2013, 15, 592-598.  | 1.0 | 10        |
| 18 | Lymphocyte–stromal cell interaction induces IL-7 expression by interferon regulatory factors.<br>Molecular Immunology, 2013, 54, 378-385.   | 1.0 | 4         |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Functional Characterization of Domains of IPS-1 Using an Inducible Oligomerization System. PLoS ONE, 2013, 8, e53578.   | 1.1 | 22        |
| 20 | Impairment of interferon regulatory factor-3 activation by hepatitis C virus core protein basic amino acid region 1. Biochemical and Biophysical Research Communications, 2012, 428, 494-499.   | 1.0 | 14        |
| 21 | Foreign RNA Induces the Degradation of Mitochondrial Antiviral Signaling Protein (MAVS): The Role of Intracellular Antiviral Factors. PLoS ONE, 2012, 7, e45136.  | 1.1 | 11        |
| 22 | Critical Role of an Antiviral Stress Granule Containing RIG-I and PKR in Viral Detection and Innate<br>Immunity. PLoS ONE, 2012, 7, e43031.   | 1.1 | 294       |
| 23 | Retinoic Acid-Inducible Gene-I-Like Receptors. Journal of Interferon and Cytokine Research, 2011, 31, 27-31.  | 0.5 | 79        |
| 24 | 55 Amino acid linker between helicase and carboxyl terminal domains of RIG-I functions as a critical repression domain and determines inter-domain conformation. Biochemical and Biophysical Research Communications, 2011, 415, 75-81. | 1.0 | 24        |
| 25 | Retinoic Acid-inducible Gene I-inducible miR-23b Inhibits Infections by Minor Group Rhinoviruses through Down-regulation of the Very Low Density Lipoprotein Receptor. Journal of Biological Chemistry, 2011, 286, 26210-26219.         | 1.6 | 45        |
| 26 | Recognition of viral nucleic acids in innate immunity. Reviews in Medical Virology, 2010, 20, 4-22.   | 3.9 | 265       |
| 27 | Ser386 phosphorylation of transcription factor IRFâ€3 induces dimerization and association with CBP/p300 without overall conformational change. Genes To Cells, 2010, 15, 901-910.  | 0.5 | 55        |
| 28 | LGP2 is a positive regulator of RIC-l– and MDA5-mediated antiviral responses. Proceedings of the National Academy of Sciences of the United States of America, 2010, 107, 1512-1517.  | 3.3 | 540       |
| 29 | Virus-Infection or 5′ppp-RNA Activates Antiviral Signal through Redistribution of IPS-1 Mediated by MFN1. PLoS Pathogens, 2010, 6, e1001012.  | 2.1 | 150       |
| 30 | Identification of Loss of Function Mutations in Human Genes Encoding RIG-I and MDA5. Journal of<br>Biological Chemistry, 2009, 284, 13348-13354.  | 1.6 | 130       |
| 31 | Solution Structures of Cytosolic RNA Sensor MDA5 and LGP2 C-terminal Domains. Journal of Biological Chemistry, 2009, 284, 17465-17474.  | 1.6 | 170       |
| 32 | RNA recognition and signal transduction by RIGâ€ <del>l</del> â€ <del>l</del> ike receptors. Immunological Reviews, 2009, 227,<br>54-65.  | 2.8 | 525       |
| 33 | Mitochondrial dynamics and innate antiviral responses regulated by RIG-I-like receptor. Cytokine, 2009, 48, 133.  | 1.4 | 0         |
| 34 | Cytoplasmic recognition of RNAâ <sup>~</sup> †. Advanced Drug Delivery Reviews, 2008, 60, 841-846.  | 6.6 | 47        |
| 35 | Nonself RNA-Sensing Mechanism of RIG-I Helicase and Activation of Antiviral Immune Responses.<br>Molecular Cell, 2008, 29, 428-440.   | 4.5 | 416       |
| 36 | Structural Mechanism of RNA Recognition by the RIG-I-like Receptors. Immunity, 2008, 29, 178-181.   | 6.6 | 226       |

| #  | Article  | IF   | CITATIONS |
|----|--|------|-----------|
| 37 | Function of RIC-I-like Receptors in Antiviral Innate Immunity. Journal of Biological Chemistry, 2007, 282, 15315-15318.  | 1.6  | 258       |
| 38 | Viral Infections Activate Types I and III Interferon Genes through a Common Mechanism. Journal of<br>Biological Chemistry, 2007, 282, 7576-7581.   | 1.6  | 300       |
| 39 | Hepatitis C virus non-structural proteins responsible for suppression of the RIG-I/Cardif-induced interferon response. Journal of General Virology, 2007, 88, 3323-3333.   | 1.3  | 34        |
| 40 | RIG-I family RNA helicases: Cytoplasmic sensor for antiviral innate immunity. Cytokine and Growth<br>Factor Reviews, 2007, 18, 545-551.  | 3.2  | 126       |
| 41 | Triggering antiviral response by RIG-I-related RNA helicases. Biochimie, 2007, 89, 754-760.  | 1.3  | 67        |
| 42 | Cytoplasmic double-stranded DNA sensor. Nature Immunology, 2007, 8, 907-908.   | 7.0  | 9         |
| 43 | Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses. Nature, 2006, 441, 101-105.  | 13.7 | 3,292     |
| 44 | Role of the Alpha/Beta Interferon Response in the Acquisition of Susceptibility to Poliovirus by Kidney<br>Cells in Culture. Journal of Virology, 2006, 80, 4313-4325.   | 1.5  | 28        |
| 45 | Viral and therapeutic control of IFN-beta promoter stimulator 1 during hepatitis C virus infection.<br>Proceedings of the National Academy of Sciences of the United States of America, 2006, 103, 6001-6006.                | 3.3  | 394       |
| 46 | Control of antiviral defenses through hepatitis C virus disruption of retinoic acid-inducible gene-I<br>signaling. Proceedings of the National Academy of Sciences of the United States of America, 2005, 102,<br>2986-2991. | 3.3  | 506       |
| 47 | The Alpha/Beta Interferon Response Controls Tissue Tropism and Pathogenicity of Poliovirus. Journal of Virology, 2005, 79, 4460-4469.  | 1.5  | 210       |
| 48 | Inhibition of RIG-I-Dependent Signaling to the Interferon Pathway during Hepatitis C Virus Expression<br>and Restoration of Signaling by IKKε. Journal of Virology, 2005, 79, 3969-3978.                                     | 1.5  | 169       |
| 49 | Shared and Unique Functions of the DExD/H-Box Helicases RIG-I, MDA5, and LGP2 in Antiviral Innate<br>Immunity. Journal of Immunology, 2005, 175, 2851-2858.  | 0.4  | 1,438     |
| 50 | Regulating Intracellular Antiviral Defense and Permissiveness to Hepatitis C Virus RNA Replication through a Cellular RNA Helicase, RIG-I. Journal of Virology, 2005, 79, 2689-2699.   | 1.5  | 830       |
| 51 | Cell Type-Specific Involvement of RIG-I in Antiviral Response. Immunity, 2005, 23, 19-28.  | 6.6  | 1,221     |
| 52 | Identification of Ser-386 of Interferon Regulatory Factor 3 as Critical Target for Inducible<br>Phosphorylation That Determines Activation. Journal of Biological Chemistry, 2004, 279, 9698-9702.                           | 1.6  | 182       |
| 53 | The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. Nature Immunology, 2004, 5, 730-737.   | 7.0  | 3,433     |
| 54 | Direct Involvement of CREB-binding Protein/p300 in Sequence-specific DNA Binding of Virus-activated<br>Interferon Regulatory Factor-3 Holocomplex. Journal of Biological Chemistry, 2002, 277, 22304-22313.                  | 1.6  | 94        |

| #  | Article  | IF   | CITATIONS |
|----|--|------|-----------|
| 55 | Review: Control of IRF-3 Activation by Phosphorylation. Journal of Interferon and Cytokine Research, 2002, 22, 73-76.  | 0.5  | 150       |
| 56 | Involvement of TIRAP/MAL in signaling for the activation of interferon regulatory factor 3 by lipopolysaccharide. FEBS Letters, 2002, 517, 251-256.  | 1.3  | 44        |
| 57 | PACT, a Double-Stranded RNA Binding Protein Acts as a Positive Regulator for Type I Interferon Gene<br>Induced by Newcastle Disease Virus. Biochemical and Biophysical Research Communications, 2001, 282,<br>515-523. | 1.0  | 43        |
| 58 | Induction of IRF-3/-7 kinase and NF-κB in response to double-stranded RNA and virus infection: common<br>and unique pathways. Genes To Cells, 2001, 6, 375-388.  | 0.5  | 242       |
| 59 | Direct triggering of the type I interferon system by virus infection: activation of a transcription factor complex containing IRF-3 and CBP/p300. EMBO Journal, 1998, 17, 1087-1095.                                   | 3.5  | 735       |
| 60 | IL-2 and EGF receptors stimulate the hematopoietic cell cycle via different signaling pathways:<br>Demonstration of a novel role for c-myc. Cell, 1992, 70, 57-67.   | 13.5 | 250       |
| 61 | The human interleukin-2 receptor $\hat{l}^2$ -chain gene: genomic organization, promoter analysis and chromosomal assignment. Nucleic Acids Research, 1990, 18, 3697-3703.   | 6.5  | 91        |
| 62 | Involvement of a common transcription factor in the regulated expression of IL-2 and IL-2 receptor genes. International Immunology, 1989, 1, 43-49.  | 1.8  | 71        |