

# MaÅ,gorzata LeÅ›niewska

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2167982/publications.pdf>

Version: 2024-02-01

15  
papers

132  
citations

1478505

6  
h-index

1281871

11  
g-index

15  
all docs

15  
docs citations

15  
times ranked

108  
citing authors

#	ARTICLE	IF	CITATIONS
1	Trunk anomalies in the centipede <i>Stigmatogaster subterranea</i> provide insight into late-embryonic segmentation. <i>Arthropod Structure and Development</i> , 2009, 38, 417-426.	1.4	33
2	Phoretic relationships between Uropodina (Acari: Mesostigmata) and centipedes (Chilopoda) as an example of evolutionary adaptation of mites to temporary microhabitats. <i>European Journal of Entomology</i> , 2006, 103, 699-707.	1.2	25
3	Effects of short- and long-term exposure to cadmium on salivary glands and fat body of soil centipede <i>Lithobius forficatus</i> (Myriapoda, Chilopoda): Histology and ultrastructure. <i>Micron</i> , 2020, 137, 102915.	2.2	15
4	Phoretic relationships between uropodid mites (Acari: Mesostigmata) and centipedes (Chilopoda) in urban agglomeration areas. <i>International Journal of Acarology</i> , 2015, 41, 250-258.	0.7	13
5	Ovaries and testes of <i>Lithobius forficatus</i> (Myriapoda, Chilopoda) react differently to the presence of cadmium in the environment. <i>Scientific Reports</i> , 2022, 12, 6705.	3.3	10
6	Centipede (Chilopoda) richness and diversity in the Bug River valley (Eastern Poland). <i>ZooKeys</i> , 2015, 510, 125-139.	1.1	7
7	Hazards related to the presence of cadmium in food – Studies on the European soil centipede, <i>Lithobius forficatus</i> . <i>Science of the Total Environment</i> , 2022, 845, 157298.	8.0	7
8	Effect of urbanization on centipede (Chilopoda) diversity in the Wielkopolska-Kujawy Lowlands of western Poland. <i>Biologia (Poland)</i> , 2008, 63, 711-719.	1.5	6
9	Reproduction and shell growth in two clausiliids with different reproductive strategies. <i>Biologia (Poland)</i> , 2015, 70, 625-631.	1.5	5
10	Variability in the sculpture of the shell aperture of <i>Ruthenica filograna</i> (Rossmassler, 1836) (Gastropoda: Clausiliidae) in specimens from natural populations and from laboratory breeding. <i>Journal of Molluscan Studies</i> , 2008, 74, 183-189.	1.2	4
11	Population Genetic Structure of a Centipede Species with High Levels of Developmental Instability. <i>PLoS ONE</i> , 2015, 10, e0126245.	2.5	2
12	Centipede (Chilopoda) richness, diversity and community structure in the forest-steppe nature reserve –Bielinek– on the Odra River (NW Poland, Central Europe). <i>Biologia (Poland)</i> , 2016, 71, 1250-1265.	1.5	2
13	The centipede community of a beech forest in Magura National Park, Poland. <i>Fragmenta Faunistica</i> , 2003, 46, 109-119.	0.0	2
14	<i>Tygarrup javanicus</i> (Chilopoda, Geophilomorpha) – an exotic species that has reached Poland. <i>Polish Journal of Entomology</i> , 2020, 1, 52-58.	0.4	1
15	Impact of windstorm on a community of centipedes (Chilopoda) in a beech forest in Western Poland. <i>Biologia (Poland)</i> , 2018, 73, 165-173.	1.5	0