

Per Hansson

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2171729/per-hansson-publications-by-citations.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

118
papers

12,448
citations

47
h-index

111
g-index

126
ext. papers

14,497
ext. citations

5.3
avg, IF

6
L-index

#	Paper	IF	Citations
118	Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. <i>Lancet Neurology, The</i> , 2015 , 14, 162-73	24.1	1971
117	EFNS guidelines on the pharmacological treatment of neuropathic pain: 2010 revision. <i>European Journal of Neurology</i> , 2010 , 17, 1113-e88	6	1190
116	NeuPSIG guidelines on neuropathic pain assessment. <i>Pain</i> , 2011 , 152, 14-27	8	694
115	EFNS guidelines on pharmacological treatment of neuropathic pain. <i>European Journal of Neurology</i> , 2006 , 13, 1153-69	6	638
114	Neuropathic pain: an updated grading system for research and clinical practice. <i>Pain</i> , 2016 , 157, 1599-1686		536
113	EFNS guidelines on neurostimulation therapy for neuropathic pain. <i>European Journal of Neurology</i> , 2007 , 14, 952-70	6	476
112	Central representation of chronic ongoing neuropathic pain studied by positron emission tomography. <i>Pain</i> , 1995 , 63, 225-236	8	460
111	Modulatory influence on somatosensory perception from vibration and heterotopic noxious conditioning stimulation (HNCS) in fibromyalgia patients and healthy subjects. <i>Pain</i> , 1997 , 70, 41-51	8	401
110	Value of quantitative sensory testing in neurological and pain disorders: NeuPSIG consensus. <i>Pain</i> , 2013 , 154, 1807-1819	8	350
109	Recommendations on terminology and practice of psychophysical DNIC testing. <i>European Journal of Pain</i> , 2010 , 14, 339	3.7	326
108	Peripheral neuropathic pain: a mechanism-related organizing principle based on sensory profiles. <i>Pain</i> , 2017 , 158, 261-272	8	310
107	Intrapataminal infusion of nerve growth factor to support adrenal medullary autografts in Parkinson disease. One-year follow-up of first clinical trial. <i>Archives of Neurology</i> , 1991 , 48, 373-81		270
106	Sensory dysfunction in fibromyalgia patients with implications for pathogenic mechanisms. <i>Pain</i> , 1996 , 68, 375-83	8	256
105	Recommendations on practice of conditioned pain modulation (CPM) testing. <i>European Journal of Pain</i> , 2015 , 19, 805-6	3.7	239
104	The IASP classification of chronic pain for ICD-11: chronic neuropathic pain. <i>Pain</i> , 2019 , 160, 53-59	8	228
103	Usefulness and limitations of quantitative sensory testing: clinical and research application in neuropathic pain states. <i>Pain</i> , 2007 , 129, 256-259	8	221
102	Patient phenotyping in clinical trials of chronic pain treatments: IMMPACT recommendations. <i>Pain</i> , 2016 , 157, 1851-1871	8	178

101	Modulation of pressure pain thresholds during and following isometric contraction in patients with fibromyalgia and in healthy controls. <i>Pain</i> , 1996 , 64, 415-423	8	168
100	Assessment of neuropathic pain in primary care. <i>American Journal of Medicine</i> , 2009 , 122, S13-21	2.4	150
99	Clinical and pre-clinical pain assessment: are we measuring the same thing?. <i>Pain</i> , 2008 , 135, 7-10	8	150
98	Somatosensory perception and function of diffuse noxious inhibitory controls (DNIC) in patients suffering from rheumatoid arthritis. <i>European Journal of Pain</i> , 2002 , 6, 161-76	3.7	131
97	Pressure pain thresholds in different tissues in one body region. The influence of skin sensitivity in pressure algometry. <i>Journal of Rehabilitation Medicine</i> , 1999 , 31, 89-93		126
96	Neuropathic pain: clinical characteristics and diagnostic workup. <i>European Journal of Pain</i> , 2002 , 6 Suppl A, 47-50	3.7	106
95	Increased pressure pain sensibility in fibromyalgia patients is located deep to the skin but not restricted to muscle tissue. <i>Pain</i> , 1995 , 63, 335-339	8	100
94	Stratifying patients with peripheral neuropathic pain based on sensory profiles: algorithm and sample size recommendations. <i>Pain</i> , 2017 , 158, 1446-1455	8	94
93	Difficulties in stratifying neuropathic pain by mechanisms. <i>European Journal of Pain</i> , 2003 , 7, 353-7	3.7	91
92	Systemic Adenosine Infusion Alleviates Spontaneous and Stimulus Evoked Pain in Patients with Peripheral Neuropathic Pain. <i>Anesthesia and Analgesia</i> , 1995 , 81, 713-717	3.9	91
91	Context-dependent deactivation of the amygdala during pain. <i>Journal of Cognitive Neuroscience</i> , 2004 , 16, 1289-301	3.1	87
90	Somatosensory perception in a remote pain-free area and function of diffuse noxious inhibitory controls (DNIC) in patients suffering from long-term trapezius myalgia. <i>European Journal of Pain</i> , 2002 , 6, 149-59	3.7	83
89	Pharmacological treatment of peripheral neuropathic pain conditions based on shared commonalities despite multiple etiologies. <i>Pain</i> , 2005 , 113, 251-254	8	80
88	Concentrations of neuropeptides substance P, neurokinin A, calcitonin gene-related peptide, neuropeptide Y and vasoactive intestinal polypeptide in synovial fluid of the human temporomandibular joint. A correlation with symptoms, signs and arthroscopic findings. <i>International Journal of Oral and Maxillofacial Surgery</i> , 1991 , 20, 228-31	2.9	80
87	Systemic adenosine infusion: a new treatment modality to alleviate neuropathic pain. <i>Pain</i> , 1995 , 61, 155-158	8	76
86	Vibratory stimulation for the relief of pain of dental origin. <i>Pain</i> , 1981 , 10, 37-45	8	73
85	Brainstem involvement in the initial response to pain. <i>NeuroImage</i> , 2004 , 22, 995-1005	7.9	69
84	Peripheral alpha-adrenoreceptors are involved in the development of capsaicin induced ongoing and stimulus evoked pain in humans. <i>Pain</i> , 1997 , 69, 79-85	8	66

83	The calcitonin gene-related peptide antagonist CGRP8-37 increases the latency to withdrawal responses bilaterally in rats with unilateral experimental mononeuropathy, an effect reversed by naloxone. <i>Neuroscience</i> , 1996 , 71, 523-31	3.9	64
82	Intrathecal CGRP8-37-induced bilateral increase in hindpaw withdrawal latency in rats with unilateral inflammation. <i>British Journal of Pharmacology</i> , 1996 , 117, 43-50	8.6	62
81	The calcitonin gene-related peptide antagonist CGRP8-37 increases the latency to withdrawal responses in rats. <i>Brain Research</i> , 1994 , 653, 223-30	3.7	62
80	Transcutaneous electrical nerve stimulation (TENS) as compared to placebo TENS for the relief of acute oro-facial pain. <i>Pain</i> , 1983 , 15, 157-65	8	62
79	The influence of pain intensity on somatosensory perception in patients suffering from subacute/chronic lateral epicondylalgia. <i>European Journal of Pain</i> , 2000 , 4, 57-71	3.7	60
78	Quantitative sensory testing using DFNS protocol in Europe: an evaluation of heterogeneity across multiple centers in patients with peripheral neuropathic pain and healthy subjects. <i>Pain</i> , 2016 , 157, 750-758	8.8	59
77	Inflammatory and Neuropathic Pain From Bench to Bedside: What Went Wrong?. <i>Journal of Pain</i> , 2018 , 19, 571-588	5.2	55
76	Injection of hypertonic saline into musculus infraspinatus resulted in referred pain and sensory disturbances in the ipsilateral upper arm. <i>European Journal of Pain</i> , 2000 , 4, 73-82	3.7	54
75	Antiretroviral therapy may improve sensory function in HIV-infected patients: a pilot study. <i>Neurology</i> , 2000 , 54, 2120-7	6.5	50
74	Painful traumatic peripheral partial nerve injury-sensory dysfunction profiles comparing outcomes of bedside examination and quantitative sensory testing. <i>European Journal of Pain</i> , 2008 , 12, 397-402	3.7	49
73	Somatosensory perception in patients suffering from long-term trapezius myalgia at the site overlying the most painful part of the muscle and in an area of pain referral. <i>European Journal of Pain</i> , 2003 , 7, 267-76	3.7	48
72	Painful and non-painful neuropathy in HIV-infected patients: an analysis of somatosensory nerve function. <i>European Journal of Pain</i> , 2003 , 7, 23-31	3.7	47
71	SUNCT may be another manifestation of orbital venous vasculitis. <i>Headache</i> , 1992 , 32, 384-9	4.2	47
70	Who is healthy? Aspects to consider when including healthy volunteers in QST-based studies-a consensus statement by the EUROPAIN and NEUROPAIN consortia. <i>Pain</i> , 2015 , 156, 2203-2211	8	42
69	Dynamic mechanical allodynia: on the relationship between temporo-spatial stimulus parameters and evoked pain in patients with peripheral neuropathy. <i>Pain</i> , 2005 , 115, 264-272	8	42
68	Neuropathic pain needs systematic classification. <i>European Journal of Pain</i> , 2013 , 17, 953-6	3.7	41
67	Cerebrospinal fluid mononuclear cell counts influence CSF HIV-1 RNA levels. <i>Journal of Acquired Immune Deficiency Syndromes</i> , 1998 , 17, 214-9		41
66	Time dependent differences in pain sensitivity during unilateral ischemic pain provocation in healthy volunteers. <i>European Journal of Pain</i> , 2006 , 10, 225-32	3.7	40

65	Increased postoperative pain and consumption of analgesics following acupuncture. <i>Pain</i> , 1991 , 44, 241-247	3.6	36
64	Mechanisms of dynamic mechanical allodynia and dysesthesia in patients with peripheral and central neuropathic pain. <i>European Journal of Pain</i> , 2011 , 15, 498-503	3.7	32
63	Effects of conditioning vibratory stimulation on pain threshold of the human tooth. <i>Acta Physiologica Scandinavica</i> , 1982 , 114, 601-4		31
62	Breast sensibility after bilateral risk-reducing mastectomy and immediate breast reconstruction: a prospective study. <i>Journal of Plastic, Reconstructive and Aesthetic Surgery</i> , 2013 , 66, 1521-7	1.7	30
61	A regression analysis study of the primary somatosensory cortex during pain. <i>NeuroImage</i> , 2002 , 16, 1142-50	7.5	30
60	Systemic adenosine infusion alleviates spontaneous and stimulus evoked pain in patients with peripheral neuropathic pain. <i>Anesthesia and Analgesia</i> , 1995 , 81, 713-7	3.9	30
59	Influence of heterotopic noxious conditioning stimulation on spontaneous pain and dynamic mechanical allodynia in central post-stroke pain patients. <i>Pain</i> , 2009 , 143, 84-91	8	29
58	The influence of experimental pain intensity in the local and referred pain area on somatosensory perception in the area of referred pain. <i>European Journal of Pain</i> , 2002 , 6, 413-25	3.7	29
57	Translational aspects of central sensitization induced by primary afferent activity: what it is and what it is not. <i>Pain</i> , 2014 , 155, 1932-1934	8	28
56	Heterotopic noxious conditioning stimulation (HNCS) reduced the intensity of spontaneous pain, but not of allodynia in painful peripheral neuropathy. <i>European Journal of Pain</i> , 2007 , 11, 452-62	3.7	28
55	Pain in ambulatory HIV-infected patients with and without intravenous drug use. <i>European Journal of Pain</i> , 1999 , 3, 157-164	3.7	28
54	Extrasegmental transcutaneous electrical nerve stimulation and mechanical vibratory stimulation as compared to placebo for the relief of acute oro-facial pain. <i>Pain</i> , 1985 , 23, 223-229	8	28
53	Effects of dextromethorphan in clinical doses on capsaicin-induced ongoing pain and mechanical hypersensitivity. <i>Journal of Pain and Symptom Management</i> , 1997 , 14, 195-201	4.8	26
52	Diagnostic work-up of neuropathic pain: computing, using questionnaires or examining the patient?. <i>European Journal of Pain</i> , 2007 , 11, 367-9	3.7	25
51	Symptom profiles in the painDETECT Questionnaire in patients with peripheral neuropathic pain stratified according to sensory loss in quantitative sensory testing. <i>Pain</i> , 2016 , 157, 1810-1818	8	25
50	Somatosensory function in patients with and without pain after traumatic peripheral nerve injury. <i>European Journal of Pain</i> , 2010 , 14, 847-53	3.7	24
49	Effects of calcitonin gene-related peptide-(8-37) on withdrawal responses in rats with inflammation. <i>European Journal of Pharmacology</i> , 1998 , 347, 275-82	5.3	24
48	Afferent stimulation induced pain relief in acute oro-facial pain and its failure to induce sufficient pain reduction in dental and oral surgery. <i>Pain</i> , 1984 , 20, 273-278	8	23

47	Influence of naloxone on relief of acute oro-facial pain by transcutaneous electrical nerve stimulation (TENS) or vibration. <i>Pain</i> , 1986 , 24, 323-329	8	23
46	Intrathecal CGRP(8-37) results in a bilateral increase in hindpaw withdrawal latency in rats with a unilateral thermal injury. <i>Neuropeptides</i> , 1997 , 31, 601-7	3.3	21
45	Pain development and consumption of analgesics after oral surgery in relation to personality characteristics. <i>Pain</i> , 1989 , 37, 271-277	8	21
44	Peripherally Administrated Morphine Attenuates Capsaicin-Induced Mechanical Hypersensitivity in Humans. <i>Anesthesia and Analgesia</i> , 1997 , 84, 595-599	3.9	19
43	A thin-section and freeze-fracture study of the pulp blood vessels in feline and human teeth. <i>Archives of Oral Biology</i> , 1984 , 29, 413-24	2.8	19
42	Opioid antagonists naloxone, beta-funaltrexamine and naltrindole, but not nor-binaltorphimine, reverse the increased hindpaw withdrawal latency in rats induced by intrathecal administration of the calcitonin gene-related peptide antagonist CGRP8-37. <i>Brain Research</i> , 1995 , 698, 23-9	3.7	18
41	Sensitivity after bilateral prophylactic mastectomy and immediate reconstruction. <i>Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery</i> , 2007 , 41, 178-83		17
40	Somatosensory status after pedicled or free TRAM flap surgery: a retrospective study. <i>Plastic and Reconstructive Surgery</i> , 1999 , 104, 1642-8	2.7	17
39	Ondansetron, a 5HT3-antagonist, does not alter dynamic mechanical allodynia or spontaneous ongoing pain in peripheral neuropathy. <i>Clinical Journal of Pain</i> , 2011 , 27, 323-9	3.5	16
38	The influence of brushing force and stroking velocity on dynamic mechanical allodynia in patients with peripheral neuropathy. <i>European Journal of Pain</i> , 2011 , 15, 389-94	3.7	16
37	On the repeatability of brush-evoked allodynia using a novel semi-quantitative method in patients with peripheral neuropathic pain. <i>Pain</i> , 2007 , 130, 40-6	8	16
36	Ultrastructural and electrophysiological changes in the olfactory epithelium following exposure to organic solvents. <i>Acta Oto-Laryngologica</i> , 1984 , 98, 351-61	1.6	16
35	The influence of intensity and duration of a painful conditioning stimulation on conditioned pain modulation in volunteers. <i>European Journal of Pain</i> , 2014 , 18, 853-61	3.7	15
34	No pain, still gain (of function): the relation between sensory profiles and the presence or absence of self-reported pain in a large multicenter cohort of patients with neuropathy. <i>Pain</i> , 2021 , 162, 718-727 ⁸		14
33	Long-term sensibility following nonautologous, immediate breast reconstruction. <i>Breast Journal</i> , 2007 , 13, 346-51	1.2	13
32	A tonic heat test stimulus yields a larger and more reliable conditioned pain modulation effect compared to a phasic heat test stimulus. <i>Pain Reports</i> , 2017 , 2, e626	3.5	12
31	Peripherally administrated morphine attenuates capsaicin-induced mechanical hypersensitivity in humans. <i>Anesthesia and Analgesia</i> , 1997 , 84, 595-9	3.9	12
30	Perceptual integration of intramuscular electrical stimulation in the focal and the referred pain area in healthy humans. <i>Pain</i> , 2003 , 105, 125-31	8	12

29	Threshold of tactile perception after nipple-sharing: a prospective study. <i>Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery</i> , 2002 , 36, 216-20		11
28	Changes of neuropeptide concentrations in the brain following experimentally induced mononeuropathy in Wistar Kyoto and spontaneously hypertensive rats. <i>Neuroscience Letters</i> , 1995 , 192, 93-6	3.3	10
27	The olfactory and respiratory epithelium in rhesus and squirrel monkeys studied with freeze-fracture technique. <i>Acta Oto-Laryngologica</i> , 1989 , 108, 259-67	1.6	10
26	L-tryptophan supplementation does not affect postoperative pain intensity or consumption of analgesics. <i>Pain</i> , 1991 , 44, 249-254	8	10
25	Chapter 34 Classification of neuropathic pain syndromes based on symptoms and signs. <i>Handbook of Clinical Neurology / Edited By P J Vinken and G W Bruyn</i> , 2006 , 81, 517-26	3	8
24	When is "pain" appropriate?. <i>Pain</i> , 1993 , 55, 403	8	5
23	Is acupuncture sufficient as the sole analgesic in oral surgery?. <i>Oral Surgery, Oral Medicine, and Oral Pathology</i> , 1987 , 64, 283-6		5
22	Acute neuropathic pain: equivalent or different to chronic neuropathic pain? A call for gathering of scientifically based information on acute neuropathic pain. <i>Pain</i> , 2019 , 160, 2413-2414	8	5
21	Patient phenotyping in clinical trials of chronic pain treatments: IMMPACT recommendations. <i>Pain Reports</i> , 2021 , 6, e899	3.5	5
20	Pain thresholds and intensities of CRPS type I and neuropathic pain in respect to sex. <i>European Journal of Pain</i> , 2020 , 24, 1058-1071	3.7	4
19	Refractory Chronic Pain Screening Tool (RCPST): a feasibility study to assess practicality and validity of identifying potential neurostimulation candidates. <i>Pain Medicine</i> , 2014 , 15, 281-91	2.8	4
18	Sensation Following Immediate Breast Reconstruction with Implants. <i>Breast Journal</i> , 2010 , 16, 633-8	1.2	4
17	The Graphical Index of Pain: a new web-based method for high-throughput screening of pain. <i>Pain</i> , 2020 , 161, 2255-2262	8	4
16	Psychophysical or spinal reflex measures when assessing conditioned pain modulation?. <i>European Journal of Pain</i> , 2019 , 23, 1879-1889	3.7	3
15	The perception threshold counterpart to dynamic and static mechanical allodynia assessed using von Frey filaments in peripheral neuropathic pain patients. <i>Scandinavian Journal of Pain</i> , 2011 , 2, 9-16	1.9	3
14	Dynamic mechanical allodynia in the secondary hyperalgesic area in the capsaicin model-Perceptually similar phenomena as in painful neuropathy?. <i>Scandinavian Journal of Pain</i> , 2011 , 2, 85-92	1.9	3
13	Yet another questionnaire is born!. <i>Pain</i> , 2010 , 150, 219	8	3
12	Opioids modulate the calcitonin gene-related peptide8-37-mediated hindpaw withdrawal latency increase in thermally injured rats. <i>Neuropeptides</i> , 1998 , 32, 173-7	3.3	2

11	Does a regional nerve block change cutaneous perception thresholds outside the anaesthetic area? Implications for the interpretation of diagnostic blocks. <i>Pain</i> , 1992 , 50, 163-167	8	2
10	Contralateral Sensory and Pain Perception Changes in Patients With Unilateral Neuropathy. <i>Neurology</i> , 2021 , 97, e389-e402	6.5	2
9	Comparing objective cognitive impairments in patients with peripheral neuropathic pain or fibromyalgia. <i>Scientific Reports</i> , 2021 , 11, 673	4.9	2
8	Letter to the Editor of Pain on Freynhagen et al.: Pseudoradicular and radicular low-back pain - a disease continuum rather than different entities? Answers from quantitative sensory testing. <i>Pain</i> 2007;135:65-74. <i>Pain</i> , 2008 , 135, 312-313	8	1
7	A tribute to Professor Ulf Lindblom, MD, PhD. <i>European Journal of Pain</i> , 2003 , 7, 299-299	3.7	1
6	Review of techniques useful for the assessment of sensory small fiber neuropathies: Report from an IFCN expert group.. <i>Clinical Neurophysiology</i> , 2022 , 136, 13-38	4.3	1
5	A narrative review on the analgesic effect of localised vibration - part 1: the neurophysiological basis.. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2022 ,	4.4	1
4	Commentary to "Challenges and opportunities in translational pain research-An opinion paper of the working group on translational pain research of the European Pain Federation (EFIC)" by Mouraux et al. <i>European Journal of Pain</i> , 2021 , 25, 1179-1180	3.7	0
3	Assessment of central sensitization in the clinic. Is it possible?. <i>Scandinavian Journal of Pain</i> , 2012 , 3, 175-176	3.7	0
2	Neuropathic pain From guidelines to clinical practice. <i>Scandinavian Journal of Pain</i> , 2012 , 3, 178-178	1.9	0
1	Comment on the commentary by H. Breivik. <i>European Journal of Pain</i> , 2010 , 14, 456	3.7	0