Per Hansson

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/2171729/per-hansson-publications-by-year.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

118 papers 12,448 47 111 g-index

126 14,497 5.3 6 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
118	Review of techniques useful for the assessment of sensory small fiber neuropathies: Report from an IFCN expert group <i>Clinical Neurophysiology</i> , 2022 , 136, 13-38	4.3	1
117	A narrative review on the analgesic effect of localised vibration - part 1: the neurophysiological basis <i>European Journal of Physical and Rehabilitation Medicine</i> , 2022 ,	4.4	1
116	No pain, still gain (of function): the relation between sensory profiles and the presence or absence of self-reported pain in a large multicenter cohort of patients with neuropathy. <i>Pain</i> , 2021 , 162, 718-72	27 ⁸	14
115	Commentary to "Challenges and opportunities in translational pain research-An opinion paper of the working group on translational pain research of the European Pain Federation (EFIC)" by Mouraux et al. <i>European Journal of Pain</i> , 2021 , 25, 1179-1180	3.7	Ο
114	Contralateral Sensory and Pain Perception Changes in Patients With Unilateral Neuropathy. <i>Neurology</i> , 2021 , 97, e389-e402	6.5	2
113	Comparing objective cognitive impairments in patients with peripheral neuropathic pain or fibromyalgia. <i>Scientific Reports</i> , 2021 , 11, 673	4.9	2
112	Patient phenotyping in clinical trials of chronic pain treatments: IMMPACT recommendations. <i>Pain Reports</i> , 2021 , 6, e899	3.5	5
111	Pain thresholds and intensities of CRPS type I and neuropathic pain in respect to sex. <i>European Journal of Pain</i> , 2020 , 24, 1058-1071	3.7	4
110	The Graphical Index of Pain: a new web-based method for high-throughput screening of pain. <i>Pain</i> , 2020 , 161, 2255-2262	8	4
109	The IASP classification of chronic pain for ICD-11: chronic neuropathic pain. <i>Pain</i> , 2019 , 160, 53-59	8	228
108	Psychophysical or spinal reflex measures when assessing conditioned pain modulation?. <i>European Journal of Pain</i> , 2019 , 23, 1879-1889	3.7	3
107	Acute neuropathic pain: equivalent or different to chronic neuropathic pain? A call for gathering of scientifically based information on acute neuropathic pain. <i>Pain</i> , 2019 , 160, 2413-2414	8	5
106	Inflammatory and Neuropathic Pain From Bench to Bedside: What Went Wrong?. <i>Journal of Pain</i> , 2018 , 19, 571-588	5.2	55
105	Peripheral neuropathic pain: a mechanism-related organizing principle based on sensory profiles. <i>Pain</i> , 2017 , 158, 261-272	8	310
104	Stratifying patients with peripheral neuropathic pain based on sensory profiles: algorithm and sample size recommendations. <i>Pain</i> , 2017 , 158, 1446-1455	8	94
103	A tonic heat test stimulus yields a larger and more reliable conditioned pain modulation effect compared to a phasic heat test stimulus. <i>Pain Reports</i> , 2017 , 2, e626	3.5	12
102	Neuropathic pain: an updated grading system for research and clinical practice. <i>Pain</i> , 2016 , 157, 1599-1	686	536

(2011-2016)

101	Symptom profiles in the painDETECT Questionnaire in patients with peripheral neuropathic pain stratified according to sensory loss in quantitative sensory testing. <i>Pain</i> , 2016 , 157, 1810-1818	8	25
100	Quantitative sensory testing using DFNS protocol in Europe: an evaluation of heterogeneity across multiple centers in patients with peripheral neuropathic pain and healthy subjects. <i>Pain</i> , 2016 , 157, 750)- <mark>8</mark> 58	59
99	Patient phenotyping in clinical trials of chronic pain treatments: IMMPACT recommendations. <i>Pain</i> , 2016 , 157, 1851-1871	8	178
98	Recommendations on practice of conditioned pain modulation (CPM) testing. <i>European Journal of Pain</i> , 2015 , 19, 805-6	3.7	239
97	Who is healthy? Aspects to consider when including healthy volunteers in QSTbased studies-a consensus statement by the EUROPAIN and NEUROPAIN consortia. <i>Pain</i> , 2015 , 156, 2203-2211	8	42
96	Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. <i>Lancet Neurology, The</i> , 2015 , 14, 162-73	24.1	1971
95	Refractory Chronic Pain Screening Tool (RCPST): a feasibility study to assess practicality and validity of identifying potential neurostimulation candidates. <i>Pain Medicine</i> , 2014 , 15, 281-91	2.8	4
94	The influence of intensity and duration of a painful conditioning stimulation on conditioned pain modulation in volunteers. <i>European Journal of Pain</i> , 2014 , 18, 853-61	3.7	15
93	Translational aspects of central sensitization induced by primary afferent activity: what it is and what it is not. <i>Pain</i> , 2014 , 155, 1932-1934	8	28
92	Value of quantitative sensory testing in neurological and pain disorders: NeuPSIG consensus. <i>Pain</i> , 2013 , 154, 1807-1819	8	350
91	Breast sensibility after bilateral risk-reducing mastectomy and immediate breast reconstruction: a prospective study. <i>Journal of Plastic, Reconstructive and Aesthetic Surgery</i> , 2013 , 66, 1521-7	1.7	30
90	Neuropathic pain needs systematic classification. European Journal of Pain, 2013, 17, 953-6	3.7	41
89	Assessment of central sensitization in the clinic. Is it possible?. Scandinavian Journal of Pain, 2012, 3, 17	511376	
88	Neuropathic pain E rom guidelines to clinical practice. <i>Scandinavian Journal of Pain</i> , 2012 , 3, 178-178	1.9	
87	The perception threshold counterpart to dynamic and static mechanical allodynia assessed using von Frey filaments in peripheral neuropathic pain patients. <i>Scandinavian Journal of Pain</i> , 2011 , 2, 9-16	1.9	3
86	Dynamic mechanical allodynia in the secondary hyperalgesic area in the capsaicin model-Perceptually similar phenomena as in painful neuropathy?. <i>Scandinavian Journal of Pain</i> , 2011 , 2, 85-92	1.9	3
85	Ondansetron, a 5HT3-antagonist, does not alter dynamic mechanical allodynia or spontaneous ongoing pain in peripheral neuropathy. <i>Clinical Journal of Pain</i> , 2011 , 27, 323-9	3.5	16
84	NeuPSIG guidelines on neuropathic pain assessment. <i>Pain</i> , 2011 , 152, 14-27	8	694

83	The influence of brushing force and stroking velocity on dynamic mechanical allodynia in patients with peripheral neuropathy. <i>European Journal of Pain</i> , 2011 , 15, 389-94	3.7	16
82	Mechanisms of dynamic mechanical allodynia and dysesthesia in patients with peripheral and central neuropathic pain. <i>European Journal of Pain</i> , 2011 , 15, 498-503	3.7	32
81	Yet another questionnaire is born!. <i>Pain</i> , 2010 , 150, 219	8	3
80	Somatosensory function in patients with and without pain after traumatic peripheral nerve injury. <i>European Journal of Pain</i> , 2010 , 14, 847-53	3.7	24
79	Recommendations on terminology and practice of psychophysical DNIC testing. <i>European Journal of Pain</i> , 2010 , 14, 339	3.7	326
78	Comment on the commentary by H. Breivik. <i>European Journal of Pain</i> , 2010 , 14, 456	3.7	
77	EFNS guidelines on the pharmacological treatment of neuropathic pain: 2010 revision. <i>European Journal of Neurology</i> , 2010 , 17, 1113-e88	6	1190
76	Sensation Following Immediate Breast Reconstruction with Implants. <i>Breast Journal</i> , 2010 , 16, 633-8	1.2	4
75	Influence of heterotopic noxious conditioning stimulation on spontaneous pain and dynamic mechanical allodynia in central post-stroke pain patients. <i>Pain</i> , 2009 , 143, 84-91	8	29
74	Assessment of neuropathic pain in primary care. American Journal of Medicine, 2009, 122, S13-21	2.4	150
73	Clinical and pre-clinical pain assessment: are we measuring the same thing?. Pain, 2008, 135, 7-10	8	150
72	Letter to the Editor of Pain on Freynhagen et al.: Pseudoradicular and radicular low-back pain - a disease continuum rather than different entities? Answers from quantitative sensory testing. Pain 2007;135:65-74. <i>Pain</i> , 2008 , 135, 312-313	8	1
71	Painful traumatic peripheral partial nerve injury-sensory dysfunction profiles comparing outcomes of bedside examination and quantitative sensory testing. <i>European Journal of Pain</i> , 2008 , 12, 397-402	3.7	49
70	Heterotopic noxious conditioning stimulation (HNCS) reduced the intensity of spontaneous pain, but not of allodynia in painful peripheral neuropathy. <i>European Journal of Pain</i> , 2007 , 11, 452-62	3.7	28
69	Diagnostic work-up of neuropathic pain: computing, using questionnaires or examining the patient?. <i>European Journal of Pain</i> , 2007 , 11, 367-9	3.7	25
68	Long-term sensibility following nonautologous, immediate breast reconstruction. <i>Breast Journal</i> , 2007 , 13, 346-51	1.2	13
67	EFNS guidelines on neurostimulation therapy for neuropathic pain. <i>European Journal of Neurology</i> , 2007 , 14, 952-70	6	476
66	Sensitivity after bilateral prophylactic mastectomy and immediate reconstruction. <i>Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery</i> , 2007 , 41, 178-83		17

(2002-2007)

65	On the repeatability of brush-evoked allodynia using a novel semi-quantitative method in patients with peripheral neuropathic pain. <i>Pain</i> , 2007 , 130, 40-6	8	16
64	Usefulness and limitations of quantitative sensory testing: clinical and research application in neuropathic pain states. <i>Pain</i> , 2007 , 129, 256-259	8	221
63	Chapter 34 Classification of neuropathic pain syndromes based on symptoms and signs. <i>Handbook of Clinical Neurology / Edited By P J Vinken and G W Bruyn</i> , 2006 , 81, 517-26	3	8
62	EFNS guidelines on pharmacological treatment of neuropathic pain. <i>European Journal of Neurology</i> , 2006 , 13, 1153-69	6	638
61	Time dependent differences in pain sensitivity during unilateral ischemic pain provocation in healthy volunteers. <i>European Journal of Pain</i> , 2006 , 10, 225-32	3.7	40
60	Pharmacological treatment of peripheral neuropathic pain conditions based on shared commonalities despite multiple etiologies. <i>Pain</i> , 2005 , 113, 251-254	8	80
59	Dynamic mechanical allodynia: on the relationship between temporo-spatial stimulus parameters and evoked pain in patients with peripheral neuropathy. <i>Pain</i> , 2005 , 115, 264-272	8	42
58	Context-dependent deactivation of the amygdala during pain. <i>Journal of Cognitive Neuroscience</i> , 2004 , 16, 1289-301	3.1	87
57	Brainstem involvement in the initial response to pain. <i>NeuroImage</i> , 2004 , 22, 995-1005	7.9	69
56	Painful and non-painful neuropathy in HIV-infected patients: an analysis of somatosensory nerve function. <i>European Journal of Pain</i> , 2003 , 7, 23-31	3.7	47
55	A tribute to Professor Ulf Lindblom, MD, PhD. European Journal of Pain, 2003, 7, 299-299	3.7	1
54	Difficulties in stratifying neuropathic pain by mechanisms. European Journal of Pain, 2003, 7, 353-7	3.7	91
53	Somatosensory perception in patients suffering from long-term trapezius myalgia at the site overlying the most painful part of the muscle and in an area of pain referral. <i>European Journal of Pain</i> , 2003 , 7, 267-76	3.7	48
52	Perceptual integration of intramuscular electrical stimulation in the focal and the referred pain area in healthy humans. <i>Pain</i> , 2003 , 105, 125-31	8	12
51	The influence of experimental pain intensity in the local and referred pain area on somatosensory perception in the area of referred pain. <i>European Journal of Pain</i> , 2002 , 6, 413-25	3.7	29
50	Somatosensory perception in a remote pain-free area and function of diffuse noxious inhibitory controls (DNIC) in patients suffering from long-term trapezius myalgia. <i>European Journal of Pain</i> , 2002 , 6, 149-59	3.7	83
49	Somatosensory perception and function of diffuse noxious inhibitory controls (DNIC) in patients suffering from rheumatoid arthritis. <i>European Journal of Pain</i> , 2002 , 6, 161-76	3.7	131
48	Neuropathic pain: clinical characteristics and diagnostic workup. <i>European Journal of Pain</i> , 2002 , 6 Suppl A, 47-50	3.7	106

47	Threshold of tactile perception after nipple-sharing: a prospective study. <i>Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery</i> , 2002 , 36, 216-20	11
46	A regression analysis study of the primary somatosensory cortex during pain. <i>NeuroImage</i> , 2002 , 16, 114 2 :5	50 30
45	The influence of pain intensity on somatosensory perception in patients suffering from subacute/chronic lateral epicondylalgia. <i>European Journal of Pain</i> , 2000 , 4, 57-71	60
44	Antiretroviral therapy may improve sensory function in HIV-infected patients: a pilot study. Neurology, 2000 , 54, 2120-7	50
43	Injection of hypertonic saline into musculus infraspinatus resulted in referred pain and sensory disturbances in the ipsilateral upper arm. <i>European Journal of Pain</i> , 2000 , 4, 73-82	54
42	Pain in ambulatory HIV-infected patients with and without intravenous drug use. <i>European Journal of Pain</i> , 1999 , 3, 157-164	28
41	Somatosensory status after pedicled or free TRAM flap surgery: a retrospective study. <i>Plastic and Reconstructive Surgery</i> , 1999 , 104, 1642-8	17
40	Pressure pain thresholds in different tissues in one body region. The influence of skin sensitivity in pressure algometry. <i>Journal of Rehabilitation Medicine</i> , 1999 , 31, 89-93	126
39	Opioids modulate the calcitonin gene-related peptide8-37-mediated hindpaw withdrawal latency increase in thermally injured rats. <i>Neuropeptides</i> , 1998 , 32, 173-7	2
38	Effects of calcitonin gene-related peptide-(8-37) on withdrawal responses in rats with inflammation. <i>European Journal of Pharmacology</i> , 1998 , 347, 275-82	24
37	Cerebrospinal fluid mononuclear cell counts influence CSF HIV-1 RNA levels. <i>Journal of Acquired Immune Deficiency Syndromes</i> , 1998 , 17, 214-9	41
36	Peripherally administrated morphine attenuates capsaicin-induced mechanical hypersensitivity in humans. <i>Anesthesia and Analgesia</i> , 1997 , 84, 595-9	12
35	Peripherally Administrated Morphine Attenuates Capsaicin-Induced Mechanical Hypersensitivity in Humans. <i>Anesthesia and Analgesia</i> , 1997 , 84, 595-599	19
34	Peripheral alpha-adrenoreceptors are involved in the development of capsaicin induced ongoing and stimulus evoked pain in humans. <i>Pain</i> , 1997 , 69, 79-85	66
33	Modulatory influence on somatosensory perception from vibration and heterotopic noxious conditioning stimulation (HNCS) in fibromyalgia patients and healthy subjects. <i>Pain</i> , 1997 , 70, 41-51	401
32	Effects of dextromethorphan in clinical doses on capsaicin-induced ongoing pain and mechanical hypersensitivity. <i>Journal of Pain and Symptom Management</i> , 1997 , 14, 195-201	3 26
31	Intrathecal CGRP(8-37) results in a bilateral increase in hindpaw withdrawal latency in rats with a unilateral thermal injury. <i>Neuropeptides</i> , 1997 , 31, 601-7	21
30	Intrathecal CGRP8-37-induced bilateral increase in hindpaw withdrawal latency in rats with unilateral inflammation. <i>British Journal of Pharmacology</i> , 1996 , 117, 43-50	6 62

(1991-1996)

29	Modulation of pressure pain thresholds during and following isometric contraction in patients with fibromyalgia and in healthy controls. <i>Pain</i> , 1996 , 64, 415-423	8	168
28	The calcitonin gene-related peptide antagonist CGRP8-37 increases the latency to withdrawal responses bilaterally in rats with unilateral experimental mononeuropathy, an effect reversed by naloxone. <i>Neuroscience</i> , 1996 , 71, 523-31	3.9	64
27	Sensory dysfunction in fibromyalgia patients with implications for pathogenic mechanisms. <i>Pain</i> , 1996 , 68, 375-83	8	256
26	Opioid antagonists naloxone, beta-funaltrexamine and naltrindole, but not nor-binaltorphimine, reverse the increased hindpaw withdrawal latency in rats induced by intrathecal administration of the calcitonin gene-related peptide antagonist CGRP8-37. <i>Brain Research</i> , 1995 , 698, 23-9	3.7	18
25	Systemic Adenosine Infusion Alleviates Spontaneous and Stimulus Evoked Pain in Patients with Peripheral Neuropathic Pain. <i>Anesthesia and Analgesia</i> , 1995 , 81, 713-717	3.9	91
24	Changes of neuropeptide concentrations in the brain following experimentally induced mononeuropathy in Wistar Kyoto and spontaneously hypertensive rats. <i>Neuroscience Letters</i> , 1995 , 192, 93-6	3.3	10
23	Systemic adenosine infusion: a new treatment modality to alleviate neuropathic pain. <i>Pain</i> , 1995 , 61, 155-158	8	76
22	Central representation of chronic ongoing neuropathic pain studied by positron emission tomography. <i>Pain</i> , 1995 , 63, 225-236	8	460
21	Increased pressure pain sensibility in fibromyalgia patients is located deep to the skin but not restricted to muscle tissue. <i>Pain</i> , 1995 , 63, 335-339	8	100
20	Systemic adenosine infusion alleviates spontaneous and stimulus evoked pain in patients with peripheral neuropathic pain. <i>Anesthesia and Analgesia</i> , 1995 , 81, 713-7	3.9	30
19	The calcitonin gene-related peptide antagonist CGRP8-37 increases the latency to withdrawal responses in rats. <i>Brain Research</i> , 1994 , 653, 223-30	3.7	62
18	When is "pain" appropriate?. <i>Pain</i> , 1993 , 55, 403	8	5
17	Does a regional nerve block change cutaneous perception thresholds outside the anaesthetic area? Implications for the interpretation of diagnostic blocks. <i>Pain</i> , 1992 , 50, 163-167	8	2
16	SUNCT may be another manifestation of orbital venous vasculitis. <i>Headache</i> , 1992 , 32, 384-9	4.2	47
15	Intraputaminal infusion of nerve growth factor to support adrenal medullary autografts in Parkinson@ disease. One-year follow-up of first clinical trial. <i>Archives of Neurology</i> , 1991 , 48, 373-81		270
14	L-tryptophan supplementation does not affect postoperative pain intensity or consumption of analgesics. <i>Pain</i> , 1991 , 44, 249-254	8	10
13	Increased postoperative pain and consumption of analgesics following acupuncture. <i>Pain</i> , 1991 , 44, 241	-847	36
12	Concentrations of neuropeptides substance P, neurokinin A, calcitonin gene-related peptide, neuropeptide Y and vasoactive intestinal polypeptide in synovial fluid of the human temporomandibular joint. A correlation with symptoms, signs and arthroscopic findings.	2.9	80

11	The olfactory and respiratory epithelium in rhesus and squirrel monkeys studied with freeze-fracture technique. <i>Acta Oto-Laryngologica</i> , 1989 , 108, 259-67	1.6	10	
10	Pain development and consumption of analgesics after oral surgery in relation to personality characteristics. <i>Pain</i> , 1989 , 37, 271-277	8	21	
9	Is acupuncture sufficient as the sole analgesic in oral surgery?. <i>Oral Surgery, Oral Medicine, and Oral Pathology</i> , 1987 , 64, 283-6		5	
8	Influence of naloxone on relief of acute oro-facial pain by transcutaneous electrical nerve stimulation (TENS) or vibration. <i>Pain</i> , 1986 , 24, 323-329	8	23	
7	Extrasegmental transcutaneous electrical nerve stimulation and mechanical vibratory stimulation as compared to placebo for the relief of acute oro-facial pain. <i>Pain</i> , 1985 , 23, 223-229	8	28	
6	Afferent stimulation induced pain relief in acute oro-facial pain and its failure to induce sufficient pain reduction in dental and oral surgery. <i>Pain</i> , 1984 , 20, 273-278	8	23	
5	A thin-section and freeze-fracture study of the pulp blood vessels in feline and human teeth. <i>Archives of Oral Biology</i> , 1984 , 29, 413-24	2.8	19	
4	Ultrastructural and electrophysiological changes in the olfactory epithelium following exposure to organic solvents. <i>Acta Oto-Laryngologica</i> , 1984 , 98, 351-61	1.6	16	
3	Transcutaneous electrical nerve stimulation (TENS) as compared to placebo TENS for the relief of acute oro-facial pain. <i>Pain</i> , 1983 , 15, 157-65	8	62	
2	Effects of conditioning vibratory stimulation on pain threshold of the human tooth. <i>Acta Physiologica Scandinavica</i> , 1982 , 114, 601-4		31	
1	Vibratory stimulation for the relief of pain of dental origin. <i>Pain</i> , 1981 , 10, 37-45	8	73	