

Rob A Cairns

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2174704/rob-a-cairns-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

26

papers

7,800

citations

20

h-index

27

g-index

27

ext. papers

8,736

ext. citations

13.1

avg, IF

6.15

L-index

#	Paper	IF	Citations
26	Loss of 5-hydroxymethylcytosine is a frequent event in peripheral T-cell lymphomas. <i>Haematologica</i> , 2018 , 103, e115-e118	6.6	19
25	p53 mutants cooperate with HIF-1 in transcriptional regulation of extracellular matrix components to promote tumor progression. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, E10869-E10878	11.5	73
24	Fire and water: Tumor cell adaptation to metabolic conditions. <i>Experimental Cell Research</i> , 2017 , 356, 204-208	4.2	11
23	Metabolic targeting of HIF-dependent glycolysis reduces lactate, increases oxygen consumption and enhances response to high-dose single-fraction radiotherapy in hypoxic solid tumors. <i>BMC Cancer</i> , 2017 , 17, 418	4.8	29
22	IDH1 deficiency attenuates gluconeogenesis in mouse liver by impairing amino acid utilization. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017 , 114, 292-297	11.5	13
21	S-2HG is an immunometabolite that shapes the T-cell response. <i>Cell Death and Differentiation</i> , 2017 , 24, 195-196	12.7	2
20	Mutant IDH1 Downregulates ATM and Alters DNA Repair and Sensitivity to DNA Damage Independent of TET2. <i>Cancer Cell</i> , 2016 , 30, 337-348	24.3	121
19	An Alternative Sugar Fuels AML. <i>Cancer Cell</i> , 2016 , 30, 660-662	24.3	4
18	PINK1 Is a Negative Regulator of Growth and the Warburg Effect in Glioblastoma. <i>Cancer Research</i> , 2016 , 76, 4708-19	10.1	80
17	Idh1 mutations contribute to the development of T-cell malignancies in genetically engineered mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016 , 113, 1387-92	11.5	13
16	The IDH2 R172K mutation associated with angioimmunoblastic T-cell lymphoma produces 2HG in T cells and impacts lymphoid development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016 , 113, 15084-15089	11.5	58
15	Lung Cancer Resets the Liver's Metabolic Clock. <i>Cell Metabolism</i> , 2016 , 23, 767-9	24.6	1
14	ALDH2(E487K) mutation increases protein turnover and promotes murine hepatocarcinogenesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 9088-93	11.5	54
13	Drivers of the Warburg phenotype. <i>Cancer Journal (Sudbury, Mass)</i> , 2015 , 21, 56-61	2.2	40
12	Mutant IDH is sufficient to initiate enchondromatosis in mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 2829-34	11.5	80
11	Oncogenic isocitrate dehydrogenase mutations: mechanisms, models, and clinical opportunities. <i>Cancer Discovery</i> , 2013 , 3, 730-41	24.4	314
10	IDH2 mutations are frequent in angioimmunoblastic T-cell lymphoma. <i>Blood</i> , 2012 , 119, 1901-3	2.2	364

9	Recurrent TET2 mutations in peripheral T-cell lymphomas correlate with TFH-like features and adverse clinical parameters. <i>Blood</i> , 2012 , 120, 1466-9	2.2	319
8	D-2-hydroxyglutarate produced by mutant IDH1 perturbs collagen maturation and basement membrane function. <i>Genes and Development</i> , 2012 , 26, 2038-49	12.6	218
7	Regulation of cancer cell metabolism. <i>Nature Reviews Cancer</i> , 2011 , 11, 85-95	31.3	3403
6	Cancer-associated metabolite 2-hydroxyglutarate accumulates in acute myelogenous leukemia with isocitrate dehydrogenase 1 and 2 mutations. <i>Journal of Experimental Medicine</i> , 2010 , 207, 339-44	16.6	564
5	Pharmacologically increased tumor hypoxia can be measured by 18F-Fluoroazomycin arabinoside positron emission tomography and enhances tumor response to hypoxic cytotoxin PR-104. <i>Clinical Cancer Research</i> , 2009 , 15, 7170-4	12.9	29
4	Metabolic targeting of hypoxia and HIF1 in solid tumors can enhance cytotoxic chemotherapy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 9445-50	11.5	134
3	HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. <i>Cell Metabolism</i> , 2006 , 3, 187-97	24.6	1595
2	A fluorescent orthotopic model of metastatic cervical carcinoma. <i>Clinical and Experimental Metastasis</i> , 2004 , 21, 275-81	4.7	30
1	Acute hypoxia enhances spontaneous lymph node metastasis in an orthotopic murine model of human cervical carcinoma. <i>Cancer Research</i> , 2004 , 64, 2054-61	10.1	186