

Thorsten Schlomm

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2182737/thorsten-schlomm-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

221
papers

10,140
citations

48
h-index

95
g-index

242
ext. papers

11,920
ext. citations

6.3
avg, IF

5.32
L-index

#	Paper	IF	Citations
221	Evolution of Targeted Prostate Biopsy by Adding Micro-Ultrasound to the Magnetic Resonance Imaging Pathway. <i>European Urology Focus</i> , 2021 , 7, 1292-1299	5.1	15
220	Chromosome 5 harbors two independent deletion hotspots at 5q13 and 5q21 that characterize biologically different subsets of aggressive prostate cancer. <i>International Journal of Cancer</i> , 2021 , 148, 748-758	7.5	2
219	Reduced anoctamin 7 (ANO7) expression is a strong and independent predictor of poor prognosis in prostate cancer. <i>Cancer Biology and Medicine</i> , 2021 , 18, 245-255	5.2	1
218	Overexpression of the TRIM24 E3 Ubiquitin Ligase is Linked to Genetic Instability and Predicts Unfavorable Prognosis in Prostate Cancer. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2021 , 29, e29-e38	1.9	0
217	Personalizing Localized Prostate Cancer: Validation of a Combined Clinical Cell-cycle Risk (CCR) Score Threshold for Prognosticating Benefit From Multimodality Therapy. <i>Clinical Genitourinary Cancer</i> , 2021 , 19, 296-304.e3	3.3	3
216	Upregulation of Phosphatase 1 Nuclear-Targeting Subunit (PNUTS) Is an Independent Predictor of Poor Prognosis in Prostate Cancer. <i>Disease Markers</i> , 2020 , 2020, 7050146	3.2	2
215	Upregulation of the heterogeneous nuclear ribonucleoprotein hnRNP1 is an independent predictor of early biochemical recurrence in TMPRSS2:ERG fusion-negative prostate cancers. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2020 , 477, 625-636	5.1	1
214	A pre-specified model based on four kallikrein markers in blood improves predictions of adverse pathology and biochemical recurrence after radical prostatectomy. <i>British Journal of Cancer</i> , 2020 , 123, 604-609	8.7	3
213	Robot-Assisted versus Laparoscopic Donor Nephrectomy: A Comparison of 250 Cases. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	2
212	Up regulation of the Hippo signalling effector YAP1 is linked to early biochemical recurrence in prostate cancers. <i>Scientific Reports</i> , 2020 , 10, 8916	4.9	5
211	The impact of long-term androgen deprivation therapy on cognitive function and socioeconomic decision making in prostate cancer patients. <i>Psycho-Oncology</i> , 2020 , 29, 1338-1346	3.9	7
210	Should We Perform Old-for-Old Kidney Transplantation during the COVID-19 Pandemic? The Risk for Post-Operative Intensive Stay. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	1
209	Upregulation of the transcription factor TFAP2D is associated with aggressive tumor phenotype in prostate cancer lacking the TMPRSS2:ERG fusion. <i>Molecular Medicine</i> , 2020 , 26, 24	6.2	2
208	Loss of the adhesion molecule CEACAM1 is associated with early biochemical recurrence in TMPRSS2:ERG fusion-positive prostate cancers. <i>International Journal of Cancer</i> , 2020 , 147, 575-583	7.5	2
207	Reduced KLK2 expression is a strong and independent predictor of poor prognosis in ERG-negative prostate cancer. <i>Prostate</i> , 2020 , 80, 1097-1107	4.2	5
206	Subcellular Compartmentalization of Survivin is Associated with Biological Aggressiveness and Prognosis in Prostate Cancer. <i>Scientific Reports</i> , 2020 , 10, 3250	4.9	3
205	High CHK2 protein expression is a strong and independent prognostic feature in ERG negative prostate cancer. <i>Pathology</i> , 2020 , 52, 421-430	1.6	2

204	Prostatakarzinom: Onkologische Kennzeichen. <i>Springer Reference Medizin</i> , 2020 , 1-18	0	
203	Repeat Radiation for Local Recurrence of Head and Neck Tumors and in Prostate Cancer. <i>Deutsches A&#x0308;rztblatt International</i> , 2020 , 117, 167-174	2.5	2
202	Therapie bei isoliertem PSA-Anstieg. <i>Springer Reference Medizin</i> , 2020 , 1-12	0	
201	Loss of cytoplasmic survivin expression is an independent predictor of poor prognosis in radically operated prostate cancer patients. <i>Cancer Medicine</i> , 2020 , 9, 1409-1418	4.8	4
200	High RSF1 protein expression is an independent prognostic feature in prostate cancer. <i>Acta Oncologica</i> , 2020 , 59, 268-273	3.2	4
199	Expression of CCCTC-binding factor (CTCF) is linked to poor prognosis in prostate cancer. <i>Molecular Oncology</i> , 2020 , 14, 129-138	7.9	8
198	Claudin-1 upregulation is associated with favorable tumor features and a reduced risk for biochemical recurrence in ERG-positive prostate cancer. <i>World Journal of Urology</i> , 2020 , 38, 2185-2196	4	6
197	Secreted Frizzled-Related Protein 4 (SFRP4) Is an Independent Prognostic Marker in Prostate Cancers Lacking TMPRSS2: ERG Fusions. <i>Pathology and Oncology Research</i> , 2020 , 26, 2709-2722	2.6	4
196	Increased Cytoplasmic CD138 Expression Is Associated with Aggressive Characteristics in Prostate Cancer and Is an Independent Predictor for Biochemical Recurrence. <i>BioMed Research International</i> , 2020 , 2020, 5845374	3	3
195	High B7-H3 expression is linked to increased risk of prostate cancer progression. <i>Pathology International</i> , 2020 , 70, 733-742	1.8	8
194	Xenograft-derived mRNA/miR and protein interaction networks of systemic dissemination in human prostate cancer. <i>European Journal of Cancer</i> , 2020 , 137, 93-107	7.5	4
193	EGFR as a stable marker of prostate cancer dissemination to bones. <i>British Journal of Cancer</i> , 2020 , 123, 1767-1774	8.7	4
192	Epithelial splicing regulatory protein 1 and 2 (ESRP1 and ESRP2) upregulation predicts poor prognosis in prostate cancer. <i>BMC Cancer</i> , 2020 , 20, 1220	4.8	3
191	Tumor-Associated Release of Prostatic Cells into the Blood after Transrectal Ultrasound-Guided Biopsy in Patients with Histologically Confirmed Prostate Cancer. <i>Clinical Chemistry</i> , 2020 , 66, 161-168	5.5	13
190	The gordian knots. <i>World Journal of Urology</i> , 2020 , 38, 1815-1817	4	
189	Analysis of the prognostic utility of the cell cycle progression (CCP) score generated from needle biopsy in men treated with definitive therapy. <i>Prostate Cancer and Prostatic Diseases</i> , 2020 , 23, 102-107	6.2	16
188	Solid organ transplantation programs facing lack of empiric evidence in the COVID-19 pandemic: A By-proxy Society Recommendation Consensus approach. <i>American Journal of Transplantation</i> , 2020 , 20, 1826-1836	8.7	62
187	Upregulation of PTTG1 is associated with poor prognosis in prostate cancer. <i>Pathology International</i> , 2020 , 70, 441-451	1.8	4

186	Random forest-based modelling to detect biomarkers for prostate cancer progression. <i>Clinical Epigenetics</i> , 2019 , 11, 148	7.7	32
185	Loss of PSP94 expression is associated with early PSA recurrence and deteriorates outcome of deleted prostate cancers. <i>Cancer Biology and Medicine</i> , 2019 , 16, 319-330	5.2	1
184	SNW1 is a prognostic biomarker in prostate cancer. <i>Diagnostic Pathology</i> , 2019 , 14, 33	3	2
183	Prostate cancer prognosis in men with other malignancies prior to radical prostatectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2019 , 37, 575.e1-575.e7	2.8	2
182	Response to olaparib in a germline mutated prostate cancer and genetic events associated with resistance. <i>Journal of Physical Education and Sports Management</i> , 2019 , 5,	2.8	21
181	Aberrant expression of the microtubule-associated protein tau is an independent prognostic feature in prostate cancer. <i>BMC Cancer</i> , 2019 , 19, 193	4.8	13
180	Active Surveillance und die moderne Diagnostik des Prostatakarzinoms. <i>Onkologe</i> , 2019 , 25, 304-314	0.1	
179	Radikale Prostatektomie beim lokal fortgeschrittenen und metastasierten Prostatakarzinom. <i>Onkologe</i> , 2019 , 25, 315-321	0.1	
178	Aktive Berwachung des Prostatakarzinoms. <i>Uro-News</i> , 2019 , 23, 46-50	0	
177	Down-Regulation of S100A8 is an Independent Predictor of PSA Recurrence in Prostate Cancer Treated by Radical Prostatectomy. <i>Neoplasia</i> , 2019 , 21, 872-881	6.4	2
176	Analysis of the Effects of Day-Time vs. Night-Time Surgery on Renal Transplant Patient Outcomes. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	7
175	Native Nephrectomy before and after Renal Transplantation in Patients with Autosomal Dominant Polycystic Kidney Disease (ADPKD). <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	4
174	High-level expression of protein tyrosine phosphatase non-receptor 12 is a strong and independent predictor of poor prognosis in prostate cancer. <i>BMC Cancer</i> , 2019 , 19, 944	4.8	0
173	Up regulation of Rho-associated coiled-coil containing kinase1 (ROCK1) is associated with genetic instability and poor prognosis in prostate cancer. <i>Aging</i> , 2019 , 11, 7859-7879	5.6	7
172	A nuclear shift of GSK3 β protein is an independent prognostic factor in prostate cancer. <i>Oncotarget</i> , 2019 , 10, 1729-1744	3.3	2
171	Nuclear ELAC2 overexpression is associated with increased hazard for relapse after radical prostatectomy. <i>Oncotarget</i> , 2019 , 10, 4973-4986	3.3	3
170	Prognostic and diagnostic role of PSA immunohistochemistry: A tissue microarray study on 21,000 normal and cancerous tissues. <i>Oncotarget</i> , 2019 , 10, 5439-5453	3.3	12
169	Nuclear up regulation of the BRCA1-associated ubiquitinase BAP1 is associated with tumor aggressiveness in prostate cancers lacking the TMPRSS2:ERG fusion. <i>Oncotarget</i> , 2019 , 10, 7096-7111	3.3	3

168	Upregulation of SPDEF is associated with poor prognosis in prostate cancer. <i>Oncology Letters</i> , 2019 , 18, 5107-5118	2.6	3
167	A Novel Gene Signature-Based Model Predicts Biochemical Recurrence-Free Survival in Prostate Cancer Patients after Radical Prostatectomy. <i>Cancers</i> , 2019 , 12,	6.6	30
166	Aktive Überwachung des Prostatakarzinoms. <i>Im Focus Onkologie</i> , 2019 , 22, 54-58	0	
165	Loss of CCAAT-enhancer-binding protein alpha (CEBPA) is linked to poor prognosis in PTEN deleted and TMPRSS2:ERG fusion type prostate cancers. <i>Prostate</i> , 2019 , 79, 302-311	4.2	2
164	A functional ex vivo assay to detect PARP1-EJ repair and radiosensitization by PARP-inhibitor in prostate cancer. <i>International Journal of Cancer</i> , 2019 , 144, 1685-1696	7.5	11
163	5q21 deletion is often heterogeneous in prostate cancer. <i>Genes Chromosomes and Cancer</i> , 2019 , 58, 509-515	3	3
162	Contemporary approach to predict early biochemical recurrence after radical prostatectomy: update of the Walz nomogram. <i>Prostate Cancer and Prostatic Diseases</i> , 2018 , 21, 386-393	6.2	8
161	Long-term cancer control outcomes in patients with biochemical recurrence and the impact of time from radical prostatectomy to biochemical recurrence. <i>Prostate</i> , 2018 , 78, 676-681	4.2	13
160	High BCAR1 expression is associated with early PSA recurrence in ERG negative prostate cancer. <i>BMC Cancer</i> , 2018 , 18, 37	4.8	14
159	Immunohistochemically detected IDH1 mutation is rare and mostly heterogeneous in prostate cancer. <i>World Journal of Urology</i> , 2018 , 36, 877-882	4	20
158	Integrating Tertiary Gleason 5 Patterns into Quantitative Gleason Grading in Prostate Biopsies and Prostatectomy Specimens. <i>European Urology</i> , 2018 , 73, 674-683	10.2	32
157	Incidence, Risk Factors, Management, and Complications of Rectal Injuries During Radical Prostatectomy. <i>European Urology Focus</i> , 2018 , 4, 554-557	5.1	22
156	Validation of Cyclic Adenosine Monophosphate Phosphodiesterase-4D7 for its Independent Contribution to Risk Stratification in a Prostate Cancer Patient Cohort with Longitudinal Biological Outcomes. <i>European Urology Focus</i> , 2018 , 4, 376-384	5.1	5
155	13q deletion is linked to an adverse phenotype and poor prognosis in prostate cancer. <i>Genes Chromosomes and Cancer</i> , 2018 , 57, 504-512	5	22
154	The Prognostic PDE4D7 Score in a Diagnostic Biopsy Prostate Cancer Patient Cohort with Longitudinal Biological Outcomes. <i>Prostate Cancer</i> , 2018 , 2018, 5821616	1.9	5
153	PSCA expression is associated with favorable tumor features and reduced PSA recurrence in operated prostate cancer. <i>BMC Cancer</i> , 2018 , 18, 612	4.8	12
152	Marked Prognostic Impact of Minimal Lymphatic Tumor Spread in Prostate Cancer. <i>European Urology</i> , 2018 , 74, 376-386	10.2	40
151	Upregulation of centromere protein F is linked to aggressive prostate cancers. <i>Cancer Management and Research</i> , 2018 , 10, 5491-5504	3.6	11

150	Development and Characterization of a Spontaneously Metastatic Patient-Derived Xenograft Model of Human Prostate Cancer. <i>Scientific Reports</i> , 2018 , 8, 17535	4.9	14
149	Deletion of 3p13 is a late event linked to progression of : fusion prostate cancer. <i>Cancer Management and Research</i> , 2018 , 10, 5909-5917	3.6	2
148	Molecular Evolution of Early-Onset Prostate Cancer Identifies Molecular Risk Markers and Clinical Trajectories. <i>Cancer Cell</i> , 2018 , 34, 996-1011.e8	24.3	89
147	High concordance of TMPRSS-ERG fusion between primary prostate cancer and its lymph node metastases. <i>Oncology Letters</i> , 2018 , 16, 6238-6244	2.6	3
146	Up regulation of the steroid hormone synthesis regulator HSD3B2 is linked to early PSA recurrence in prostate cancer. <i>Experimental and Molecular Pathology</i> , 2018 , 105, 50-56	4.4	5
145	Oncologic and Functional Outcomes after Radical Prostatectomy for High or Very High Risk Prostate Cancer: European Validation of the Current NCCN Guideline. <i>Journal of Urology</i> , 2017 , 198, 354-361	2.5	27
144	Apurinic/aprimidinic endonuclease 1 (APE1/Ref-1) overexpression is an independent prognostic marker in prostate cancer without TMPRSS2:ERG fusion. <i>Molecular Carcinogenesis</i> , 2017 , 56, 2135-2145	5	12
143	Prognostic utility of biopsy-derived cell cycle progression score in patients with National Comprehensive Cancer Network low-risk prostate cancer undergoing radical prostatectomy: implications for treatment guidance. <i>BJU International</i> , 2017 , 120, 808-814	5.6	37
142	Androgen Receptor Deregulation Drives Bromodomain-Mediated Chromatin Alterations in Prostate Cancer. <i>Cell Reports</i> , 2017 , 19, 2045-2059	10.6	72
141	Overexpression of the A Disintegrin and Metalloproteinase ADAM15 is linked to a Small but Highly Aggressive Subset of Prostate Cancers. <i>Neoplasia</i> , 2017 , 19, 279-287	6.4	11
140	High concordance of findings obtained from transgluteal magnetic resonance imaging - and transrectal ultrasonography-guided biopsy as compared with prostatectomy specimens. <i>BJU International</i> , 2017 , 120, 365-376	5.6	2
139	High-Level Glyoxalase 1 (GLO1) expression is linked to poor prognosis in prostate cancer. <i>Prostate</i> , 2017 , 77, 1528-1538	4.2	6
138	FOXA1 expression is a strong independent predictor of early PSA recurrence in ERG negative prostate cancers treated by radical prostatectomy. <i>Carcinogenesis</i> , 2017 , 38, 1180-1187	4.6	10
137	Increased ERCC1 expression is linked to chromosomal aberrations and adverse tumor biology in prostate cancer. <i>BMC Cancer</i> , 2017 , 17, 504	4.8	5
136	Comparison of 11 Active Surveillance Protocols in Contemporary European Men Treated With Radical Prostatectomy. <i>Clinical Genitourinary Cancer</i> , 2017 ,	3.3	9
135	Up-regulation of Biglycan is Associated with Poor Prognosis and PTEN Deletion in Patients with Prostate Cancer. <i>Neoplasia</i> , 2017 , 19, 707-715	6.4	43
134	Mitochondrial mutations drive prostate cancer aggression. <i>Nature Communications</i> , 2017 , 8, 656	17.4	66
133	Does Cytoreductive Prostatectomy Really Have an Impact on Prognosis in Prostate Cancer Patients with Low-volume Bone Metastasis? Results from a Prospective Case-Control Study. <i>European Urology Focus</i> , 2017 , 3, 646-649	5.1	42

132	Up regulation and nuclear translocation of Y-box binding protein 1 (YB-1) is linked to poor prognosis in ERG-negative prostate cancer. <i>Scientific Reports</i> , 2017 , 7, 2056	4.9	23
131	Up-regulation of mismatch repair genes MSH6, PMS2 and MLH1 parallels development of genetic instability and is linked to tumor aggressiveness and early PSA recurrence in prostate cancer. <i>Carcinogenesis</i> , 2017 , 38, 19-27	4.6	29
130	Adjuvant radiation therapy is associated with better oncological outcome compared with salvage radiation therapy in patients with pN1 prostate cancer treated with radical prostatectomy. <i>BJU International</i> , 2017 , 119, 717-723	5.6	25
129	Functional Outcomes and Quality of Life After Radical Prostatectomy Only Versus a Combination of Prostatectomy with Radiation and Hormonal Therapy. <i>European Urology</i> , 2017 , 71, 330-336	10.2	42
128	High-Level βGlutamyl-Hydrolase (GGH) Expression is Linked to Poor Prognosis in ERG Negative Prostate Cancer. <i>International Journal of Molecular Sciences</i> , 2017 , 18,	6.3	15
127	Deletion of 8p is an independent prognostic parameter in prostate cancer. <i>Oncotarget</i> , 2017 , 8, 379-392	3.3	30
126	Family with sequence similarity 13C (FAM13C) overexpression is an independent prognostic marker in prostate cancer. <i>Oncotarget</i> , 2017 , 8, 31494-31508	3.3	3
125	PTEN loss detection in prostate cancer: comparison of PTEN immunohistochemistry and PTEN FISH in a large retrospective prostatectomy cohort. <i>Oncotarget</i> , 2017 , 8, 65566-65576	3.3	35
124	Deletion lengthening at chromosomes 6q and 16q targets multiple tumor suppressor genes and is associated with an increasingly poor prognosis in prostate cancer. <i>Oncotarget</i> , 2017 , 8, 108923-108935	3.3	21
123	Nerve-sparing Surgery Technique, Not the Preservation of the Neurovascular Bundles, Leads to Improved Long-term Continence Rates After Radical Prostatectomy. <i>European Urology</i> , 2016 , 69, 584-589	10.2	86
122	Salvage radical prostatectomy for recurrent prostate cancer: verification of European Association of Urology guideline criteria. <i>BJU International</i> , 2016 , 117, 55-61	5.6	32
121	p16 upregulation is linked to poor prognosis in ERG negative prostate cancer. <i>Tumor Biology</i> , 2016 , 37, 12655-12663	2.9	16
120	Heterogeneity of ERG expression in prostate cancer: a large section mapping study of entire prostatectomy specimens from 125 patients. <i>BMC Cancer</i> , 2016 , 16, 641	4.8	19
119	Quantitative comparison of DNA methylation assays for biomarker development and clinical applications. <i>Nature Biotechnology</i> , 2016 , 34, 726-37	44.5	204
118	Reduced AZGP1 expression is an independent predictor of early PSA recurrence and associated with ERG-fusion positive and PTEN deleted prostate cancers. <i>International Journal of Cancer</i> , 2016 , 138, 1199-206	7.5	21
117	Limited prognostic value of preoperative circulating tumor cells for early biochemical recurrence in patients with localized prostate cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2016 , 34, 235.e11-6	2.8	48
116	Phosphodiesterase Type 5 Inhibitor Use and Disease Recurrence After Prostate Cancer Treatment. <i>European Urology</i> , 2016 , 70, 824-828	10.2	14
115	Clinical Utility of Quantitative Gleason Grading in Prostate Biopsies and Prostatectomy Specimens. <i>European Urology</i> , 2016 , 69, 592-598	10.2	167

114	Aquaporin 5 expression is frequent in prostate cancer and shows a dichotomous correlation with tumor phenotype and PSA recurrence. <i>Human Pathology</i> , 2016 , 48, 102-10	3.7	13
113	Deletion of 18q is a strong and independent prognostic feature in prostate cancer. <i>Oncotarget</i> , 2016 , 7, 86339-86349	3.3	23
112	Heterogeneity and chronology of 6q15 deletion and ERG-fusion in prostate cancer. <i>Oncotarget</i> , 2016 , 7, 3897-904	3.3	6
111	Toxicity in a multimodality approach of radical prostatectomy with radiation and hormonal therapy.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 107-107	2.2	
110	Cytoplasmic accumulation of ELAVL1 is an independent predictor of biochemical recurrence associated with genomic instability in prostate cancer. <i>Prostate</i> , 2016 , 76, 259-72	4.2	18
109	Improved detection of circulating tumor cells in non-metastatic high-risk prostate cancer patients. <i>Scientific Reports</i> , 2016 , 6, 39736	4.9	73
108	A Clinician@ Guide to Avoiding and Managing Common Complications During and After Robot-assisted Laparoscopic Radical Prostatectomy. <i>European Urology Focus</i> , 2016 , 2, 30-48	5.1	8
107	The Combination of DNA Ploidy Status and PTEN/6q15 Deletions Provides Strong and Independent Prognostic Information in Prostate Cancer. <i>Clinical Cancer Research</i> , 2016 , 22, 2802-11	12.9	13
106	Re: The Molecular Taxonomy of Primary Prostate Cancer. <i>European Urology</i> , 2016 , 69, 1157	10.2	4
105	β-tubulin overexpression is linked to aggressive tumor features and shortened survival in clear cell renal cell carcinoma. <i>World Journal of Urology</i> , 2015 , 33, 1561-9	4	10
104	Prostate cancer-associated autoantibodies in serum against tumor-associated antigens as potential new biomarkers. <i>Journal of Proteomics</i> , 2015 , 119, 218-29	3.9	24
103	VEGFR-1 overexpression identifies a small subgroup of aggressive prostate cancers in patients treated by prostatectomy. <i>International Journal of Molecular Sciences</i> , 2015 , 16, 8591-606	6.3	3
102	Genomic Predictors of Outcome in Prostate Cancer. <i>European Urology</i> , 2015 , 68, 1033-44	10.2	136
101	HDAC1 overexpression independently predicts biochemical recurrence and is associated with rapid tumor cell proliferation and genomic instability in prostate cancer. <i>Experimental and Molecular Pathology</i> , 2015 , 98, 419-26	4.4	20
100	Tumor volume in insignificant prostate cancer: increasing threshold gains increasing risk. <i>Prostate</i> , 2015 , 75, 45-9	4.2	17
99	Saccharomyces cerevisiae-like 1 overexpression is frequent in prostate cancer and has markedly different effects in Ets-related gene fusion-positive and fusion-negative cancers. <i>Human Pathology</i> , 2015 , 46, 514-23	3.7	7
98	Overexpression of enhancer of zeste homolog 2 (EZH2) characterizes an aggressive subset of prostate cancers and predicts patient prognosis independently from pre- and postoperatively assessed clinicopathological parameters. <i>Carcinogenesis</i> , 2015 , 36, 1333-40	4.6	37
97	The Molecular Taxonomy of Primary Prostate Cancer. <i>Cell</i> , 2015 , 163, 1011-25	56.2	1713

96	The prognostic value of SUMO1/Sentrin specific peptidase 1 (SEN1) in prostate cancer is limited to ERG-fusion positive tumors lacking PTEN deletion. <i>BMC Cancer</i> , 2015 , 15, 538	4.8	27
95	Cytoplasmic Accumulation of Sequestosome 1 (p62) Is a Predictor of Biochemical Recurrence, Rapid Tumor Cell Proliferation, and Genomic Instability in Prostate Cancer. <i>Clinical Cancer Research</i> , 2015 , 21, 3471-9	12.9	36
94	BAZZA (TIP5) is involved in epigenetic alterations in prostate cancer and its overexpression predicts disease recurrence. <i>Nature Genetics</i> , 2015 , 47, 22-30	36.3	99
93	Safe-R: a novel score, accounting for oncological safe nerve-sparing at radical prostatectomy for localized prostate cancer. <i>World Journal of Urology</i> , 2015 , 33, 77-83	4	2
92	Oncological outcome after radical prostatectomy: Marital status does not make a difference. <i>International Journal of Urology</i> , 2015 , 22, 484-9	2.3	4
91	PSMA expression is highly homogenous in primary prostate cancer. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2015 , 23, 449-55	1.9	19
90	Concurrent deletion of 16q23 and PTEN is an independent prognostic feature in prostate cancer. <i>International Journal of Cancer</i> , 2015 , 137, 2354-63	7.5	33
89	Identification of pathologically favorable disease in intermediate-risk prostate cancer patients: Implications for active surveillance candidates selection. <i>Prostate</i> , 2015 , 75, 1484-91	4.2	19
88	Loss of SOX9 Expression Is Associated with PSA Recurrence in ERG-Positive and PTEN Deleted Prostate Cancers. <i>PLoS ONE</i> , 2015 , 10, e0128525	3.7	24
87	High-Level HOOK3 Expression Is an Independent Predictor of Poor Prognosis Associated with Genomic Instability in Prostate Cancer. <i>PLoS ONE</i> , 2015 , 10, e0134614	3.7	8
86	HOXB13 overexpression is an independent predictor of early PSA recurrence in prostate cancer treated by radical prostatectomy. <i>Oncotarget</i> , 2015 , 6, 12822-34	3.3	26
85	Heterogeneity in D'Amico classification-based low-risk prostate cancer: Differences in upgrading and upstaging according to active surveillance eligibility. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 329.e13-9	2.8	29
84	A visual-interactive system for prostate cancer cohort analysis. <i>IEEE Computer Graphics and Applications</i> , 2015 , 35, 44-55	1.7	21
83	Prevalence of chromosomal rearrangements involving non-ETS genes in prostate cancer. <i>International Journal of Oncology</i> , 2015 , 46, 1637-42	4.4	9
82	Expression of DNA ligase IV is linked to poor prognosis and characterizes a subset of prostate cancers harboring TMPRSS2:ERG fusion and PTEN deletion. <i>Oncology Reports</i> , 2015 , 34, 1211-20	3.5	6
81	Use of phosphodiesterase type 5 inhibitors may adversely impact biochemical recurrence after radical prostatectomy. <i>Journal of Urology</i> , 2015 , 193, 479-83	2.5	32
80	External validation of the CAPRA-S score to predict biochemical recurrence, metastasis and mortality after radical prostatectomy in a European cohort. <i>Journal of Urology</i> , 2015 , 193, 1970-5	2.5	43
79	Overexpression of thymidylate synthase (TYMS) is associated with aggressive tumor features and early PSA recurrence in prostate cancer. <i>Oncotarget</i> , 2015 , 6, 8377-87	3.3	33

78	Genomic deletion of chromosome 12p is an independent prognostic marker in prostate cancer. <i>Oncotarget</i> , 2015 , 6, 27966-79	3.3	28
77	βII-tubulin overexpression is an independent predictor of prostate cancer progression tightly linked to ERG fusion status and PTEN deletion. <i>American Journal of Pathology</i> , 2014 , 184, 609-17	5.8	44
76	NY-ESO-1 expression is tightly linked to TMPRSS2-ERG fusion in prostate cancer. <i>Prostate</i> , 2014 , 74, 1012-22	9	
75	High RNA-binding motif protein 3 expression is an independent prognostic marker in operated prostate cancer and tightly linked to ERG activation and PTEN deletions. <i>European Journal of Cancer</i> , 2014 , 50, 852-61	7.5	33
74	TMPRSS2-ERG fusions are strongly linked to young patient age in low-grade prostate cancer. <i>European Urology</i> , 2014 , 66, 978-81	10.2	48
73	Intratumor DNA methylation heterogeneity reflects clonal evolution in aggressive prostate cancer. <i>Cell Reports</i> , 2014 , 8, 798-806	10.6	177
72	Prognostic utility of the cell cycle progression score generated from biopsy in men treated with prostatectomy. <i>Journal of Urology</i> , 2014 , 192, 409-14	2.5	153
71	A feasible and time-efficient adaptation of NeuroSAFE for da Vinci robot-assisted radical prostatectomy. <i>European Urology</i> , 2014 , 66, 138-44	10.2	71
70	Patterns of TPD52 overexpression in multiple human solid tumor types analyzed by quantitative PCR. <i>International Journal of Oncology</i> , 2014 , 44, 609-15	4.4	22
69	Loss of somatostatin receptor subtype 2 in prostate cancer is linked to an aggressive cancer phenotype, high tumor cell proliferation and predicts early metastatic and biochemical relapse. <i>PLoS ONE</i> , 2014 , 9, e100469	3.7	15
68	CD57 expression in incidental, clinically manifest, and metastatic carcinoma of the prostate. <i>BioMed Research International</i> , 2014 , 2014, 356427	3	10
67	Comparison between complication rates of laser prostatectomy electrocautery transurethral resection of the prostate: A population-based study. <i>Canadian Urological Association Journal</i> , 2014 , 8, E419-24	1.2	4
66	High nuclear karyopherin β expression is a strong and independent predictor of biochemical recurrence in prostate cancer patients treated by radical prostatectomy. <i>Modern Pathology</i> , 2014 , 27, 96-106	9.8	24
65	The prognostic impact of high Nijmegen breakage syndrome (NBS1) gene expression in ERG-negative prostate cancers lacking PTEN deletion is driven by KPNA2 expression. <i>International Journal of Cancer</i> , 2014 , 135, 1399-407	7.5	29
64	Heterogeneity and chronology of PTEN deletion and ERG fusion in prostate cancer. <i>Modern Pathology</i> , 2014 , 27, 1612-20	9.8	59
63	Clinical significance of different types of p53 gene alteration in surgically treated prostate cancer. <i>International Journal of Cancer</i> , 2014 , 135, 1369-80	7.5	85
62	Aberrant presentation of HPA-reactive carbohydrates implies Selectin-independent metastasis formation in human prostate cancer. <i>Clinical Cancer Research</i> , 2014 , 20, 1791-802	12.9	23
61	Qualitative and Quantitative Requirements for Assessing Prognostic Markers in Prostate Cancer. <i>Microarrays (Basel, Switzerland)</i> , 2014 , 3, 137-58		2

60	Predicting radical prostatectomy outcome: Cell cycle progression (CCP) score compared with primary Gleason grade among men with clinical Gleason less than 7 who are upgraded to Gleason 7.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 13-13	2.2	2
59	Prognostic utility of the cell cycle progression (CCP) score generated from needle biopsy in men treated with prostatectomy.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 17-17	2.2	
58	MALDI imaging on tissue microarrays identifies molecular features associated with renal cell cancer phenotype. <i>Anticancer Research</i> , 2014 , 34, 2255-61	2.3	27
57	Recurrent deletion of 3p13 targets multiple tumour suppressor genes and defines a distinct subgroup of aggressive ERG fusion-positive prostate cancers. <i>Journal of Pathology</i> , 2013 , 231, 130-41	9.4	112
56	SPINK1 expression is tightly linked to 6q15- and 5q21-deleted ERG-fusion negative prostate cancers but unrelated to PSA recurrence. <i>Prostate</i> , 2013 , 73, 1690-8	4.2	34
55	Overexpression of the chromatin remodeler death-domain-associated protein in prostate cancer is an independent predictor of early prostate-specific antigen recurrence. <i>Human Pathology</i> , 2013 , 44, 1789-96	2.7	17
54	High mitochondria content is associated with prostate cancer disease progression. <i>Molecular Cancer</i> , 2013 , 12, 145	42.1	39
53	High lysophosphatidylcholine acyltransferase 1 expression independently predicts high risk for biochemical recurrence in prostate cancers. <i>Molecular Oncology</i> , 2013 , 7, 1001-11	7.9	40
52	High tissue density of FOXP3+ T cells is associated with clinical outcome in prostate cancer. <i>European Journal of Cancer</i> , 2013 , 49, 1273-9	7.5	83
51	Reduced CD147 expression is linked to ERG fusion-positive prostate cancers but lacks substantial impact on PSA recurrence in patients treated by radical prostatectomy. <i>Experimental and Molecular Pathology</i> , 2013 , 95, 227-34	4.4	13
50	Integrative genomic analyses reveal an androgen-driven somatic alteration landscape in early-onset prostate cancer. <i>Cancer Cell</i> , 2013 , 23, 159-70	24.3	259
49	Loss of pSer2448-mTOR expression is linked to adverse prognosis and tumor progression in ERG-fusion-positive cancers. <i>International Journal of Cancer</i> , 2013 , 132, 1333-40	7.5	32
48	Cysteine-rich secretory protein 3 overexpression is linked to a subset of PTEN-deleted ERG fusion-positive prostate cancers with early biochemical recurrence. <i>Modern Pathology</i> , 2013 , 26, 733-42	9.8	32
47	Genomic deletion of MAP3K7 at 6q12-22 is associated with early PSA recurrence in prostate cancer and absence of TMPRSS2:ERG fusions. <i>Modern Pathology</i> , 2013 , 26, 975-83	9.8	121
46	Marked heterogeneity of ERG expression in large primary prostate cancers. <i>Modern Pathology</i> , 2013 , 26, 106-16	9.8	56
45	Loss of CDKN1B/p27Kip1 expression is associated with ERG fusion-negative prostate cancer, but is unrelated to patient prognosis. <i>Oncology Letters</i> , 2013 , 6, 1245-1252	2.6	18
44	CHD1 is a 5q21 tumor suppressor required for ERG rearrangement in prostate cancer. <i>Cancer Research</i> , 2013 , 73, 2795-805	10.1	169
43	Strong expression of the neuronal transcription factor FOXP2 is linked to an increased risk of early PSA recurrence in ERG fusion-negative cancers. <i>Journal of Clinical Pathology</i> , 2013 , 66, 563-8	3.9	25

42	High c-MET expression is frequent but not associated with early PSA recurrence in prostate cancer. <i>Experimental and Therapeutic Medicine</i> , 2013 , 5, 102-106	2.1	12
41	Value of cell cycle progression (CCP) score to predict biochemical recurrence and definitive post-surgical pathology.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 5043-5043	2.2	2
40	Neurovascular structure-adjacent frozen-section examination (NeuroSAFE) increases nerve-sparing frequency and reduces positive surgical margins in open and robot-assisted laparoscopic radical prostatectomy: experience after 11,069 consecutive patients. <i>European Urology</i> , 2012 , 62, 333-40	10.2	162
39	Risk assessment of metastatic recurrence in patients with prostate cancer by using the Cancer of the Prostate Risk Assessment score: results from 2937 European patients. <i>BJU International</i> , 2012 , 110, 1714-20	5.6	11
38	Peroxiredoxins 3 and 4 are overexpressed in prostate cancer tissue and affect the proliferation of prostate cancer cells in vitro. <i>Journal of Proteome Research</i> , 2012 , 11, 2452-66	5.6	38
37	Genome-wide DNA methylation events in TMPRSS2-ERG fusion-negative prostate cancers implicate an EZH2-dependent mechanism with miR-26a hypermethylation. <i>Cancer Discovery</i> , 2012 , 2, 1024-35	24.4	107
36	Genomic deletion of PTEN is associated with tumor progression and early PSA recurrence in ERG fusion-positive and fusion-negative prostate cancer. <i>American Journal of Pathology</i> , 2012 , 181, 401-12	5.8	256
35	The impact of the number of cores on tissue microarray studies investigating prostate cancer biomarkers. <i>International Journal of Oncology</i> , 2012 , 40, 261-8	4.4	21
34	Human prostate cancer in a clinically relevant xenograft mouse model: identification of (1,6)-branched oligosaccharides as a marker of tumor progression. <i>Clinical Cancer Research</i> , 2012 , 18, 1364-73	12.9	64
33	Association of nuclear accumulation of p53 with ERG fusion and poor prognosis in prostate cancer.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 124-124	2.2	
32	Inverse stage migration in patients undergoing radical prostatectomy: results of 8916 European patients treated within the last decade. <i>BJU International</i> , 2011 , 108, 1256-61	5.6	92
31	MicroRNA in prostate, bladder, and kidney cancer: a systematic review. <i>European Urology</i> , 2011 , 59, 671-80.	10.2	355
30	Full functional-length urethral sphincter preservation during radical prostatectomy. <i>European Urology</i> , 2011 , 60, 320-9	10.2	138
29	TMPRSS2-ERG -specific transcriptional modulation is associated with prostate cancer biomarkers and TGF- β signaling. <i>BMC Cancer</i> , 2011 , 11, 507	4.8	117
28	High level PSMA expression is associated with early PSA recurrence in surgically treated prostate cancer. <i>Prostate</i> , 2011 , 71, 281-8	4.2	179
27	Circulating miRNAs are correlated with tumor progression in prostate cancer. <i>International Journal of Cancer</i> , 2011 , 128, 608-16	7.5	448
26	ERG status is unrelated to PSA recurrence in radically operated prostate cancer in the absence of antihormonal therapy. <i>Clinical Cancer Research</i> , 2011 , 17, 5878-88	12.9	225
25	Identification of clinically relevant protein targets in prostate cancer with 2D-DIGE coupled mass spectrometry and systems biology network platform. <i>PLoS ONE</i> , 2011 , 6, e16833	3.7	55

24	The presence of prostate cancer on saturation biopsy can be accurately predicted. <i>BJU International</i> , 2010 , 105, 636-41	5.6	16
23	Chromosome 8p deletions and 8q gains are associated with tumor progression and poor prognosis in prostate cancer. <i>Clinical Cancer Research</i> , 2010 , 16, 56-64	12.9	98
22	Low level HER2 overexpression is associated with rapid tumor cell proliferation and poor prognosis in prostate cancer. <i>Clinical Cancer Research</i> , 2010 , 16, 1553-60	12.9	113
21	Biochemical recurrence after radical prostatectomy: multiplicative interaction between surgical margin status and pathological stage. <i>Journal of Urology</i> , 2010 , 184, 1341-6	2.5	68
20	Long-term data on the survival of patients with prostate cancer treated with radical prostatectomy in the prostate-specific antigen era. <i>BJU International</i> , 2010 , 106, 37-43	5.6	60
19	Head-to-head comparison of the three most commonly used preoperative models for prediction of biochemical recurrence after radical prostatectomy. <i>European Urology</i> , 2010 , 57, 562-8	10.2	58
18	From gene to clinic: TMA-based clinical validation of molecular markers in prostate cancer. <i>Methods in Molecular Biology</i> , 2010 , 664, 177-89	1.4	8
17	Molecular cancer phenotype in normal prostate tissue. <i>European Urology</i> , 2009 , 55, 885-90	10.2	21
16	Prevalence of a tertiary Gleason grade and its impact on adverse histopathologic parameters in a contemporary radical prostatectomy series. <i>European Urology</i> , 2009 , 55, 394-401	10.2	31
15	Current technique of open intrafascial nerve-sparing retropubic prostatectomy. <i>European Urology</i> , 2009 , 56, 317-24	10.2	102
14	Immunological microenvironment in prostate cancer: high mast cell densities are associated with favorable tumor characteristics and good prognosis. <i>Prostate</i> , 2009 , 69, 976-81	4.2	100
13	TMPRSS2:ERG fusion transcripts in urine from prostate cancer patients correlate with a less favorable prognosis. <i>Apmis</i> , 2009 , 117, 575-82	3.4	59
12	Predictive value of prostate-specific antigen expression in prostate cancer: a tissue microarray study. <i>Urology</i> , 2009 , 74, 1169-73	1.6	17
11	Clinical significance of p53 alterations in surgically treated prostate cancers. <i>Modern Pathology</i> , 2008 , 21, 1371-8	9.8	165
10	Contemporary prostate cancer prevalence among T1c biopsy-referred men with a prostate-specific antigen level <i>European Urology</i> , 2008 , 53, 750-7	10.2	19
9	Assessment of biochemical recurrence rate in patients with pathologically confirmed insignificant prostate cancer. <i>Urology</i> , 2008 , 72, 1208-11; discussion 1212-3	1.6	10
8	Distinct subcellular expression patterns of neutral endopeptidase (CD10) in prostate cancer predict diverging clinical courses in surgically treated patients. <i>Clinical Cancer Research</i> , 2008 , 14, 7838-42	12.9	55
7	Currently used criteria for active surveillance in men with low-risk prostate cancer: an analysis of pathologic features. <i>Cancer</i> , 2008 , 113, 2068-72	6.4	88

6	Development and external validation of an extended repeat biopsy nomogram. <i>Journal of Urology</i> , 2007 , 177, 510-5	2.5	72
5	Molecular staging of prostate cancer in the year 2007. <i>World Journal of Urology</i> , 2007 , 25, 19-30	4	35
4	Clinical significance of epidermal growth factor receptor protein overexpression and gene copy number gains in prostate cancer. <i>Clinical Cancer Research</i> , 2007 , 13, 6579-84	12.9	128
3	The 2002 AJCC pT2 substages confer no prognostic information on the rate of biochemical recurrence after radical prostatectomy. <i>European Urology</i> , 2006 , 49, 273-8; discussion 278-9	10.2	41
2	Significant upgrading affects a third of men diagnosed with prostate cancer: predictive nomogram and internal validation. <i>BJU International</i> , 2006 , 98, 329-34	5.6	110
1	Random forest-based modelling to detect biomarkers for prostate cancer progression		1