

Frank J Snoek

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2185682/publications.pdf>

Version: 2024-02-01

270
papers

15,081
citations

17440

63
h-index

22832

112
g-index

282
all docs

282
docs citations

282
times ranked

12894
citing authors

#	ARTICLE	IF	CITATIONS
1	Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. <i>Diabetologia</i> , 2006, 49, 837-845.	6.3	815
2	Psychosocial problems and barriers to improved diabetes management: results of the Cross-National Diabetes Attitudes, Wishes and Needs (DAWN) Study. <i>Diabetic Medicine</i> , 2005, 22, 1379-1385.	2.3	728
3	Resistance to Insulin Therapy Among Patients and Providers: Results of the cross-national Diabetes Attitudes, Wishes, and Needs (DAWN) study. <i>Diabetes Care</i> , 2005, 28, 2673-2679.	8.6	709
4	Diabetes-related emotional distress in Dutch and U.S. diabetic patients: cross-cultural validity of the problem areas in diabetes scale.. <i>Diabetes Care</i> , 2000, 23, 1305-1309.	8.6	361
5	Constructs of depression and distress in diabetes: time for an appraisal. <i>Lancet Diabetes and Endocrinology</i> , 2015, 3, 450-460.	11.4	304
6	Short-form measures of diabetes-related emotional distress: the Problem Areas in Diabetes Scale (PAID)-5 and PAID-1. <i>Diabetologia</i> , 2010, 53, 66-69.	6.3	290
7	Effect of interventions for major depressive disorder and significant depressive symptoms in patients with diabetes mellitus: a systematic review and meta-analysis. <i>General Hospital Psychiatry</i> , 2010, 32, 380-395.	2.4	290
8	The prevalence of diabetes-specific emotional distress in people with Type 2 diabetes: a systematic review and meta-analysis. <i>Diabetic Medicine</i> , 2017, 34, 1508-1520.	2.3	285
9	Continuous glucose monitoring for patients with type 1 diabetes and impaired awareness of hypoglycaemia (IN CONTROL): a randomised, open-label, crossover trial. <i>Lancet Diabetes and Endocrinology</i> , 2016, 4, 893-902.	11.4	284
10	The Management of Type 1 Diabetes in Adults. A Consensus Report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). <i>Diabetes Care</i> , 2021, 44, 2589-2625.	8.6	244
11	Validation of the WHO-5 Well-Being Index in Adolescents With Type 1 Diabetes. <i>Diabetes Care</i> , 2007, 30, 2003-2006.	8.6	241
12	Do guided internet-based interventions result in clinically relevant changes for patients with depression? An individual participant data meta-analysis. <i>Clinical Psychology Review</i> , 2018, 63, 80-92.	11.4	239
13	Symptoms and Well-Being in Relation to Glycemic Control in Type II Diabetes. <i>Diabetes Care</i> , 1996, 19, 204-210.	8.6	235
14	Diabetes structured self-management education programmes: a narrative review and current innovations. <i>Lancet Diabetes and Endocrinology</i> , 2018, 6, 130-142.	11.4	233
15	Monitoring and Discussing Health-Related Quality of Life in Adolescents With Type 1 Diabetes Improve Psychosocial Well-Being. <i>Diabetes Care</i> , 2008, 31, 1521-1526.	8.6	207
16	Web-Based Depression Treatment for Type 1 and Type 2 Diabetic Patients. <i>Diabetes Care</i> , 2011, 34, 320-325.	8.6	184
17	Diabetic encephalopathy: a concept in need of a definition. <i>Diabetologia</i> , 2006, 49, 1447-1448.	6.3	176
18	A Randomized Trial of Continuous Subcutaneous Insulin Infusion and Intensive Injection Therapy in Type 1 Diabetes for Patients With Long-Standing Poor Glycemic Control. <i>Diabetes Care</i> , 2002, 25, 2074-2080.	8.6	174

#	ARTICLE	IF	CITATIONS
19	Rates and risks for co-morbid depression in patients with Type 2 diabetes mellitus: results from a community-based study. <i>Diabetologia</i> , 2003, 46, 892-898.	6.3	174
20	Development of a Type 2 Diabetes Symptom Checklist: a Measure of Symptom Severity. <i>Diabetic Medicine</i> , 1994, 11, 253-261.	2.3	164
21	Psychometric and screening properties of the WHO's well-being index in adult outpatients with Type 1 or Type 2 diabetes mellitus. <i>Diabetic Medicine</i> , 2013, 30, e63-9.	2.3	158
22	Voxel-based morphometry demonstrates reduced grey matter density on brain MRI in patients with diabetic retinopathy. <i>Diabetologia</i> , 2006, 49, 2474-2480.	6.3	156
23	The Confidence in Diabetes Self-Care Scale: Psychometric properties of a new measure of diabetes-specific self-efficacy in Dutch and U.S. patients with type 1 diabetes. <i>Diabetes Care</i> , 2003, 26, 713-718.	8.6	147
24	A 24-Week, Randomized, Treat-to-Target Trial Comparing Initiation of Insulin Glargine Once-Daily With Insulin Detemir Twice-Daily in Patients With Type 2 Diabetes Inadequately Controlled on Oral Glucose-Lowering Drugs. <i>Diabetes Care</i> , 2010, 33, 1176-1178.	8.6	145
25	Efficacy of a Web-Based Intervention With Mobile Phone Support in Treating Depressive Symptoms in Adults With Type 1 and Type 2 Diabetes: A Randomized Controlled Trial. <i>Diabetes Care</i> , 2015, 38, 776-783.	8.6	143
26	Effect of physical activity and/or healthy eating on GDM risk: The DALI Lifestyle Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, jc.2016-3455.	3.6	140
27	Diabetes-specific emotional distress mediates the association between depressive symptoms and glycaemic control in Type 1 and Type 2 diabetes. <i>Diabetic Medicine</i> , 2010, 27, 798-803.	2.3	136
28	Barriers to insulin initiation and intensification and how to overcome them. <i>International Journal of Clinical Practice</i> , 2009, 63, 6-10.	1.7	134
29	Prevalence of comorbid depression is high in outpatients with Type 1 or Type 2 diabetes mellitus. Results from three outpatient clinics in the Netherlands. <i>Diabetic Medicine</i> , 2010, 27, 217-224.	2.3	131
30	Does Internet-based guided-self-help for depression cause harm? An individual participant data meta-analysis on deterioration rates and its moderators in randomized controlled trials. <i>Psychological Medicine</i> , 2016, 46, 2679-2693.	4.5	129
31	The management of type 1 diabetes in adults. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). <i>Diabetologia</i> , 2021, 64, 2609-2652.	6.3	128
32	Epidemiology of gestational diabetes mellitus according to IADPSG/WHO 2013 criteria among obese pregnant women in Europe. <i>Diabetologia</i> , 2017, 60, 1913-1921.	6.3	117
33	Insulin-treated diabetes patients with fear of self-injecting or fear of self-testing. <i>Journal of Psychosomatic Research</i> , 2001, 51, 665-672.	2.6	115
34	Nurses' recognition and registration of depression, anxiety and diabetes-specific emotional problems in outpatients with diabetes mellitus. <i>Patient Education and Counseling</i> , 2006, 60, 235-240.	2.2	114
35	Monitoring of Psychological Well-Being in Outpatients With Diabetes: Effects on mood, HbA1c, and the patient's evaluation of the quality of diabetes care: a randomized controlled trial. <i>Diabetes Care</i> , 2001, 24, 1929-1935.	8.6	111
36	Resting-State Brain Networks in Type 1 Diabetic Patients With and Without Microangiopathy and Their Relation to Cognitive Functions and Disease Variables. <i>Diabetes</i> , 2012, 61, 1814-1821.	0.6	109

#	ARTICLE	IF	CITATIONS
37	Development and validation of the insulin treatment appraisal scale (ITAS) in patients with type 2 diabetes. <i>Health and Quality of Life Outcomes</i> , 2007, 5, 69.	2.4	107
38	Cognitive performance in type 1 diabetes patients is associated with cerebral white matter volume. <i>Diabetologia</i> , 2007, 50, 1763-1769.	6.3	105
39	Monitoring of Individual Needs in Diabetes (MIND): Baseline Data From the Cross-National Diabetes Attitudes, Wishes, and Needs (DAWN) MIND Study. <i>Diabetes Care</i> , 2011, 34, 601-603.	8.6	103
40	Systematic review and meta-analysis of psychological interventions in people with diabetes and elevated diabetes-related distress. <i>Diabetic Medicine</i> , 2018, 35, 1157-1172.	2.3	103
41	Use of Behavioral Change Techniques in Web-Based Self-Management Programs for Type 2 Diabetes Patients: Systematic Review. <i>Journal of Medical Internet Research</i> , 2013, 15, e279.	4.3	100
42	Are patients with type 2 diabetes reluctant to start insulin therapy? An examination of the scope and underpinnings of psychological insulin resistance in a large, international population. <i>Current Medical Research and Opinion</i> , 2011, 27, 1169-1174.	1.9	96
43	Psychological counselling in problematic diabetes: does it help?. <i>Diabetic Medicine</i> , 2002, 19, 265-273.	2.3	95
44	Results From a European Multicenter Randomized Trial of Physical Activity and/or Healthy Eating to Reduce the Risk of Gestational Diabetes Mellitus: The DALI Lifestyle Pilot. <i>Diabetes Care</i> , 2015, 38, 1650-1656.	8.6	93
45	Monitoring of Individual Needs in Diabetes (MIND)-2. <i>Diabetes Care</i> , 2012, 35, 2128-2132.	8.6	92
46	Screening, evaluation and management of depression in people with diabetes in primary care. <i>Primary Care Diabetes</i> , 2013, 7, 1-10.	1.8	91
47	Patient and provider perceptions of care for diabetes: results of the cross-national DAWN Study. <i>Diabetologia</i> , 2006, 49, 279-288.	6.3	89
48	Depression, Anxiety and Glucose Metabolism in the General Dutch Population: The New Hoorn Study. <i>PLoS ONE</i> , 2010, 5, e9971.	2.5	88
49	Fatigue, sleep disturbances and circadian rhythm in multiple sclerosis. <i>Journal of Neurology</i> , 1993, 240, 446-448.	3.6	85
50	DALI: Vitamin D and lifestyle intervention for gestational diabetes mellitus (GDM) prevention: an European multicentre, randomised trial "study protocol. <i>BMC Pregnancy and Childbirth</i> , 2013, 13, 142.	2.4	85
51	Phobia of self-injecting and self-testing in insulin-treated diabetes patients: opportunities for screening. <i>Diabetic Medicine</i> , 2001, 18, 671-674.	2.3	82
52	IADPSG and WHO 2013 Gestational Diabetes Mellitus Criteria Identify Obese Women With Marked Insulin Resistance in Early Pregnancy. <i>Diabetes Care</i> , 2016, 39, e90-e92.	8.6	79
53	Diabetes Fear of Injecting and Self-Testing Questionnaire: a psychometric evaluation. <i>Diabetes Care</i> , 2000, 23, 765-769.	8.6	77
54	Differential associations between depressive symptoms and glycaemic control in outpatients with diabetes. <i>Diabetic Medicine</i> , 2013, 30, e115-22.	2.3	77

#	ARTICLE	IF	CITATIONS
55	Well-Being and Symptoms in Relation to Insulin Therapy in Type 2 Diabetes. <i>Diabetes Care</i> , 1998, 21, 919-924.	8.6	76
56	The 12-item well-being questionnaire. An evaluation of its validity and reliability in Dutch people with diabetes. <i>Diabetes Care</i> , 1999, 22, 2004-2010.	8.6	74
57	Serious diabetes-specific emotional problems and depression in a Croatianâ€“Dutchâ€“English Survey from the European Depression in Diabetes [EDID] Research Consortium. <i>Diabetes Research and Clinical Practice</i> , 2005, 70, 166-173.	2.8	73
58	Symptoms of depression and diabetesâ€“specific emotional distress are associated with a negative appraisal of insulin therapy in insulinâ€“naïve patients with Type 2 diabetes mellitus. A study from the European Depression in Diabetes [EDID] Research Consortium. <i>Diabetic Medicine</i> , 2009, 26, 28-33.	2.3	71
59	Monitoring health related quality of life in adolescents with diabetes: a review of measures. <i>Archives of Disease in Childhood</i> , 2007, 92, 434-439.	1.9	69
60	Substitution of night-time continuous subcutaneous insulin infusion therapy for bedtime NPH insulin in a multiple injection regimen improves counterregulatory hormonal responses and warning symptoms of hypoglycaemia in IDDM. <i>Diabetologia</i> , 1998, 41, 322-329.	6.3	68
61	Health-related quality of life in patients with systemic lupus erythematosus: development and validation of a lupus specific symptom checklist. <i>Quality of Life Research</i> , 2003, 12, 635-644.	3.1	68
62	Cognitive behavioural therapy (CBT) compared with blood glucose awareness training (BGAT) in poorly controlled Type 1 diabetic patients: long-term effects on HbA _{1c} moderated by depression. A randomized controlled trial. <i>Diabetic Medicine</i> , 2008, 25, 1337-1342.	2.3	68
63	Functional Brain Connectivity and Neurocognitive Functioning in Patients With Long-Standing Type 1 Diabetes With and Without Microvascular Complications. <i>Diabetes</i> , 2009, 58, 2335-2343.	0.6	67
64	Persistent poor glycaemic control in adult Type 1 diabetes. A closer look at the problem. <i>Diabetic Medicine</i> , 2004, 21, 1263-1268.	2.3	65
65	Psychometric Evaluation of the Diabetes Symptom Checklist-Revised (DSC-R)â€“A Measure of Symptom Distress. <i>Value in Health</i> , 2009, 12, 1168-1175.	0.3	65
66	Assessing diabetes-related quality of life of youth with type 1 diabetes in routine clinical care: the MIND Youth Questionnaire (MY-Q). <i>Pediatric Diabetes</i> , 2012, 13, 638-646.	2.9	64
67	Microvascular Disease in Type 1 Diabetes Alters Brain Activation: A Functional Magnetic Resonance Imaging Study. <i>Diabetes</i> , 2006, 55, 334-340.	0.6	63
68	Effects of cognitive behavioural group training (CBGT) in adult patients with poorly controlled insulin-dependent (type 1) diabetes: a pilot study. <i>Patient Education and Counseling</i> , 2001, 45, 143-148.	2.2	61
69	Proliferative Retinopathy in Type 1 Diabetes Is Associated With Cerebral Microbleeds, Which Is Part of Generalized Microangiopathy. <i>Diabetes Care</i> , 2014, 37, 1165-1168.	8.6	61
70	Decision aids that facilitate elements of shared decision making in chronic illnesses: a systematic review. <i>Systematic Reviews</i> , 2019, 8, 121.	5.3	60
71	Short-term effects of cognitive behavioural group training (CBGT) in adult Type 1 diabetes patients in prolonged poor glycaemic control. A randomized controlled trial. <i>Diabetic Medicine</i> , 2005, 22, 1619-1623.	2.3	59
72	Self-report and parent-report of physical and psychosocial well-being in Dutch adolescents with type 1 diabetes in relation to glycemic control. <i>Health and Quality of Life Outcomes</i> , 2007, 5, 10.	2.4	59

#	ARTICLE	IF	CITATIONS
73	Efficacy and Moderators of Internet-Based Interventions in Adults with Subthreshold Depression: An Individual Participant Data Meta-Analysis of Randomized Controlled Trials. <i>Psychotherapy and Psychosomatics</i> , 2021, 90, 94-106.	8.8	59
74	Diffusion tensor imaging in type 1 diabetes: decreased white matter integrity relates to cognitive functions. <i>Diabetologia</i> , 2012, 55, 1218-1220.	6.3	58
75	The Well-being Questionnaire: evidence for a three-factor structure with 12 items (W-BQ12). <i>Psychological Medicine</i> , 2000, 30, 455-462.	4.5	57
76	Examining the Behaviour subscale of the Hypoglycaemia Fear Survey: an international study. <i>Diabetic Medicine</i> , 2013, 30, 603-609.	2.3	57
77	Follow-up results on monitoring and discussing health-related quality of life in adolescent diabetes care: benefits do not sustain in routine practice. <i>Pediatric Diabetes</i> , 2010, 11, 175-181.	2.9	56
78	Diabetes-specific emotional distress in people with Type 2 diabetes: a comparison between primary and secondary care. <i>Diabetic Medicine</i> , 2014, 31, 1252-1259.	2.3	56
79	Limited effect of screening for depression with written feedback in outpatients with diabetes mellitus: a randomised controlled trial. <i>Diabetologia</i> , 2011, 54, 741-748.	6.3	54
80	Associations between depressive symptoms and insulin resistance: The Hoorn Study. <i>Diabetologia</i> , 2006, 49, 2874-2877.	6.3	53
81	Perceived risk for Type 2 diabetes in participants in a stepwise population-screening programme. <i>Diabetic Medicine</i> , 2003, 20, 210-215.	2.3	52
82	Efficacy and cost-effectiveness of a web-based intervention with mobile phone support to treat depressive symptoms in adults with diabetes mellitus type 1 and type 2: design of a randomised controlled trial. <i>BMC Psychiatry</i> , 2013, 13, 306.	2.6	52
83	Assessing impaired hypoglycemia awareness in type 1 diabetes: agreement of self-report but not of field study data with the autonomic symptom threshold during experimental hypoglycemia. <i>Diabetes Care</i> , 2000, 23, 529-532.	8.6	51
84	Sleep and HbA1c in Patients With Type 2 Diabetes: Which Sleep Characteristics Matter Most?. <i>Diabetes Care</i> , 2020, 43, 235-243.	8.6	51
85	Development and validation of the diabetes fear of injecting and self-testing questionnaire (D-FISQ): first findings. , 1997, 14, 871-876.		50
86	A reduction in sedentary behaviour in obese women during pregnancy reduces neonatal adiposity: the DALI randomised controlled trial. <i>Diabetologia</i> , 2019, 62, 915-925.	6.3	50
87	Association between symptoms of depression and glycaemic control may be unstable across gender. <i>Diabetic Medicine</i> , 2001, 18, 595-598.	2.3	49
88	Symptoms of depression in people with impaired glucose metabolism or Type 2 diabetes mellitus: The Hoorn Study. <i>Diabetic Medicine</i> , 2008, 25, 843-849.	2.3	49
89	Eicosapentaenoic acid as an add-on to antidepressant medication for co-morbid major depression in patients with diabetes mellitus: A randomized, double-blind placebo-controlled study. <i>Journal of Affective Disorders</i> , 2010, 126, 282-286.	4.1	49
90	The psychological impact of screening for type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2006, 22, 20-25.	4.0	48

#	ARTICLE	IF	CITATIONS
91	Diabetes-Related Symptom Distress in Association With Glucose Metabolism and Comorbidity: The Hoorn Study. <i>Diabetes Care</i> , 2008, 31, 2268-2270.	8.6	47
92	The cognitive and psychological effects of living with type 1 diabetes: a narrative review. <i>Diabetic Medicine</i> , 2020, 37, 555-563.	2.3	47
93	Internet-Based Interventions in Chronic Somatic Disease. <i>Deutsches A&#x0308;rztblatt International</i> , 2018, 115, 659-665.	0.9	47
94	Toward Defining a Cutoff Score for Elevated Fear of Hypoglycemia on the Hypoglycemia Fear Survey Worry Subscale in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2014, 37, 102-108.	8.6	46
95	Role of Continuous Glucose Monitoring in Clinical Trials: Recommendations on Reporting. <i>Diabetes Technology and Therapeutics</i> , 2017, 19, 391-399.	4.4	45
96	Depressive symptoms and unmet psychological needs of Dutch youth with type 1 diabetes: results of a web-survey. <i>Pediatric Diabetes</i> , 2011, 12, 172-176.	2.9	44
97	Screening for Type 2 diabetes: an exploration of subjectsâ€™ perceptions regarding diagnosis and procedure. <i>Diabetic Medicine</i> , 2002, 19, 406-411.	2.3	43
98	Web-based cognitive behavioural therapy (W-CBT) for diabetes patients with co-morbid depression: Design of a randomised controlled trial. <i>BMC Psychiatry</i> , 2008, 8, 9.	2.6	43
99	The DALI vitamin D randomized controlled trial for gestational diabetes mellitus prevention: No major benefit shown besides vitamin D sufficiency. <i>Clinical Nutrition</i> , 2020, 39, 976-984.	5.0	42
100	Web-based intervention for depressive symptoms in adults with types 1 and 2 diabetes mellitus: a health economic evaluation. <i>British Journal of Psychiatry</i> , 2018, 212, 199-206.	2.8	41
101	Cerebrospinal fluid levels of Alzheimerâ€™s disease biomarkers in middle-aged patients with type 1 diabetes. <i>Diabetologia</i> , 2014, 57, 2208-2214.	6.3	40
102	Continuous intraperitoneal insulin infusion in patients with â€˜brittleâ€™ diabetes: favourable effects on glycaemic control and hospital stay. <i>Diabetic Medicine</i> , 2002, 19, 496-501.	2.3	39
103	Diabetes-related symptoms and negative mood in participants of a targeted population-screening program for type 2 diabetes: The Hoorn Screening study. <i>Quality of Life Research</i> , 2005, 14, 1501-1509.	3.1	39
104	Development and reach of a web-based cognitive behavioural therapy programme to reduce symptoms of depression and diabetes-specific distress. <i>Patient Education and Counseling</i> , 2011, 84, 49-55.	2.2	39
105	Health-related quality of life in the first year following diagnosis of Type 2 diabetes: newly diagnosed patients in general practice compared with screening-detected patients. <i>The Hoorn Screening Study. Diabetic Medicine</i> , 2004, 21, 1075-1081.	2.3	38
106	No substantial psychological impact of the diagnosis of Type 2 diabetes following targeted population screening: The Hoorn Screening Study. <i>Diabetic Medicine</i> , 2004, 21, 992-998.	2.3	37
107	Physical activity, depressed mood and pregnancy worries in European obese pregnant women: results from the DALI study. <i>BMC Pregnancy and Childbirth</i> , 2015, 15, 158.	2.4	36
108	Youth With Type 1 Diabetes Taking Responsibility for Self-Management: The Importance of Executive Functioning in Achieving Glycemic Control. <i>Diabetes Care</i> , 2019, 42, 225-231.	8.6	36

#	ARTICLE	IF	CITATIONS
109	Effectiveness of HypoAware, a Brief Partly Web-Based Psychoeducational Intervention for Adults With Type 1 and Insulin-Treated Type 2 Diabetes and Problematic Hypoglycemia: A Cluster Randomized Controlled Trial. <i>Diabetes Care</i> , 2016, 39, 2190-2196.	8.6	35
110	Randomized Study of Two Different Target Levels of Glycemic Control Within the Acceptable Range in Type 2 Diabetes: Effects on well-being at 1 year. <i>Diabetes Care</i> , 1998, 21, 2085-2093.	8.6	34
111	Cost-effectiveness of healthy eating and/or physical activity promotion in pregnant women at increased risk of gestational diabetes mellitus: economic evaluation alongside the DALI study, a European multicenter randomized controlled trial. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2018, 15, 23.	4.6	34
112	Back to the future: 25 years of "Guidelines for encouraging psychological well-being" among people affected by diabetes. <i>Diabetic Medicine</i> , 2020, 37, 1225-1229.	2.3	34
113	Altered eigenvector centrality is related to local resting-state network functional connectivity in patients with longstanding type 1 diabetes mellitus. <i>Human Brain Mapping</i> , 2017, 38, 3623-3636.	3.6	33
114	Initiation of insulin glargine in patients with Type 2 diabetes in suboptimal glycaemic control positively impacts health-related quality of life. A prospective cohort study in primary care. <i>Diabetic Medicine</i> , 2011, 28, 1096-1102.	2.3	31
115	Ventral Striatum, but Not Cortical Volume Loss, Is Related to Cognitive Dysfunction in Type 1 Diabetic Patients With and Without Microangiopathy. <i>Diabetes Care</i> , 2014, 37, 2483-2490.	8.6	31
116	Beliefs, Barriers, and Preferences of European Overweight Women to Adopt a Healthier Lifestyle in Pregnancy to Minimize Risk of Developing Gestational Diabetes Mellitus: An Explorative Study. <i>Journal of Pregnancy</i> , 2016, 2016, 1-11.	2.4	31
117	Diabetes distress is associated with adverse pregnancy outcomes in women with gestational diabetes: a prospective cohort study. <i>BMC Pregnancy and Childbirth</i> , 2019, 19, 223.	2.4	31
118	Cognitive behavioural group training (CBGT) for patients with type 1 diabetes in persistent poor glycaemic control: who do we reach?. <i>Patient Education and Counseling</i> , 2005, 56, 313-322.	2.2	30
119	Do physicians understand Type 2 diabetes patients' perceptions of seriousness; the emotional impact and needs for care improvement? A cross-national survey. <i>Patient Education and Counseling</i> , 2011, 85, 258-263.	2.2	29
120	Uptake and Effects of the e-Vita Personal Health Record with Self-Management Support and Coaching, for Type 2 Diabetes Patients Treated in Primary Care. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-9.	2.3	29
121	Light therapy: is it safe for the eyes?. <i>Acta Psychiatrica Scandinavica</i> , 2017, 136, 534-548.	4.5	29
122	Is a Severe Clinical Profile an Effect Modifier in a Web-Based Depression Treatment for Adults With Type 1 or Type 2 Diabetes? Secondary Analyses From a Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2012, 14, e2.	4.3	29
123	Improved glycaemic control in type 1 diabetes patients following participation per se in a clinical trial? mechanisms and implications. <i>Diabetes/Metabolism Research and Reviews</i> , 2003, 19, 357-362.	4.0	28
124	Clinical Utility of SMBG: Recommendations on the Use and Reporting of SMBG in Clinical Research. <i>Diabetes Care</i> , 2015, 38, 1627-1633.	8.6	28
125	Parental Diabetes Behaviors and Distress Are Related to Glycemic Control in Youth with Type 1 Diabetes: Longitudinal Data from the DINO Study. <i>Journal of Diabetes Research</i> , 2017, 2017, 1-7.	2.3	28
126	The longitudinal association between glycaemic control and health-related quality of life following insulin therapy optimisation in type 2 diabetes patients. A prospective observational study in secondary care. <i>Quality of Life Research</i> , 2012, 21, 1359-1365.	3.1	27

#	ARTICLE	IF	CITATIONS
127	Fat food for a bad mood. Could we treat and prevent depression in Type 2 diabetes by means of omega-3 polyunsaturated fatty acids? A review of the evidence. <i>Diabetic Medicine</i> , 2005, 22, 1465-1475.	2.3	26
128	Effects of Light Therapy on Mood and Insulin Sensitivity in Patients With Type 2 Diabetes and Depression: Results From a Randomized Placebo-Controlled Trial. <i>Diabetes Care</i> , 2019, 42, 529-538.	8.6	26
129	Hypoglycaemia induces emotional disruption. <i>Patient Education and Counseling</i> , 1996, 29, 117-122.	2.2	25
130	Disrupted subject-specific gray matter network properties and cognitive dysfunction in type 1 diabetes patients with and without proliferative retinopathy. <i>Human Brain Mapping</i> , 2016, 37, 1194-1208.	3.6	25
131	Keeping safe. Continuous glucose monitoring (CGM) in persons with Type 1 diabetes and impaired awareness of hypoglycaemia: a qualitative study. <i>Diabetic Medicine</i> , 2017, 34, 1470-1476.	2.3	24
132	Effectiveness of psychoeducational interventions for the treatment of diabetes-specific emotional distress and glycaemic control in people with type 2 diabetes: A systematic review and meta-analysis. <i>Primary Care Diabetes</i> , 2019, 13, 556-567.	1.8	24
133	Perceptions of risk in adults with a low or high risk profile of developing type 2 diabetes; a cross-sectional population-based study. <i>Patient Education and Counseling</i> , 2008, 73, 307-312.	2.2	23
134	Changing the odds. What do we learn from prevention studies targeted at people with a positive family history of type 2 diabetes?. <i>Primary Care Diabetes</i> , 2011, 5, 215-221.	1.8	23
135	High Diabetes Distress Among Ethnic Minorities Is Not Explained by Metabolic, Cardiovascular, or Lifestyle Factors: Findings From the Dutch Diabetes Pearl Cohort. <i>Diabetes Care</i> , 2018, 41, 1854-1861.	8.6	23
136	Identifying solutions to psychological insulin resistance: An international study. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 307-314.	2.3	23
137	Metabolic phenotypes of early gestational diabetes mellitus and their association with adverse pregnancy outcomes. <i>Diabetic Medicine</i> , 2021, 38, e14413.	2.3	23
138	Ceiling effect reduces the validity of the Diabetes Treatment Satisfaction Questionnaire. <i>Diabetes Care</i> , 1998, 21, 2039-2039.	8.6	22
139	Biological and behavioural determinants of the frequency of mild, biochemical hypoglycaemia in patients with Type 1 diabetes on multiple insulin injection therapy. <i>Diabetes/Metabolism Research and Reviews</i> , 2000, 16, 157-163.	4.0	22
140	Optimized basal-bolus therapy using a fixed mixture of 75% lispro and 25% NPL insulin in type 1 diabetes patients: no favorable effects on glycemic control, physiological responses to hypoglycemia, well-being, or treatment satisfaction. <i>Diabetes Care</i> , 2000, 23, 629-633.	8.6	22
141	Prevalence and Correlates of Sexual Dysfunction in Men and Women With Type 2 Diabetes. <i>Journal of Sex and Marital Therapy</i> , 2015, 41, 680-690.	1.5	22
142	Association between Gestational Weight Gain, Gestational Diabetes Risk, and Obstetric Outcomes: A Randomized Controlled Trial Post Hoc Analysis. <i>Nutrients</i> , 2018, 10, 1568.	4.1	22
143	Improving quality of life in diabetes: how effective is education?. <i>Patient Education and Counseling</i> , 2003, 51, 1-3.	2.2	21
144	Implementation of quality of life monitoring in Dutch routine care of adolescents with type 1 diabetes: appreciated but difficult. <i>Pediatric Diabetes</i> , 2016, 17, 112-119.	2.9	21

#	ARTICLE	IF	CITATIONS
145	Nutritional Lifestyle Intervention in Obese Pregnant Women, Including Lower Carbohydrate Intake, Is Associated With Increased Maternal Free Fatty Acids, 3- β -Hydroxybutyrate, and Fasting Glucose Concentrations: A Secondary Factorial Analysis of the European Multicenter, Randomized Controlled DALI Lifestyle Intervention Trial. <i>Diabetes Care</i> , 2019, 42, 1380-1389.	8.6	21
146	Blood Glucose Awareness Training. , 0, , 169-206.		20
147	Risk factors for hyperglycemia in pregnancy in the DALI study differ by period of pregnancy and OGTT time point. <i>European Journal of Endocrinology</i> , 2018, 179, 39-49.	3.7	20
148	Changes in MEG resting-state networks are related to cognitive decline in type 1 diabetes mellitus patients. <i>NeuroImage: Clinical</i> , 2014, 5, 69-76.	2.7	19
149	Effectiveness of a PLISSIT model intervention in patients with type 2 diabetes mellitus in primary care: design of a cluster-randomised controlled trial. <i>BMC Family Practice</i> , 2015, 16, 69.	2.9	19
150	HypoAware: development and pilot study of a brief and partly web-based psychoeducational group intervention for adults with Type 1 and insulin-treated Type 2 diabetes and problematic hypoglycaemia. <i>Diabetic Medicine</i> , 2016, 33, 184-191.	2.3	19
151	Type 2 Diabetes Patients' Needs and Preferences for Care Concerning Sexual Problems: A Cross-Sectional Survey and Qualitative Interviews. <i>Journal of Sex and Marital Therapy</i> , 2016, 42, 324-337.	1.5	19
152	Disturbed eating behaviors in adolescents with type 1 diabetes. How to screen for yellow flags in clinical practice?. <i>Pediatric Diabetes</i> , 2017, 18, 376-383.	2.9	19
153	The statistical approach in trial-based economic evaluations matters: get your statistics together!. <i>BMC Health Services Research</i> , 2021, 21, 475.	2.2	19
154	Glucose variability and mood in adults with diabetes: A systematic review. <i>Endocrinology, Diabetes and Metabolism</i> , 2021, 4, e00152.	2.4	19
155	DiAlert: a lifestyle education programme aimed at people with a positive family history of type 2 diabetes and overweight, study protocol of a randomised controlled trial. <i>BMC Public Health</i> , 2011, 11, 751.	2.9	18
156	Rationale, Design, and Baseline Data of the Insulin Glargine (Lantus) Versus Insulin Detemir (Levemir) Treat-To-Target (L2T3) Study: A Multinational, Randomized Noninferiority Trial of Basal Insulin Initiation in Type 2 Diabetes. <i>Diabetes Technology and Therapeutics</i> , 2009, 11, 739-743.	4.4	17
157	Effects of the PRo-active Interdisciplinary Self-MANagement (PRISMA, Dutch DESMOND) program on dietary intake in type 2 diabetes outpatients: A pilot study. <i>Clinical Nutrition</i> , 2010, 29, 199-205.	5.0	17
158	Light therapy for better mood and insulin sensitivity in patients with major depression and type 2 diabetes: a randomised, double-blind, parallel-arm trial. <i>BMC Psychiatry</i> , 2015, 15, 169.	2.6	17
159	A systematic review of decision aids that facilitate elements of shared decision-making in chronic illnesses: a review protocol. <i>Systematic Reviews</i> , 2017, 6, 155.	5.3	17
160	Conversations and Reactions Around Severe Hypoglycemia Study: Results of Hypoglycemia Experiences in Canadian Adults With Insulin-Treated Diabetes and Their Caregivers. <i>Canadian Journal of Diabetes</i> , 2021, 45, 236-242.	0.8	17
161	Breaking the barriers to optimal glycaemic control-what physicians need to know from patients' perspectives. <i>International Journal of Clinical Practice, Supplement</i> , 2002, , 80-4.	0.3	17
162	Expert Panel Guidance and Narrative Review of Treatment Simplification of Complex Insulin Regimens to Improve Outcomes in Type 2 Diabetes. <i>Diabetes Therapy</i> , 2022, 13, 619-634.	2.5	17

#	ARTICLE	IF	CITATIONS
163	Assessment of perceived barriers in self-care of insulinrequiring diabetic patients. <i>Patient Education and Counseling</i> , 1996, 29, 277-281.	2.2	16
164	Diabetes risk reduction in overweight first degree relatives of type 2 diabetes patients: Effects of a low-intensive lifestyle education program (DiAlert) A randomized controlled trial. <i>Patient Education and Counseling</i> , 2015, 98, 476-483.	2.2	16
165	Subgenual Cingulate Cortex Functional Connectivity in Relation to Depressive Symptoms and Cognitive Functioning in Type 1 Diabetes Mellitus Patients. <i>Psychosomatic Medicine</i> , 2016, 78, 740-749.	2.0	16
166	Quality of life and neuropsychological functions in long-term low-grade glioma survivors. <i>International Journal of Radiation Oncology Biology Physics</i> , 1994, 29, 1201-1202.	0.8	15
167	Family communication as strategy in diabetes prevention: An observational study in families with Dutch and Surinamese South-Asian ancestry. <i>Patient Education and Counseling</i> , 2012, 87, 23-29.	2.2	15
168	Health care providersâ€™ perspective on using family history in the prevention of type 2 diabetes: a qualitative study including different disciplines. <i>BMC Family Practice</i> , 2013, 14, 31.	2.9	15
169	Comment on â€œPsychological distress in patients with cancer: is screening the effective solution?â€™. <i>British Journal of Cancer</i> , 2013, 108, 2628-2630.	6.4	15
170	Ethnic Minorities with Diabetes Differ in Depressive and Anxiety Symptoms and Diabetes-Distress. <i>Journal of Diabetes Research</i> , 2017, 2017, 1-11.	2.3	15
171	How has psychoâ€behavioural research advanced our understanding of hypoglycaemia in type 1 diabetes?. <i>Diabetic Medicine</i> , 2019, 37, 409-417.	2.3	15
172	Combined bedtime insulin-daytime sulphonylurea regimen compared with two different daily insulin regimens in type 2 diabetes: effects on HbA1c and hypoglycaemia rate-a randomised trial. <i>Diabetes/Metabolism Research and Reviews</i> , 2003, 19, 148-152.	4.0	14
173	Type 2 diabetes and inheritance: what information do diabetes organizations provide on the Internet?. <i>Diabetic Medicine</i> , 2006, 23, 1233-1238.	2.3	14
174	Assessment of Parent-Adolescent Partnership in Diabetes Care. <i>The Diabetes Educator</i> , 2010, 36, 205-215.	2.5	14
175	Ethnic aspects of emotional distress in patients with diabetes â€œ the Amsterdam Health Monitor Study. <i>Diabetic Medicine</i> , 2013, 30, e25-31.	2.3	14
176	Diabetes IN deVelOpment (DINO): the bio-psychosocial, family functioning and parental well-being of youth with type 1 diabetes: a longitudinal cohort study design. <i>BMC Pediatrics</i> , 2015, 15, 82.	1.7	14
177	Do Nonsuicidal Severely Depressed Individuals with Diabetes Profit from Internet-Based Guided Self-Help? Secondary Analyses of a Pragmatic Randomized Trial. <i>Journal of Diabetes Research</i> , 2019, 2019, 1-11.	2.3	14
178	Performance of early pregnancy HbA1c for predicting gestational diabetes mellitus and adverse pregnancy outcomes in obese European women. <i>Diabetes Research and Clinical Practice</i> , 2020, 168, 108378.	2.8	14
179	Effects of self-monitoring of glucose in non-insulin treated patients with type 2 diabetes: design of the IN CONTROL-trial. <i>BMC Family Practice</i> , 2009, 10, 26.	2.9	13
180	Web-based self-management with and without coaching for type 2 diabetes patients in primary care: design of a randomized controlled trial. <i>BMC Endocrine Disorders</i> , 2013, 13, 53.	2.2	13

#	ARTICLE	IF	CITATIONS
181	HypoAware-a brief and partly web-based psycho-educational group intervention for adults with type 1 and insulin-treated type 2 diabetes and problematic hypoglycaemia: design of a cost-effectiveness randomised controlled trial. <i>BMC Endocrine Disorders</i> , 2015, 15, 43.	2.2	13
182	Continuous Glucose Monitoring in Patients with Type 1 Diabetes and Impaired Awareness of Hypoglycemia: Also Effective in Patients with Psychological Distress?. <i>Diabetes Technology and Therapeutics</i> , 2017, 19, 595-599.	4.4	13
183	How to identify clinically significant diabetes distress using the Problem Areas in Diabetes (PAID) scale in adults with diabetes treated in primary or secondary care? Evidence for new cut points based on latent class analyses. <i>BMJ Open</i> , 2022, 12, e056304.	1.9	13
184	No evidence for increased self-reported cognitive failure in Type 1 and Type 2 diabetes: a cross-sectional study. <i>Diabetic Medicine</i> , 2007, 24, 735-740.	2.3	12
185	Illness representations of type 2 diabetes patients are associated with perceptions of diabetes threat in relatives. <i>Journal of Health Psychology</i> , 2014, 19, 358-368.	2.3	12
186	Key factors for overcoming psychological insulin resistance: an examination of patient perspectives through content analysis. <i>BMJ Open Diabetes Research and Care</i> , 2019, 7, e000723.	2.8	12
187	Less sedentary time is associated with a more favourable glucose-insulin axis in obese pregnant women—a secondary analysis of the DALI study. <i>International Journal of Obesity</i> , 2021, 45, 296-307.	3.4	12
188	Clinical assessment of emotions in patients with cancer: Diagnostic accuracy compared to two reference standards.. <i>Journal of Clinical Oncology</i> , 2017, 35, 24-24.	1.6	12
189	Web-based guided insulin self-titration in patients with type 2 diabetes: the Di@log study. Design of a cluster randomised controlled trial [TC1316]. <i>BMC Family Practice</i> , 2009, 10, 40.	2.9	11
190	Design and rationale of the IN CONTROL trial: the effects of real-time continuous glucose monitoring on glycemia and quality of life in patients with type 1 diabetes mellitus and impaired awareness of hypoglycemia. <i>BMC Endocrine Disorders</i> , 2015, 15, 42.	2.2	11
191	Correlates of poor mental health in early pregnancy in obese European women. <i>BMC Pregnancy and Childbirth</i> , 2017, 17, 404.	2.4	11
192	Improved diabetes medication convenience and satisfaction in persons with type 2 diabetes after switching to insulin glargine 300 U/mL: results of the observational OPTIN-D study. <i>BMJ Open Diabetes Research and Care</i> , 2018, 6, e000548.	2.8	11
193	Terminology matters: “diabulimia” is insufficient to describe eating disorders in individuals with Type 1 diabetes. <i>Diabetic Medicine</i> , 2020, 37, 1075-1076.	2.3	11
194	Temporal relationships between maternal metabolic parameters with neonatal adiposity in women with obesity differ by neonatal sex: Secondary analysis of the DALI study. <i>Pediatric Obesity</i> , 2020, 15, e12628.	2.8	11
195	The importance of maternal insulin resistance throughout pregnancy on neonatal adiposity. <i>Paediatric and Perinatal Epidemiology</i> , 2021, 35, 83-91.	1.7	11
196	Does low well-being modify the effects of PRISMA (Dutch DESMOND), a structured self-management-education program for people with type 2 diabetes?. <i>Primary Care Diabetes</i> , 2016, 10, 103-110.	1.8	10
197	Does hypoglycaemia affect the improvement in QoL after the transition to insulin in people with type 2 diabetes?. <i>Journal of Endocrinological Investigation</i> , 2018, 41, 249-258.	3.3	10
198	Low Self-Confidence and Diabetes Mismanagement in Youth with Type 1 Diabetes Mediate the Relationship between Behavioral Problems and Elevated HbA1c. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-6.	2.3	9

#	ARTICLE	IF	CITATIONS
199	Screening for sexual dissatisfaction among people with type 2 diabetes in primary care. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1614-1619.	2.3	9
200	Accelerated executive functions decline and gray matter structural changes in middle-aged type 1 diabetes mellitus patients with proliferative retinopathy. <i>Journal of Diabetes</i> , 2018, 10, 835-846.	1.8	9
201	"I am pregnant and my husband has diabetes. Is there a risk for my child?" A qualitative study of questions asked by email about the role of genetic susceptibility to diabetes. <i>BMC Public Health</i> , 2010, 10, 688.	2.9	8
202	Patients' intentions to inform relatives about Type 2 diabetes risk: the role of worry in the process of family risk disclosure. <i>Diabetic Medicine</i> , 2012, 29, e461-7.	2.3	8
203	The presence of cerebral white matter lesions and lower skin microvascular perfusion predicts lower cognitive performance in type 1 diabetes patients with retinopathy but not in healthy controls: A longitudinal study. <i>Microcirculation</i> , 2019, 26, e12530.	1.8	8
204	The Effects of Lifestyle and/or Vitamin D Supplementation Interventions on Pregnancy Outcomes: What Have We Learned from the DALI Studies?. <i>Current Diabetes Reports</i> , 2019, 19, 162.	4.2	8
205	Management of chronic cardiometabolic conditions and mental health during COVID-19. <i>Primary Care Diabetes</i> , 2021, 15, 21-23.	1.8	8
206	Self management of type 2 diabetes. <i>BMJ: British Medical Journal</i> , 2007, 335, 458-459.	2.3	7
207	Effects of self-monitoring of glucose on distress and self-efficacy in people with non-insulin-treated Type 2 diabetes: a randomized controlled trial. <i>Diabetic Medicine</i> , 2016, 33, 537-546.	2.3	7
208	Comparing the EQ-5D-5L crosswalks and value sets for England, the Netherlands and Spain: Exploring their impact on cost-utility results. <i>Health Economics (United Kingdom)</i> , 2020, 29, 640-651.	1.7	7
209	Diabetes and psychological well-being: crossing borders to achieve optimum care. <i>Practical Diabetes International: the International Journal for Diabetes Care Teams Worldwide</i> , 2006, 23, 241-242.	0.2	6
210	The DAWN MIND Youth program. <i>Pediatric Diabetes</i> , 2009, 10, 46-49.	2.9	6
211	Differential impact of subclinical carotid artery disease on cerebral structure and functioning in type 1 diabetes patients with versus those without proliferative retinopathy. <i>Cardiovascular Diabetology</i> , 2014, 13, 58.	6.8	6
212	Psychometric evaluation of the Korean version of the Diabetes Symptom Checklist-Revised (DSC-R) for Patients with Type 2 Diabetes. <i>Health and Quality of Life Outcomes</i> , 2014, 12, 77.	2.4	6
213	Is a motivational interviewing based lifestyle intervention for obese pregnant women across Europe implemented as planned? Process evaluation of the DALI study. <i>BMC Pregnancy and Childbirth</i> , 2017, 17, 293.	2.4	6
214	Comment on Young-Hyman et al. Psychosocial Care for People With Diabetes: A Position Statement of the American Diabetes Association. <i>Diabetes Care</i> 2016;39:2126-2140. <i>Diabetes Care</i> , 2018, 41, e31-e32.	8.6	6
215	Cost-effectiveness of the psychoeducational blended (group and online) intervention HypoAware compared with usual care for people with Type 1 and insulin-treated Type 2 diabetes with problematic hypoglycaemia: analyses of a cluster-randomized controlled trial. <i>Diabetic Medicine</i> , 2018, 35, 214-222.	2.3	6
216	The challenge of treating comorbid mental health problems in patients with a somatic illness. <i>Lancet Psychiatry</i> , 2018, 5, 465.	7.4	6

#	ARTICLE	IF	CITATIONS
217	Mediators of Lifestyle Behaviour Changes in Obese Pregnant Women. Secondary Analyses from the DALI Lifestyle Randomised Controlled Trial. <i>Nutrients</i> , 2019, 11, 311.	4.1	6
218	Improving interpretability of individual Diabetes Symptom Checklist-Revised (DSC-R) scores: the role of patient characteristics. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001146.	2.8	6
219	Improvement in Patient-Reported Outcomes in Adults with Type 1 Diabetes Treated with Sotagliflozin plus Insulin Versus Insulin Alone. <i>Diabetes Technology and Therapeutics</i> , 2021, 23, 70-77.	4.4	6
220	The unexplored role of sedentary time and physical activity in glucose and lipid metabolism-related placental mRNAs in pregnant women who are obese: the DALI lifestyle randomised controlled trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2022, 129, 708-721.	2.3	6
221	Feasibility and user experience of the unguided web-based self-help app "MyDiaMate"™ aimed to prevent and reduce psychological distress and fatigue in adults with diabetes. <i>Internet Interventions</i> , 2021, 25, 100414.	2.7	6
222	Insulin resistance as a marker for the immune-metabolic subtype of depression. <i>Journal of Affective Disorders</i> , 2021, 295, 1371-1376.	4.1	6
223	Re: Vitamin D and gestational diabetes mellitus: a systematic review based on data free of Hawthorne effect. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2018, 125, 1338-1339.	2.3	5
224	Interaction between rs10830962 polymorphism in MTNR1B and lifestyle intervention on maternal and neonatal outcomes: secondary analyses of the DALI lifestyle randomized controlled trial. <i>American Journal of Clinical Nutrition</i> , 2022, 115, 388-396.	4.7	5
225	DiAlert: a prevention program for overweight first degree relatives of type 2 diabetes patients: results of a pilot study to test feasibility and acceptability. <i>Trials</i> , 2012, 13, 178.	1.6	4
226	Validation of a quick screening instrument for measuring fear of hypoglycaemia in persons with diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1360-1361.	2.3	4
227	Clinical assessment of emotions in patients with cancer: Diagnostic accuracy compared with two reference standards. <i>Psycho-Oncology</i> , 2020, 29, 775-780.	2.3	4
228	Data on diabetes-specific distress are needed to improve the quality of diabetes care. <i>Lancet, The</i> , 2021, 397, 2149.	13.7	4
229	Conversations and Reactions Around Severe Hypoglycemia (CRASH): Japan Results From a Global Survey of People with T1DM or Insulin-Treated T2DM and Caregivers. <i>Diabetes Therapy</i> , 2022, 13, 517-533.	2.5	4
230	Conversations and Reactions Around Severe Hypoglycemia (CRASH) Study: Results From People With Diabetes and Caregivers in the United States. <i>Clinical Diabetes</i> , 2022, 40, 477-488.	2.2	4
231	No Differences in Attentional Functioning Between Type 1 Diabetic Patients With and Without a History of Severe Hypoglycemia. <i>Diabetes Care</i> , 1998, 21, 1568-1569.	8.6	3
232	Looking back on 25 years of the PSAD study group. <i>Diabetic Medicine</i> , 2019, 37, 380-382.	2.3	3
233	Key Strategies for Overcoming Psychological Insulin Resistance in Adults with Type 2 Diabetes: The UK Subgroup in the EMOTION Study. <i>Diabetes Therapy</i> , 2020, 11, 1735-1744.	2.5	3
234	Cognitive Functioning and Hippocampal Connectivity in Patients With Longstanding Type 1 Diabetes and Apolipoprotein E ϵ 4. <i>Diabetes Care</i> , 2021, 44, 2388-2396.	8.6	3

#	ARTICLE	IF	CITATIONS
235	Key Factors for Overcoming Psychological Insulin Resistance—An Examination of a Large International Sample through Content Analysis. <i>Diabetes</i> , 2018, 67, .	0.6	3
236	285-OR: Conversations and Reactions around Severe Hypoglycemia (CRASH): U.S. Results from a Global Survey of People with T1DM or Insulin-Treated T2DM and Caregivers. <i>Diabetes</i> , 2019, 68, .	0.6	3
237	Conversations and Reactions Around Severe Hypoglycaemia (CRASH): Results from the German Cohort of a Global Survey of People with Type 1 Diabetes or Insulin-Treated Type 2 Diabetes and Caregivers. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2020, , .	1.2	3
238	Conversations and Reactions Around Severe Hypoglycaemia (CRASH): Spanish results of a global survey of people with type 1 diabetes or insulin-treated type 2 diabetes and caregivers. <i>Endocrinología y Nutrición (English Ed)</i> , 2021, 68, 557-566.	0.2	3
239	Symptoms of depression and insomnia in older age: A within-individual analysis over 20 years. <i>Journal of the American Geriatrics Society</i> , 2022, 70, 2051-2059.	2.6	3
240	#DiabetesPsychologyMatters. <i>Diabetes Spectrum</i> , 2020, 33, 95-98.	1.0	2
241	Effects of Diabetes Prevention Education Program for Overweight and Obese Subjects with a Family History of Type 2 Diabetes Mellitus: A Pilot Study from the United Arab Emirates. <i>Oman Medical Journal</i> , 2021, 36, e268-e268.	1.0	2
242	Conversations and Reactions Around Severe Hypoglycaemia (CRASH): Spanish results of a global survey of people with type 1 diabetes or insulin-treated type 2 diabetes and caregivers. <i>Endocrinología y Nutrición</i> , 2021, 68, 557-566.	0.3	2
243	Biological and behavioural determinants of the frequency of mild, biochemical hypoglycaemia in patients with Type 1 diabetes on multiple insulin injection therapy. <i>Diabetes/Metabolism Research and Reviews</i> , 2000, 16, 157-163.	4.0	2
244	Conversations and Reactions Around Severe Hypoglycemia (CRASH) global survey of people with type-1 diabetes or insulin-treated type-2 diabetes and caregivers: findings from the French cohort. <i>Annales D'Endocrinologie</i> , 2021, 83, 16-16.	1.4	2
245	Evaluatie en bespreking van de kwaliteit van leven van adolescenten met type-1-diabetes. <i>Tijdschrift Voor Kindergeneeskunde</i> , 2008, 76, 180-189.	0.0	1
246	Routine Psychological Screening in Youth With Type 1 Diabetes and Their Parents: a Notion Whose Time Has Come?: Response to Cameron et al.. <i>Diabetes Care</i> , 2008, 31, e14-e14.	8.6	1
247	Symptom burden and its association with change in glucose metabolism status over a 7-year period: the Hoorn Study. <i>Diabetic Medicine</i> , 2014, 31, 747-753.	2.3	1
248	PRM202 - COMPARING THE EQ-5D-5L CROSSWALKS AND VALUE SETS FOR ENGLAND, THE NETHERLANDS AND SPAIN: DO CONCLUSIONS CHANGE?. <i>Value in Health</i> , 2018, 21, S391.	0.3	1
249	Comment on the consensus report on the management of hyperglycaemia in Type 2 diabetes by the American Diabetes Association and the European Association for the Study of Diabetes. <i>Diabetic Medicine</i> , 2019, 36, 911-912.	2.3	1
250	115 - Conversations and Reactions Around Severe Hypoglycemia (CRASH Study): Canadian Results. <i>Canadian Journal of Diabetes</i> , 2019, 43, S40.	0.8	1
251	Cognition in elderly with type 1 diabetes: Is there an interaction between glycemia and aging?. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 4-5.	2.3	1
252	Executive function mediates the link between externalizing behavior and HbA1c in children and adolescents with type 1 diabetes: A cross-national investigation. <i>Pediatric Diabetes</i> , 2021, 22, 503-510.	2.9	1

#	ARTICLE	IF	CITATIONS
253	50-OR: Validation of Fear of Hypoglycemia Screener: Results from the T1D Exchange Registry. <i>Diabetes</i> , 2021, 70, .	0.6	1
254	PS1 - 6. Cerebral resting-state network changes in patients with type 1 diabetes with and without microangiopathy relate to cognitive functions. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2011, 9, 94-94.	0.0	0
255	PS11 - 56. Implementation of quality of life in routine care of adolescents with diabetes mellitus type 1. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2011, 9, 129-129.	0.0	0
256	PS11 - 50. Depressive symptoms exacerbate the negative effects of type 1 diabetes and microangiopathy on brain functioning. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2012, 10, 133-133.	0.0	0
257	PS11 - 51. Prevalence and determinants of diabetes-specific emotional distress: primary care and secondary care patients with type 2 diabetes mellitus compared. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2012, 10, 133-134.	0.0	0
258	PS12 - 3. Use of behavioral change techniques in web-based self-management programs for type 2 diabetes patients: a systematic review. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2013, 11, 167-168.	0.0	0
259	PS12 - 5. Development and evaluation of a light psycho-educational group intervention with internet for insulin-dependent diabetes patients and problematic hypoglycaemia. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2013, 11, 179-179.	0.0	0
260	Technology and chronic disease management – Authors' reply. <i>Lancet Diabetes and Endocrinology</i> , the, 2018, 6, 91-92.	11.4	0
261	Psychology, Mental Health, and Quality of Life. , 2018, , 941-984.		0
262	Overcoming psychological insulin resistance: A practical guide for healthcare professionals. <i>Primary Care Diabetes</i> , 2021, 15, 619-621.	1.8	0
263	Diabetes Mellitus: A Biopsychosocial Perspective. , 2021, , .		0
264	Behavioral Medicine of the Future: From Disease to Health and Well Being, from the Individual to the Community and the Global, from Causes to Complexity. , 2018, , 1111-1119.		0
265	Improved Diabetes Medication Convenience in Persons with Type 2 Diabetes after Switching to Insulin Glargine 300 U/mL (U-300) – The Observational OPTIN-D Study. <i>Diabetes</i> , 2018, 67, 876-P.	0.6	0
266	The Presence of Cerebral White Matter Lesions and Lower Skin Microvascular Perfusion Predict Decline in General Cognitive Ability in Type 1 Diabetes Patients but Not in Healthy Controls. <i>Diabetes</i> , 2018, 67, .	0.6	0
267	Attitudes among Adults with Type 2 Diabetes Affecting Insulin Initiation and Discontinuation. <i>Diabetes</i> , 2018, 67, 838-P.	0.6	0
268	268-OR: Conversations and Reactions around Severe Hypoglycemia (CRASH): Survey Responses of People Aged 65+ with T1DM or Insulin-Treated T2DM and Caregivers. <i>Diabetes</i> , 2020, 69, .	0.6	0
269	Individual-Level Intervention Approaches. , 2020, , 381-388.		0
270	The multinational conversations and reactions round severe hypoglycemia (CRASH) study: Impact of health care provider communications and recommendations on people with diabetes. <i>Journal of Clinical and Translational Endocrinology</i> , 2022, 27, 100295.	1.4	0