

Frank J Snoek

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2185682/publications.pdf>

Version: 2024-02-01

270
papers

15,081
citations

20036

63
h-index

26792

111
g-index

282
all docs

282
docs citations

282
times ranked

13678
citing authors

#	ARTICLE	IF	CITATIONS
1	The unexplored role of sedentary time and physical activity in glucose and lipid metabolism-related placental mRNAs in pregnant women who are obese: the DALI lifestyle randomised controlled trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2022, 129, 708-721.	1.1	6
2	Interaction between rs10830962 polymorphism in MTNR1B and lifestyle intervention on maternal and neonatal outcomes: secondary analyses of the DALI lifestyle randomized controlled trial. <i>American Journal of Clinical Nutrition</i> , 2022, 115, 388-396.	2.2	5
3	Conversations and Reactions Around Severe Hypoglycemia (CRASH): Japan Results From a Global Survey of People with T1DM or Insulin-Treated T2DM and Caregivers. <i>Diabetes Therapy</i> , 2022, 13, 517-533.	1.2	4
4	Expert Panel Guidance and Narrative Review of Treatment Simplification of Complex Insulin Regimens to Improve Outcomes in Type 2 Diabetes. <i>Diabetes Therapy</i> , 2022, 13, 619-634.	1.2	17
5	How to identify clinically significant diabetes distress using the Problem Areas in Diabetes (PAID) scale in adults with diabetes treated in primary or secondary care? Evidence for new cut points based on latent class analyses. <i>BMJ Open</i> , 2022, 12, e056304.	0.8	13
6	The multinational conversations and reactions round severe hypoglycemia (CRASH) study: Impact of health care provider communications and recommendations on people with diabetes. <i>Journal of Clinical and Translational Endocrinology</i> , 2022, 27, 100295.	1.0	0
7	Symptoms of depression and insomnia in older age: A within-individual analysis over 20 years. <i>Journal of the American Geriatrics Society</i> , 2022, 70, 2051-2059.	1.3	3
8	Conversations and Reactions Around Severe Hypoglycemia (CRASH) Study: Results From People With Diabetes and Caregivers in the United States. <i>Clinical Diabetes</i> , 2022, 40, 477-488.	1.2	4
9	Efficacy and Moderators of Internet-Based Interventions in Adults with Subthreshold Depression: An Individual Participant Data Meta-Analysis of Randomized Controlled Trials. <i>Psychotherapy and Psychosomatics</i> , 2021, 90, 94-106.	4.0	59
10	The importance of maternal insulin resistance throughout pregnancy on neonatal adiposity. <i>Paediatric and Perinatal Epidemiology</i> , 2021, 35, 83-91.	0.8	11
11	Improvement in Patient-Reported Outcomes in Adults with Type 1 Diabetes Treated with Sotagliflozin plus Insulin Versus Insulin Alone. <i>Diabetes Technology and Therapeutics</i> , 2021, 23, 70-77.	2.4	6
12	Less sedentary time is associated with a more favourable glucose-insulin axis in obese pregnant women—a secondary analysis of the DALI study. <i>International Journal of Obesity</i> , 2021, 45, 296-307.	1.6	12
13	Metabolic phenotypes of early gestational diabetes mellitus and their association with adverse pregnancy outcomes. <i>Diabetic Medicine</i> , 2021, 38, e14413.	1.2	23
14	Management of chronic cardiometabolic conditions and mental health during COVID-19. <i>Primary Care Diabetes</i> , 2021, 15, 21-23.	0.9	8
15	Conversations and Reactions Around Severe Hypoglycemia Study: Results of Hypoglycemia Experiences in Canadian Adults With Insulin-Treated Diabetes and Their Caregivers. <i>Canadian Journal of Diabetes</i> , 2021, 45, 236-242.	0.4	17
16	Executive function mediates the link between externalizing behavior and HbA1c in children and adolescents with type 1 diabetes: A cross-national investigation. <i>Pediatric Diabetes</i> , 2021, 22, 503-510.	1.2	1
17	The statistical approach in trial-based economic evaluations matters: get your statistics together!. <i>BMC Health Services Research</i> , 2021, 21, 475.	0.9	19
18	Effects of Diabetes Prevention Education Program for Overweight and Obese Subjects with a Family History of Type 2 Diabetes Mellitus: A Pilot Study from the United Arab Emirates. <i>Oman Medical Journal</i> , 2021, 36, e268-e268.	0.3	2

#	ARTICLE	IF	CITATIONS
19	50-OR: Validation of Fear of Hypoglycemia Screener: Results from the T1D Exchange Registry. <i>Diabetes</i> , 2021, 70, .	0.3	1
20	Data on diabetes-specific distress are needed to improve the quality of diabetes care. <i>Lancet</i> , The, 2021, 397, 2149.	6.3	4
21	Overcoming psychological insulin resistance: A practical guide for healthcare professionals. <i>Primary Care Diabetes</i> , 2021, 15, 619-621.	0.9	0
22	Cognitive Functioning and Hippocampal Connectivity in Patients With Longstanding Type 1 Diabetes and Apolipoprotein E Îµ4. <i>Diabetes Care</i> , 2021, 44, 2388-2396.	4.3	3
23	The Management of Type 1 Diabetes in Adults. A Consensus Report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). <i>Diabetes Care</i> , 2021, 44, 2589-2625.	4.3	244
24	Feasibility and user experience of the unguided web-based self-help app "MyDiaMate"™ aimed to prevent and reduce psychological distress and fatigue in adults with diabetes. <i>Internet Interventions</i> , 2021, 25, 100414.	1.4	6
25	The management of type 1 diabetes in adults. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). <i>Diabetologia</i> , 2021, 64, 2609-2652.	2.9	128
26	Insulin resistance as a marker for the immune-metabolic subtype of depression. <i>Journal of Affective Disorders</i> , 2021, 295, 1371-1376.	2.0	6
27	Conversations and Reactions Around Severe Hypoglycaemia (CRASH): Spanish results of a global survey of people with type 1 diabetes or insulin-treated type 2 diabetes and caregivers. <i>Endocrinología, Diabetes Y Nutrición</i> , 2021, 68, 557-566.	0.1	2
28	Glucose variability and mood in adults with diabetes: A systematic review. <i>Endocrinology, Diabetes and Metabolism</i> , 2021, 4, e00152.	1.0	19
29	Diabetes Mellitus: A Biopsychosocial Perspective. , 2021, , .		0
30	Conversations and Reactions Around Severe Hypoglycaemia (CRASH): Spanish results of a global survey of people with type 1 diabetes or insulin-treated type 2 diabetes and caregivers. <i>Endocrinología y Nutrición (English Ed)</i> , 2021, 68, 557-566.	0.1	3
31	Conversations and Reactions Around Severe Hypoglycemia (CRASH) global survey of people with type-1 diabetes or insulin-treated type-2 diabetes and caregivers: findings from the French cohort. <i>Annales D'Endocrinologie</i> , 2021, 83, 16-16.	0.6	2
32	The DALI vitamin D randomized controlled trial for gestational diabetes mellitus prevention: No major benefit shown besides vitamin D sufficiency. <i>Clinical Nutrition</i> , 2020, 39, 976-984.	2.3	42
33	Terminology matters: "diabulimia"™ is insufficient to describe eating disorders in individuals with Type 1 diabetes. <i>Diabetic Medicine</i> , 2020, 37, 1075-1076.	1.2	11
34	Back to the future: 25 years of "Guidelines for encouraging psychological well-being"™ among people affected by diabetes. <i>Diabetic Medicine</i> , 2020, 37, 1225-1229.	1.2	34
35	Sleep and HbA1c in Patients With Type 2 Diabetes: Which Sleep Characteristics Matter Most?. <i>Diabetes Care</i> , 2020, 43, 235-243.	4.3	51
36	The cognitive and psychological effects of living with type 1 diabetes: a narrative review. <i>Diabetic Medicine</i> , 2020, 37, 555-563.	1.2	47

#	ARTICLE	IF	CITATIONS
37	Performance of early pregnancy HbA1c for predicting gestational diabetes mellitus and adverse pregnancy outcomes in obese European women. <i>Diabetes Research and Clinical Practice</i> , 2020, 168, 108378.	1.1	14
38	Improving interpretability of individual Diabetes Symptom Checklist-Revised (DSC-R) scores: the role of patient characteristics. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001146.	1.2	6
39	Temporal relationships between maternal metabolic parameters with neonatal adiposity in women with obesity differ by neonatal sex: Secondary analysis of the DALI study. <i>Pediatric Obesity</i> , 2020, 15, e12628.	1.4	11
40	Key Strategies for Overcoming Psychological Insulin Resistance in Adults with Type 2 Diabetes: The UK Subgroup in the EMOTION Study. <i>Diabetes Therapy</i> , 2020, 11, 1735-1744.	1.2	3
41	Comparing the EQ-5D crosswalks and value sets for England, the Netherlands and Spain: Exploring their impact on cost-utility results. <i>Health Economics (United Kingdom)</i> , 2020, 29, 640-651.	0.8	7
42	Clinical assessment of emotions in patients with cancer: Diagnostic accuracy compared with two reference standards. <i>Psycho-Oncology</i> , 2020, 29, 775-780.	1.0	4
43	#DiabetesPsychologyMatters. <i>Diabetes Spectrum</i> , 2020, 33, 95-98.	0.4	2
44	268-OR: Conversations and Reactions around Severe Hypoglycemia (CRASH): Survey Responses of People Aged 65+ with T1DM or Insulin-Treated T2DM and Caregivers. <i>Diabetes</i> , 2020, 69, .	0.3	0
45	Conversations and Reactions Around Severe Hypoglycaemia (CRASH): Results from the German Cohort of a Global Survey of People with Type 1 Diabetes or Insulin-Treated Type 2 Diabetes and Caregivers. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2020, , .	0.6	3
46	Individual-Level Intervention Approaches. , 2020, , 381-388.		0
47	Diabetes distress is associated with adverse pregnancy outcomes in women with gestational diabetes: a prospective cohort study. <i>BMC Pregnancy and Childbirth</i> , 2019, 19, 223.	0.9	31
48	Identifying solutions to psychological insulin resistance: An international study. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 307-314.	1.2	23
49	The presence of cerebral white matter lesions and lower skin microvascular perfusion predicts lower cognitive performance in type 1 diabetes patients with retinopathy but not in healthy controls: A longitudinal study. <i>Microcirculation</i> , 2019, 26, e12530.	1.0	8
50	Decision aids that facilitate elements of shared decision making in chronic illnesses: a systematic review. <i>Systematic Reviews</i> , 2019, 8, 121.	2.5	60
51	Do Nonsuicidal Severely Depressed Individuals with Diabetes Profit from Internet-Based Guided Self-Help? Secondary Analyses of a Pragmatic Randomized Trial. <i>Journal of Diabetes Research</i> , 2019, 2019, 1-11.	1.0	14
52	Nutritional Lifestyle Intervention in Obese Pregnant Women, Including Lower Carbohydrate Intake, Is Associated With Increased Maternal Free Fatty Acids, 3-Î²-Hydroxybutyrate, and Fasting Glucose Concentrations: A Secondary Factorial Analysis of the European Multicenter, Randomized Controlled DALI Lifestyle Intervention Trial. <i>Diabetes Care</i> , 2019, 42, 1380-1389.	4.3	21
53	Effectiveness of psychoeducational interventions for the treatment of diabetes-specific emotional distress and glycaemic control in people with type 2 diabetes: A systematic review and meta-analysis. <i>Primary Care Diabetes</i> , 2019, 13, 556-567.	0.9	24
54	A reduction in sedentary behaviour in obese women during pregnancy reduces neonatal adiposity: the DALI randomised controlled trial. <i>Diabetologia</i> , 2019, 62, 915-925.	2.9	50

#	ARTICLE	IF	CITATIONS
55	Effects of Light Therapy on Mood and Insulin Sensitivity in Patients With Type 2 Diabetes and Depression: Results From a Randomized Placebo-Controlled Trial. <i>Diabetes Care</i> , 2019, 42, 529-538.	4.3	26
56	Mediators of Lifestyle Behaviour Changes in Obese Pregnant Women. Secondary Analyses from the DALI Lifestyle Randomised Controlled Trial. <i>Nutrients</i> , 2019, 11, 311.	1.7	6
57	Comment on the consensus report on the management of hyperglycaemia in Type 2 diabetes by the American Diabetes Association and the European Association for the Study of Diabetes. <i>Diabetic Medicine</i> , 2019, 36, 911-912.	1.2	1
58	Key factors for overcoming psychological insulin resistance: an examination of patient perspectives through content analysis. <i>BMJ Open Diabetes Research and Care</i> , 2019, 7, e000723.	1.2	12
59	115 - Conversations and Reactions Around Severe Hypoglycemia (CRASH Study): Canadian Results. <i>Canadian Journal of Diabetes</i> , 2019, 43, S40.	0.4	1
60	Looking back on 25 years of the PSAD study group. <i>Diabetic Medicine</i> , 2019, 37, 380-382.	1.2	3
61	The Effects of Lifestyle and/or Vitamin D Supplementation Interventions on Pregnancy Outcomes: What Have We Learned from the DALI Studies?. <i>Current Diabetes Reports</i> , 2019, 19, 162.	1.7	8
62	How has psycho-behavioural research advanced our understanding of hypoglycaemia in type 1 diabetes?. <i>Diabetic Medicine</i> , 2019, 37, 409-417.	1.2	15
63	Youth With Type 1 Diabetes Taking Responsibility for Self-Management: The Importance of Executive Functioning in Achieving Glycemic Control. <i>Diabetes Care</i> , 2019, 42, 225-231.	4.3	36
64	Cognition in elderly with type 1 diabetes: Is there an interaction between glycemia and aging?. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 4-5.	1.2	1
65	285-OR: Conversations and Reactions around Severe Hypoglycemia (CRASH): U.S. Results from a Global Survey of People with T1DM or Insulin-Treated T2DM and Caregivers. <i>Diabetes</i> , 2019, 68, .	0.3	3
66	Accelerated executive functions decline and gray matter structural changes in middle-aged type 1 diabetes mellitus patients with proliferative retinopathy. <i>Journal of Diabetes</i> , 2018, 10, 835-846.	0.8	9
67	Cost-effectiveness of healthy eating and/or physical activity promotion in pregnant women at increased risk of gestational diabetes mellitus: economic evaluation alongside the DALI study, a European multicenter randomized controlled trial. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2018, 15, 23.	2.0	34
68	Web-based intervention for depressive symptoms in adults with types 1 and 2 diabetes mellitus: a health economic evaluation. <i>British Journal of Psychiatry</i> , 2018, 212, 199-206.	1.7	41
69	Comment on Young-Hyman et al. Psychosocial Care for People With Diabetes: A Position Statement of the American Diabetes Association. <i>Diabetes Care</i> 2016;39:2126-2140. <i>Diabetes Care</i> , 2018, 41, e31-e32.	4.3	6
70	Diabetes structured self-management education programmes: a narrative review and current innovations. <i>Lancet Diabetes and Endocrinology</i> , 2018, 6, 130-142.	5.5	233
71	Cost-effectiveness of the psycho-educational blended (group and online) intervention HypoAware compared with usual care for people with Type 1 and insulin-treated Type 2 diabetes with problematic hypoglycaemia: analyses of a cluster-randomized controlled trial. <i>Diabetic Medicine</i> , 2018, 35, 214-222.	1.2	6
72	Technology and chronic disease management – Authors' reply. <i>Lancet Diabetes and Endocrinology</i> , 2018, 6, 91-92.	5.5	0

#	ARTICLE	IF	CITATIONS
73	PRM202 - COMPARING THE EQ-5D-5L CROSSWALKS AND VALUE SETS FOR ENGLAND, THE NETHERLANDS AND SPAIN: DO CONCLUSIONS CHANGE?. Value in Health, 2018, 21, S391.	0.1	1
74	Association between Gestational Weight Gain, Gestational Diabetes Risk, and Obstetric Outcomes: A Randomized Controlled Trial Post Hoc Analysis. Nutrients, 2018, 10, 1568.	1.7	22
75	Psychology, Mental Health, and Quality of Life. , 2018, , 941-984.		0
76	Improved diabetes medication convenience and satisfaction in persons with type 2 diabetes after switching to insulin glargine 300 U/mL: results of the observational OPTIN-D study. BMJ Open Diabetes Research and Care, 2018, 6, e000548.	1.2	11
77	High Diabetes Distress Among Ethnic Minorities Is Not Explained by Metabolic, Cardiovascular, or Lifestyle Factors: Findings From the Dutch Diabetes Pearl Cohort. Diabetes Care, 2018, 41, 1854-1861.	4.3	23
78	Does hypoglycaemia affect the improvement in QoL after the transition to insulin in people with type 2 diabetes?. Journal of Endocrinological Investigation, 2018, 41, 249-258.	1.8	10
79	Risk factors for hyperglycemia in pregnancy in the DALI study differ by period of pregnancy and OGTT time point. European Journal of Endocrinology, 2018, 179, 39-49.	1.9	20
80	The challenge of treating comorbid mental health problems in patients with a somatic illness. Lancet Psychiatry, the, 2018, 5, 465.	3.7	6
81	Re: Vitamin D and gestational diabetes mellitus: a systematic review based on data free of Hawthorne effect. BJOG: an International Journal of Obstetrics and Gynaecology, 2018, 125, 1338-1339.	1.1	5
82	Systematic review and meta-analysis of psychological interventions in people with diabetes and elevated diabetes distress. Diabetic Medicine, 2018, 35, 1157-1172.	1.2	103
83	Do guided internet-based interventions result in clinically relevant changes for patients with depression? An individual participant data meta-analysis. Clinical Psychology Review, 2018, 63, 80-92.	6.0	239
84	Key Factors for Overcoming Psychological Insulin Resistance—An Examination of a Large International Sample through Content Analysis. Diabetes, 2018, 67, .	0.3	3
85	Internet-Based Interventions in Chronic Somatic Disease. Deutsches Ärztblatt International, 2018, 115, 659-665.	0.6	47
86	Behavioral Medicine of the Future: From Disease to Health and Well Being, from the Individual to the Community and the Global, from Causes to Complexity. , 2018, , 1111-1119.		0
87	Improved Diabetes Medication Convenience in Persons with Type 2 Diabetes after Switching to Insulin Glargine 300 U/mL (U-300)—The Observational OPTIN-D Study. Diabetes, 2018, 67, 876-P.	0.3	0
88	The Presence of Cerebral White Matter Lesions and Lower Skin Microvascular Perfusion Predict Decline in General Cognitive Ability in Type 1 Diabetes Patients but Not in Healthy Controls. Diabetes, 2018, 67, .	0.3	0
89	Attitudes among Adults with Type 2 Diabetes Affecting Insulin Initiation and Discontinuation. Diabetes, 2018, 67, 838-P.	0.3	0
90	Effect of physical activity and/or healthy eating on GDM risk: The DALI Lifestyle Study. Journal of Clinical Endocrinology and Metabolism, 2017, 102, jc.2016-3455.	1.8	140

#	ARTICLE	IF	CITATIONS
91	Altered eigenvector centrality is related to local resting-state network functional connectivity in patients with longstanding type 1 diabetes mellitus. <i>Human Brain Mapping</i> , 2017, 38, 3623-3636.	1.9	33
92	Validation of a quick screening instrument for measuring fear of hypoglycaemia in persons with diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1360-1361.	1.2	4
93	Role of Continuous Glucose Monitoring in Clinical Trials: Recommendations on Reporting. <i>Diabetes Technology and Therapeutics</i> , 2017, 19, 391-399.	2.4	45
94	Continuous Glucose Monitoring in Patients with Type 1 Diabetes and Impaired Awareness of Hypoglycemia: Also Effective in Patients with Psychological Distress?. <i>Diabetes Technology and Therapeutics</i> , 2017, 19, 595-599.	2.4	13
95	Light therapy: is it safe for the eyes?. <i>Acta Psychiatrica Scandinavica</i> , 2017, 136, 534-548.	2.2	29
96	Epidemiology of gestational diabetes mellitus according to IADPSG/WHO 2013 criteria among obese pregnant women in Europe. <i>Diabetologia</i> , 2017, 60, 1913-1921.	2.9	117
97	The prevalence of diabetes-specific emotional distress in people with Type 2 diabetes: a systematic review and meta-analysis. <i>Diabetic Medicine</i> , 2017, 34, 1508-1520.	1.2	285
98	Screening for sexual dissatisfaction among people with type 2 diabetes in primary care. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1614-1619.	1.2	9
99	Disturbed eating behaviors in adolescents with type 1 diabetes. How to screen for yellow flags in clinical practice?. <i>Pediatric Diabetes</i> , 2017, 18, 376-383.	1.2	19
100	Keeping safe. Continuous glucose monitoring (CGM) in persons with Type 1 diabetes and impaired awareness of hypoglycaemia: a qualitative study. <i>Diabetic Medicine</i> , 2017, 34, 1470-1476.	1.2	24
101	Ethnic Minorities with Diabetes Differ in Depressive and Anxiety Symptoms and Diabetes-Distress. <i>Journal of Diabetes Research</i> , 2017, 2017, 1-11.	1.0	15
102	Is a motivational interviewing based lifestyle intervention for obese pregnant women across Europe implemented as planned? Process evaluation of the DALI study. <i>BMC Pregnancy and Childbirth</i> , 2017, 17, 293.	0.9	6
103	Correlates of poor mental health in early pregnancy in obese European women. <i>BMC Pregnancy and Childbirth</i> , 2017, 17, 404.	0.9	11
104	A systematic review of decision aids that facilitate elements of shared decision-making in chronic illnesses: a review protocol. <i>Systematic Reviews</i> , 2017, 6, 155.	2.5	17
105	Parental Diabetes Behaviors and Distress Are Related to Glycemic Control in Youth with Type 1 Diabetes: Longitudinal Data from the DINO Study. <i>Journal of Diabetes Research</i> , 2017, 2017, 1-7.	1.0	28
106	Clinical assessment of emotions in patients with cancer: Diagnostic accuracy compared to two reference standards.. <i>Journal of Clinical Oncology</i> , 2017, 35, 24-24.	0.8	12
107	Beliefs, Barriers, and Preferences of European Overweight Women to Adopt a Healthier Lifestyle in Pregnancy to Minimize Risk of Developing Gestational Diabetes Mellitus: An Explorative Study. <i>Journal of Pregnancy</i> , 2016, 2016, 1-11.	1.1	31
108	Low Self-Confidence and Diabetes Mismanagement in Youth with Type 1 Diabetes Mediate the Relationship between Behavioral Problems and Elevated HbA1c. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-6.	1.0	9

#	ARTICLE	IF	CITATIONS
109	Uptake and Effects of the e-Vita Personal Health Record with Self-Management Support and Coaching, for Type 2 Diabetes Patients Treated in Primary Care. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-9.	1.0	29
110	Subgenual Cingulate Cortex Functional Connectivity in Relation to Depressive Symptoms and Cognitive Functioning in Type 1 Diabetes Mellitus Patients. <i>Psychosomatic Medicine</i> , 2016, 78, 740-749.	1.3	16
111	Effects of self-monitoring of glucose on distress and self-efficacy in people with non-insulin-treated Type 2 diabetes: a randomized controlled trial. <i>Diabetic Medicine</i> , 2016, 33, 537-546.	1.2	7
112	HypoAware: development and pilot study of a brief and partly web-based psychoeducational group intervention for adults with Type 1 and insulin-treated Type 2 diabetes and problematic hypoglycaemia. <i>Diabetic Medicine</i> , 2016, 33, 184-191.	1.2	19
113	Does Internet-based guided-self-help for depression cause harm? An individual participant data meta-analysis on deterioration rates and its moderators in randomized controlled trials. <i>Psychological Medicine</i> , 2016, 46, 2679-2693.	2.7	129
114	IADPSG and WHO 2013 Gestational Diabetes Mellitus Criteria Identify Obese Women With Marked Insulin Resistance in Early Pregnancy. <i>Diabetes Care</i> , 2016, 39, e90-e92.	4.3	79
115	Continuous glucose monitoring for patients with type 1 diabetes and impaired awareness of hypoglycaemia (IN CONTROL): a randomised, open-label, crossover trial. <i>Lancet Diabetes and Endocrinology</i> , 2016, 4, 893-902.	5.5	284
116	Effectiveness of HypoAware, a Brief Partly Web-Based Psychoeducational Intervention for Adults With Type 1 and Insulin-Treated Type 2 Diabetes and Problematic Hypoglycemia: A Cluster Randomized Controlled Trial. <i>Diabetes Care</i> , 2016, 39, 2190-2196.	4.3	35
117	Disrupted subject-specific gray matter network properties and cognitive dysfunction in type 1 diabetes patients with and without proliferative retinopathy. <i>Human Brain Mapping</i> , 2016, 37, 1194-1208.	1.9	25
118	Implementation of quality of life monitoring in Dutch routine care of adolescents with type 1 diabetes: appreciated but difficult. <i>Pediatric Diabetes</i> , 2016, 17, 112-119.	1.2	21
119	Does low well-being modify the effects of PRISMA (Dutch DESMOND), a structured self-management-education program for people with type 2 diabetes?. <i>Primary Care Diabetes</i> , 2016, 10, 103-110.	0.9	10
120	Type 2 Diabetes Patients' Needs and Preferences for Care Concerning Sexual Problems: A Cross-Sectional Survey and Qualitative Interviews. <i>Journal of Sex and Marital Therapy</i> , 2016, 42, 324-337.	1.0	19
121	Constructs of depression and distress in diabetes: time for an appraisal. <i>Lancet Diabetes and Endocrinology</i> , 2015, 3, 450-460.	5.5	304
122	Efficacy of a Web-Based Intervention With Mobile Phone Support in Treating Depressive Symptoms in Adults With Type 1 and Type 2 Diabetes: A Randomized Controlled Trial. <i>Diabetes Care</i> , 2015, 38, 776-783.	4.3	143
123	Results From a European Multicenter Randomized Trial of Physical Activity and/or Healthy Eating to Reduce the Risk of Gestational Diabetes Mellitus: The DALI Lifestyle Pilot. <i>Diabetes Care</i> , 2015, 38, 1650-1656.	4.3	93
124	Diabetes risk reduction in overweight first degree relatives of type 2 diabetes patients: Effects of a low-intensive lifestyle education program (DiAlert) A randomized controlled trial. <i>Patient Education and Counseling</i> , 2015, 98, 476-483.	1.0	16
125	Effectiveness of a PLISSIT model intervention in patients with type 2 diabetes mellitus in primary care: design of a cluster-randomised controlled trial. <i>BMC Family Practice</i> , 2015, 16, 69.	2.9	19
126	Light therapy for better mood and insulin sensitivity in patients with major depression and type 2 diabetes: a randomised, double-blind, parallel-arm trial. <i>BMC Psychiatry</i> , 2015, 15, 169.	1.1	17

#	ARTICLE	IF	CITATIONS
127	Design and rationale of the IN CONTROL trial: the effects of real-time continuous glucose monitoring on glycemia and quality of life in patients with type 1 diabetes mellitus and impaired awareness of hypoglycemia. <i>BMC Endocrine Disorders</i> , 2015, 15, 42.	0.9	11
128	Diabetes IN develOpment (DINO): the bio-psychosocial, family functioning and parental well-being of youth with type 1 diabetes: a longitudinal cohort study design. <i>BMC Pediatrics</i> , 2015, 15, 82.	0.7	14
129	HypoAware-a brief and partly web-based psycho-educational group intervention for adults with type 1 and insulin-treated type 2 diabetes and problematic hypoglycaemia: design of a cost-effectiveness randomised controlled trial. <i>BMC Endocrine Disorders</i> , 2015, 15, 43.	0.9	13
130	Physical activity, depressed mood and pregnancy worries in European obese pregnant women: results from the DALI study. <i>BMC Pregnancy and Childbirth</i> , 2015, 15, 158.	0.9	36
131	Clinical Utility of SMBG: Recommendations on the Use and Reporting of SMBG in Clinical Research. <i>Diabetes Care</i> , 2015, 38, 1627-1633.	4.3	28
132	Prevalence and Correlates of Sexual Dysfunction in Men and Women With Type 2 Diabetes. <i>Journal of Sex and Marital Therapy</i> , 2015, 41, 680-690.	1.0	22
133	Diabetes-specific emotional distress in people with Type 2 diabetes: a comparison between primary and secondary care. <i>Diabetic Medicine</i> , 2014, 31, 1252-1259.	1.2	56
134	Toward Defining a Cutoff Score for Elevated Fear of Hypoglycemia on the Hypoglycemia Fear Survey Worry Subscale in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2014, 37, 102-108.	4.3	46
135	Differential impact of subclinical carotid artery disease on cerebral structure and functioning in type 1 diabetes patients with versus those without proliferative retinopathy. <i>Cardiovascular Diabetology</i> , 2014, 13, 58.	2.7	6
136	Illness representations of type 2 diabetes patients are associated with perceptions of diabetes threat in relatives. <i>Journal of Health Psychology</i> , 2014, 19, 358-368.	1.3	12
137	Symptom burden and its association with change in glucose metabolism status over a 7-year period: the Hoorn Study. <i>Diabetic Medicine</i> , 2014, 31, 747-753.	1.2	1
138	Cerebrospinal fluid levels of Alzheimer's disease biomarkers in middle-aged patients with type 1 diabetes. <i>Diabetologia</i> , 2014, 57, 2208-2214.	2.9	40
139	Ventral Striatum, but Not Cortical Volume Loss, Is Related to Cognitive Dysfunction in Type 1 Diabetic Patients With and Without Microangiopathy. <i>Diabetes Care</i> , 2014, 37, 2483-2490.	4.3	31
140	Changes in MEG resting-state networks are related to cognitive decline in type 1 diabetes mellitus patients. <i>NeuroImage: Clinical</i> , 2014, 5, 69-76.	1.4	19
141	Psychometric evaluation of the Korean version of the Diabetes Symptom Checklist-Revised (DSC-R) for Patients with Type 2 Diabetes. <i>Health and Quality of Life Outcomes</i> , 2014, 12, 77.	1.0	6
142	Proliferative Retinopathy in Type 1 Diabetes Is Associated With Cerebral Microbleeds, Which Is Part of Generalized Microangiopathy. <i>Diabetes Care</i> , 2014, 37, 1165-1168.	4.3	61
143	Health care providers' perspective on using family history in the prevention of type 2 diabetes: a qualitative study including different disciplines. <i>BMC Family Practice</i> , 2013, 14, 31.	2.9	15
144	DALI: Vitamin D and lifestyle intervention for gestational diabetes mellitus (GDM) prevention: an European multicentre, randomised trial "study protocol. <i>BMC Pregnancy and Childbirth</i> , 2013, 13, 142.	0.9	85

#	ARTICLE	IF	CITATIONS
145	Efficacy and cost-effectiveness of a web-based intervention with mobile phone support to treat depressive symptoms in adults with diabetes mellitus type 1 and type 2: design of a randomised controlled trial. <i>BMC Psychiatry</i> , 2013, 13, 306.	1.1	52
146	Web-based self-management with and without coaching for type 2 diabetes patients in primary care: design of a randomized controlled trial. <i>BMC Endocrine Disorders</i> , 2013, 13, 53.	0.9	13
147	Screening, evaluation and management of depression in people with diabetes in primary care. <i>Primary Care Diabetes</i> , 2013, 7, 1-10.	0.9	91
148	Ethnic aspects of emotional distress in patients with diabetes – the Amsterdam Health Monitor Study. <i>Diabetic Medicine</i> , 2013, 30, e25-31.	1.2	14
149	Psychometric and screening properties of the WHO’s well-being index in adult outpatients with Type 1 or Type 2 diabetes mellitus. <i>Diabetic Medicine</i> , 2013, 30, e63-9.	1.2	158
150	PS12 - 3. Use of behavioral change techniques in web-based self-management programs for type 2 diabetes patients: a systematic review. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2013, 11, 167-168.	0.0	0
151	PS12 - 5. Development and evaluation of a light psycho-educational group intervention with internet for insulin-dependent diabetes patients and problematic hypoglycaemia. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2013, 11, 179-179.	0.0	0
152	Differential associations between depressive symptoms and glycaemic control in outpatients with diabetes. <i>Diabetic Medicine</i> , 2013, 30, e115-22.	1.2	77
153	Examining the Behaviour subscale of the Hypoglycaemia Fear Survey: an international study. <i>Diabetic Medicine</i> , 2013, 30, 603-609.	1.2	57
154	Comment on “Psychological distress in patients with cancer: is screening the effective solution?”. <i>British Journal of Cancer</i> , 2013, 108, 2628-2630.	2.9	15
155	Use of Behavioral Change Techniques in Web-Based Self-Management Programs for Type 2 Diabetes Patients: Systematic Review. <i>Journal of Medical Internet Research</i> , 2013, 15, e279.	2.1	100
156	Monitoring of Individual Needs in Diabetes (MIND)-2. <i>Diabetes Care</i> , 2012, 35, 2128-2132.	4.3	92
157	PS11 - 50. Depressive symptoms exacerbate the negative effects of type 1 diabetes and microangiopathy on brain functioning. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2012, 10, 133-133.	0.0	0
158	PS11 - 51. Prevalence and determinants of diabetes-specific emotional distress: primary care and secondary care patients with type 2 diabetes mellitus compared. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2012, 10, 133-134.	0.0	0
159	Resting-State Brain Networks in Type 1 Diabetic Patients With and Without Microangiopathy and Their Relation to Cognitive Functions and Disease Variables. <i>Diabetes</i> , 2012, 61, 1814-1821.	0.3	109
160	The longitudinal association between glycaemic control and health-related quality of life following insulin therapy optimisation in type 2 diabetes patients. A prospective observational study in secondary care. <i>Quality of Life Research</i> , 2012, 21, 1359-1365.	1.5	27
161	DiAlert: a prevention program for overweight first degree relatives of type 2 diabetes patients: results of a pilot study to test feasibility and acceptability. <i>Trials</i> , 2012, 13, 178.	0.7	4
162	Patients’ intentions to inform relatives about Type 2 diabetes risk: the role of worry in the process of family risk disclosure. <i>Diabetic Medicine</i> , 2012, 29, e461-7.	1.2	8

#	ARTICLE	IF	CITATIONS
163	Family communication as strategy in diabetes prevention: An observational study in families with Dutch and Surinamese South-Asian ancestry. <i>Patient Education and Counseling</i> , 2012, 87, 23-29.	1.0	15
164	Assessing diabetes-related quality of life of youth with type 1 diabetes in routine clinical care: the MIND Youth Questionnaire (MY-Q). <i>Pediatric Diabetes</i> , 2012, 13, 638-646.	1.2	64
165	Diffusion tensor imaging in type 1 diabetes: decreased white matter integrity relates to cognitive functions. <i>Diabetologia</i> , 2012, 55, 1218-1220.	2.9	58
166	Is a Severe Clinical Profile an Effect Modifier in a Web-Based Depression Treatment for Adults With Type 1 or Type 2 Diabetes? Secondary Analyses From a Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2012, 14, e2.	2.1	29
167	Are patients with type 2 diabetes reluctant to start insulin therapy? An examination of the scope and underpinnings of psychological insulin resistance in a large, international population. <i>Current Medical Research and Opinion</i> , 2011, 27, 1169-1174.	0.9	96
168	Changing the odds. What do we learn from prevention studies targeted at people with a positive family history of type 2 diabetes?. <i>Primary Care Diabetes</i> , 2011, 5, 215-221.	0.9	23
169	Depressive symptoms and unmet psychological needs of Dutch youth with type 1 diabetes: results of a web-survey. <i>Pediatric Diabetes</i> , 2011, 12, 172-176.	1.2	44
170	Initiation of insulin glargine in patients with Type 2 diabetes in suboptimal glycaemic control positively impacts health-related quality of life. A prospective cohort study in primary care. <i>Diabetic Medicine</i> , 2011, 28, 1096-1102.	1.2	31
171	Do physicians understand Type 2 diabetes patients' perceptions of seriousness; the emotional impact and needs for care improvement? A cross-national survey. <i>Patient Education and Counseling</i> , 2011, 85, 258-263.	1.0	29
172	DiAlert: a lifestyle education programme aimed at people with a positive family history of type 2 diabetes and overweight, study protocol of a randomised controlled trial. <i>BMC Public Health</i> , 2011, 11, 751.	1.2	18
173	Limited effect of screening for depression with written feedback in outpatients with diabetes mellitus: a randomised controlled trial. <i>Diabetologia</i> , 2011, 54, 741-748.	2.9	54
174	Development and reach of a web-based cognitive behavioural therapy programme to reduce symptoms of depression and diabetes-specific distress. <i>Patient Education and Counseling</i> , 2011, 84, 49-55.	1.0	39
175	Web-Based Depression Treatment for Type 1 and Type 2 Diabetic Patients. <i>Diabetes Care</i> , 2011, 34, 320-325.	4.3	184
176	PS1 - 6. Cerebral resting-state network changes in patients with type 1 diabetes with and without microangiopathy relate to cognitive functions. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2011, 9, 94-94.	0.0	0
177	PS11 - 56. Implementation of quality of life in routine care of adolescents with diabetes mellitus type 1. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2011, 9, 129-129.	0.0	0
178	Monitoring of Individual Needs in Diabetes (MIND): Baseline Data From the Cross-National Diabetes Attitudes, Wishes, and Needs (DAWN) MIND Study. <i>Diabetes Care</i> , 2011, 34, 601-603.	4.3	103
179	Follow-up results on monitoring and discussing health-related quality of life in adolescent diabetes care: benefits do not sustain in routine practice. <i>Pediatric Diabetes</i> , 2010, 11, 175-181.	1.2	56
180	Short-form measures of diabetes-related emotional distress: the Problem Areas in Diabetes Scale (PAID)-5 and PAID-1. <i>Diabetologia</i> , 2010, 53, 66-69.	2.9	290

#	ARTICLE	IF	CITATIONS
181	"I am pregnant and my husband has diabetes. Is there a risk for my child?" A qualitative study of questions asked by email about the role of genetic susceptibility to diabetes. BMC Public Health, 2010, 10, 688.	1.2	8
182	Effects of the PRO-active Interdisciplinary Self-MANagement (PRISMA, Dutch DESMOND) program on dietary intake in type 2 diabetes outpatients: A pilot study. Clinical Nutrition, 2010, 29, 199-205.	2.3	17
183	Effect of interventions for major depressive disorder and significant depressive symptoms in patients with diabetes mellitus: a systematic review and meta-analysis. General Hospital Psychiatry, 2010, 32, 380-395.	1.2	290
184	Eicosapentaenoic acid as an add-on to antidepressant medication for co-morbid major depression in patients with diabetes mellitus: A randomized, double-blind placebo-controlled study. Journal of Affective Disorders, 2010, 126, 282-286.	2.0	49
185	Prevalence of comorbid depression is high in outpatients with Type 1 or Type 2 diabetes mellitus. Results from three outpatient clinics in the Netherlands. Diabetic Medicine, 2010, 27, 217-224.	1.2	131
186	Diabetes-specific emotional distress mediates the association between depressive symptoms and glycaemic control in Type 1 and Type 2 diabetes. Diabetic Medicine, 2010, 27, 798-803.	1.2	136
187	Depression, Anxiety and Glucose Metabolism in the General Dutch Population: The New Hoorn Study. PLoS ONE, 2010, 5, e9971.	1.1	88
188	Assessment of Parent-Adolescent Partnership in Diabetes Care. The Diabetes Educator, 2010, 36, 205-215.	2.6	14
189	A 24-Week, Randomized, Treat-to-Target Trial Comparing Initiation of Insulin Glargine Once-Daily With Insulin Detemir Twice-Daily in Patients With Type 2 Diabetes Inadequately Controlled on Oral Glucose-Lowering Drugs. Diabetes Care, 2010, 33, 1176-1178.	4.3	145
190	Functional Brain Connectivity and Neurocognitive Functioning in Patients With Long-Standing Type 1 Diabetes With and Without Microvascular Complications. Diabetes, 2009, 58, 2335-2343.	0.3	67
191	Effects of self-monitoring of glucose in non-insulin treated patients with type 2 diabetes: design of the IN CONTROL-trial. BMC Family Practice, 2009, 10, 26.	2.9	13
192	Web-based guided insulin self-titration in patients with type 2 diabetes: the Di@log study. Design of a cluster randomised controlled trial [TC1316]. BMC Family Practice, 2009, 10, 40.	2.9	11
193	Barriers to insulin initiation and intensification and how to overcome them. International Journal of Clinical Practice, 2009, 63, 6-10.	0.8	134
194	The DAWN MIND Youth program. Pediatric Diabetes, 2009, 10, 46-49.	1.2	6
195	Symptoms of depression and diabetes-specific emotional distress are associated with a negative appraisal of insulin therapy in insulin-naïve patients with Type 2 diabetes mellitus. A study from the European Depression in Diabetes [EDID] Research Consortium. Diabetic Medicine, 2009, 26, 28-33.	1.2	71
196	Psychometric Evaluation of the Diabetes Symptom Checklist-Revised (DSC-R) – A Measure of Symptom Distress. Value in Health, 2009, 12, 1168-1175.	0.1	65
197	Rationale, Design, and Baseline Data of the Insulin Glargine (Lantus) Versus Insulin Detemir (Levemir) Treat-To-Target (L2T3) Study: A Multinational, Randomized Noninferiority Trial of Basal Insulin Initiation in Type 2 Diabetes. Diabetes Technology and Therapeutics, 2009, 11, 739-743.	2.4	17
198	Evaluatie en bespreking van de kwaliteit van leven van adolescenten met type-1-diabetes. Tijdschrift Voor Kindergeneeskunde, 2008, 76, 180-189.	0.0	1

#	ARTICLE	IF	CITATIONS
199	Web-based cognitive behavioural therapy (W-CBT) for diabetes patients with co-morbid depression: Design of a randomised controlled trial. <i>BMC Psychiatry</i> , 2008, 8, 9.	1.1	43
200	Perceptions of risk in adults with a low or high risk profile of developing type 2 diabetes; a cross-sectional population-based study. <i>Patient Education and Counseling</i> , 2008, 73, 307-312.	1.0	23
201	Symptoms of depression in people with impaired glucose metabolism or Type 2 diabetes mellitus: The Hoorn Study. <i>Diabetic Medicine</i> , 2008, 25, 843-849.	1.2	49
202	Cognitive behavioural therapy (CBT) compared with blood glucose awareness training (BGAT) in poorly controlled Type 1 diabetic patients: long-term effects on HbA _{1c} moderated by depression. A randomized controlled trial. <i>Diabetic Medicine</i> , 2008, 25, 1337-1342.	1.2	68
203	Routine Psychological Screening in Youth With Type 1 Diabetes and Their Parents: a Notion Whose Time Has Come?: Response to Cameron et al.. <i>Diabetes Care</i> , 2008, 31, e14-e14.	4.3	1
204	Diabetes-Related Symptom Distress in Association With Glucose Metabolism and Comorbidity: The Hoorn Study. <i>Diabetes Care</i> , 2008, 31, 2268-2270.	4.3	47
205	Monitoring and Discussing Health-Related Quality of Life in Adolescents With Type 1 Diabetes Improve Psychosocial Well-Being. <i>Diabetes Care</i> , 2008, 31, 1521-1526.	4.3	207
206	Self management of type 2 diabetes. <i>BMJ: British Medical Journal</i> , 2007, 335, 458-459.	2.4	7
207	Validation of the WHO-5 Well-Being Index in Adolescents With Type 1 Diabetes. <i>Diabetes Care</i> , 2007, 30, 2003-2006.	4.3	241
208	Monitoring health related quality of life in adolescents with diabetes: a review of measures. <i>Archives of Disease in Childhood</i> , 2007, 92, 434-439.	1.0	69
209	Self-report and parent-report of physical and psychosocial well-being in Dutch adolescents with type 1 diabetes in relation to glycemic control. <i>Health and Quality of Life Outcomes</i> , 2007, 5, 10.	1.0	59
210	Development and validation of the insulin treatment appraisal scale (ITAS) in patients with type 2 diabetes. <i>Health and Quality of Life Outcomes</i> , 2007, 5, 69.	1.0	107
211	No evidence for increased self-reported cognitive failure in Type 1 and Type 2 diabetes: a cross-sectional study. <i>Diabetic Medicine</i> , 2007, 24, 735-740.	1.2	12
212	Cognitive performance in type 1 diabetes patients is associated with cerebral white matter volume. <i>Diabetologia</i> , 2007, 50, 1763-1769.	2.9	105
213	Diabetes and psychological well-being: crossing borders to achieve optimum care. <i>Practical Diabetes International: the International Journal for Diabetes Care Teams Worldwide</i> , 2006, 23, 241-242.	0.2	6
214	Type 2 diabetes and inheritance: what information do diabetes organizations provide on the Internet?. <i>Diabetic Medicine</i> , 2006, 23, 1233-1238.	1.2	14
215	Nurses' recognition and registration of depression, anxiety and diabetes-specific emotional problems in outpatients with diabetes mellitus. <i>Patient Education and Counseling</i> , 2006, 60, 235-240.	1.0	114
216	Patient and provider perceptions of care for diabetes: results of the cross-national DAWN Study. <i>Diabetologia</i> , 2006, 49, 279-288.	2.9	89

#	ARTICLE	IF	CITATIONS
217	Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. <i>Diabetologia</i> , 2006, 49, 837-845.	2.9	815
218	Diabetic encephalopathy: a concept in need of a definition. <i>Diabetologia</i> , 2006, 49, 1447-1448.	2.9	176
219	Voxel-based morphometry demonstrates reduced grey matter density on brain MRI in patients with diabetic retinopathy. <i>Diabetologia</i> , 2006, 49, 2474-2480.	2.9	156
220	Associations between depressive symptoms and insulin resistance: The Hoorn Study. <i>Diabetologia</i> , 2006, 49, 2874-2877.	2.9	53
221	The psychological impact of screening for type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2006, 22, 20-25.	1.7	48
222	Microvascular Disease in Type 1 Diabetes Alters Brain Activation: A Functional Magnetic Resonance Imaging Study. <i>Diabetes</i> , 2006, 55, 334-340.	0.3	63
223	Cognitive behavioural group training (CBGT) for patients with type 1 diabetes in persistent poor glycaemic control: who do we reach?. <i>Patient Education and Counseling</i> , 2005, 56, 313-322.	1.0	30
224	Psychosocial problems and barriers to improved diabetes management: results of the Cross-National Diabetes Attitudes, Wishes and Needs (DAWN) Study. <i>Diabetic Medicine</i> , 2005, 22, 1379-1385.	1.2	728
225	Fat food for a bad mood. Could we treat and prevent depression in Type 2 diabetes by means of omega-3 polyunsaturated fatty acids? A review of the evidence. <i>Diabetic Medicine</i> , 2005, 22, 1465-1475.	1.2	26
226	Short-term effects of cognitive behavioural group training (CBGT) in adult Type 1 diabetes patients in prolonged poor glycaemic control. A randomized controlled trial. <i>Diabetic Medicine</i> , 2005, 22, 1619-1623.	1.2	59
227	Diabetes-related symptoms and negative mood in participants of a targeted population-screening program for type 2 diabetes: The Hoorn Screening study. <i>Quality of Life Research</i> , 2005, 14, 1501-1509.	1.5	39
228	Resistance to Insulin Therapy Among Patients and Providers: Results of the cross-national Diabetes Attitudes, Wishes, and Needs (DAWN) study. <i>Diabetes Care</i> , 2005, 28, 2673-2679.	4.3	709
229	Serious diabetes-specific emotional problems and depression in a Croatian "Dutch" English Survey from the European Depression in Diabetes [EDID] Research Consortium. <i>Diabetes Research and Clinical Practice</i> , 2005, 70, 166-173.	1.1	73
230	No substantial psychological impact of the diagnosis of Type 2 diabetes following targeted population screening: The Hoorn Screening Study. <i>Diabetic Medicine</i> , 2004, 21, 992-998.	1.2	37
231	Health-related quality of life in the first year following diagnosis of Type 2 diabetes: newly diagnosed patients in general practice compared with screening-detected patients. The Hoorn Screening Study. <i>Diabetic Medicine</i> , 2004, 21, 1075-1081.	1.2	38
232	Persistent poor glycaemic control in adult Type 1 diabetes. A closer look at the problem. <i>Diabetic Medicine</i> , 2004, 21, 1263-1268.	1.2	65
233	Health-related quality of life in patients with systemic lupus erythematosus: development and validation of a lupus specific symptom checklist. <i>Quality of Life Research</i> , 2003, 12, 635-644.	1.5	68
234	Rates and risks for co-morbid depression in patients with Type 2 diabetes mellitus: results from a community-based study. <i>Diabetologia</i> , 2003, 46, 892-898.	2.9	174

#	ARTICLE	IF	CITATIONS
235	Combined bedtime insulin-daytime sulphonylurea regimen compared with two different daily insulin regimens in type 2 diabetes: effects on HbA1c and hypoglycaemia rate—a randomised trial. <i>Diabetes/Metabolism Research and Reviews</i> , 2003, 19, 148-152.	1.7	14
236	Improved glycaemic control in type 1 diabetes patients following participation per se in a clinical trial? mechanisms and implications. <i>Diabetes/Metabolism Research and Reviews</i> , 2003, 19, 357-362.	1.7	28
237	Improving quality of life in diabetes: how effective is education?. <i>Patient Education and Counseling</i> , 2003, 51, 1-3.	1.0	21
238	Perceived risk for Type 2 diabetes in participants in a stepwise population-screening programme. <i>Diabetic Medicine</i> , 2003, 20, 210-215.	1.2	52
239	The Confidence in Diabetes Self-Care Scale: Psychometric properties of a new measure of diabetes-specific self-efficacy in Dutch and U.S. patients with type 1 diabetes. <i>Diabetes Care</i> , 2003, 26, 713-718.	4.3	147
240	A Randomized Trial of Continuous Subcutaneous Insulin Infusion and Intensive Injection Therapy in Type 1 Diabetes for Patients With Long-Standing Poor Glycemic Control. <i>Diabetes Care</i> , 2002, 25, 2074-2080.	4.3	174
241	Psychological counselling in problematic diabetes: does it help?. <i>Diabetic Medicine</i> , 2002, 19, 265-273.	1.2	95
242	Screening for Type 2 diabetes: an exploration of subjects' perceptions regarding diagnosis and procedure. <i>Diabetic Medicine</i> , 2002, 19, 406-411.	1.2	43
243	Continuous intraperitoneal insulin infusion in patients with "brittle" diabetes: favourable effects on glycaemic control and hospital stay. <i>Diabetic Medicine</i> , 2002, 19, 496-501.	1.2	39
244	Breaking the barriers to optimal glycaemic control—what physicians need to know from patients' perspectives. <i>International Journal of Clinical Practice, Supplement</i> , 2002, , 80-4.	0.3	17
245	Insulin-treated diabetes patients with fear of self-injecting or fear of self-testing. <i>Journal of Psychosomatic Research</i> , 2001, 51, 665-672.	1.2	115
246	Effects of cognitive behavioural group training (CBGT) in adult patients with poorly controlled insulin-dependent (type 1) diabetes: a pilot study. <i>Patient Education and Counseling</i> , 2001, 45, 143-148.	1.0	61
247	Association between symptoms of depression and glycaemic control may be unstable across gender. <i>Diabetic Medicine</i> , 2001, 18, 595-598.	1.2	49
248	Phobia of self-injecting and self-testing in insulin-treated diabetes patients: opportunities for screening. <i>Diabetic Medicine</i> , 2001, 18, 671-674.	1.2	82
249	Monitoring of Psychological Well-Being in Outpatients With Diabetes: Effects on mood, HbA1c, and the patient's evaluation of the quality of diabetes care: a randomized controlled trial. <i>Diabetes Care</i> , 2001, 24, 1929-1935.	4.3	111
250	The Well-being Questionnaire: evidence for a three-factor structure with 12 items (W-BQ12). <i>Psychological Medicine</i> , 2000, 30, 455-462.	2.7	57
251	Biological and behavioural determinants of the frequency of mild, biochemical hypoglycaemia in patients with Type 1 diabetes on multiple insulin injection therapy. <i>Diabetes/Metabolism Research and Reviews</i> , 2000, 16, 157-163.	1.7	22
252	Diabetes-related emotional distress in Dutch and U.S. diabetic patients: cross-cultural validity of the problem areas in diabetes scale. <i>Diabetes Care</i> , 2000, 23, 1305-1309.	4.3	361

#	ARTICLE	IF	CITATIONS
253	Optimized basal-bolus therapy using a fixed mixture of 75% lispro and 25% NPL insulin in type 1 diabetes patients: no favorable effects on glycemic control, physiological responses to hypoglycemia, well-being, or treatment satisfaction. <i>Diabetes Care</i> , 2000, 23, 629-633.	4.3	22
254	Diabetes Fear of Injecting and Self-Testing Questionnaire: a psychometric evaluation. <i>Diabetes Care</i> , 2000, 23, 765-769.	4.3	77
255	Assessing impaired hypoglycemia awareness in type 1 diabetes: agreement of self-report but not of field study data with the autonomic symptom threshold during experimental hypoglycemia. <i>Diabetes Care</i> , 2000, 23, 529-532.	4.3	51
256	Biological and behavioural determinants of the frequency of mild, biochemical hypoglycaemia in patients with Type 1 diabetes on multiple insulin injection therapy. , 2000, 16, 157.		2
257	The 12-item well-being questionnaire. An evaluation of its validity and reliability in Dutch people with diabetes. <i>Diabetes Care</i> , 1999, 22, 2004-2010.	4.3	74
258	Substitution of night-time continuous subcutaneous insulin infusion therapy for bedtime NPH insulin in a multiple injection regimen improves counterregulatory hormonal responses and warning symptoms of hypoglycaemia in IDDM. <i>Diabetologia</i> , 1998, 41, 322-329.	2.9	68
259	Randomized Study of Two Different Target Levels of Glycemic Control Within the Acceptable Range in Type 2 Diabetes: Effects on well-being at 1 year. <i>Diabetes Care</i> , 1998, 21, 2085-2093.	4.3	34
260	Well-Being and Symptoms in Relation to Insulin Therapy in Type 2 Diabetes. <i>Diabetes Care</i> , 1998, 21, 919-924.	4.3	76
261	No Differences in Attentional Functioning Between Type 1 Diabetic Patients With and Without a History of Severe Hypoglycemia. <i>Diabetes Care</i> , 1998, 21, 1568-1569.	4.3	3
262	Ceiling effect reduces the validity of the Diabetes Treatment Satisfaction Questionnaire. <i>Diabetes Care</i> , 1998, 21, 2039-2039.	4.3	22
263	Development and validation of the diabetes fear of injecting and self-testing questionnaire (D-FISQ): first findings. , 1997, 14, 871-876.		50
264	Hypoglycaemia induces emotional disruption. <i>Patient Education and Counseling</i> , 1996, 29, 117-122.	1.0	25
265	Assessment of perceived barriers in self-care of insulinrequiring diabetic patients. <i>Patient Education and Counseling</i> , 1996, 29, 277-281.	1.0	16
266	Symptoms and Well-Being in Relation to Glycemic Control in Type II Diabetes. <i>Diabetes Care</i> , 1996, 19, 204-210.	4.3	235
267	Quality of life and neuropsychological functions in long-term low-grade glioma survivors. <i>International Journal of Radiation Oncology Biology Physics</i> , 1994, 29, 1201-1202.	0.4	15
268	Development of a Type 2 Diabetes Symptom Checklist: a Measure of Symptom Severity. <i>Diabetic Medicine</i> , 1994, 11, 253-261.	1.2	164
269	Fatigue, sleep disturbances and circadian rhythm in multiple sclerosis. <i>Journal of Neurology</i> , 1993, 240, 446-448.	1.8	85
270	Blood Glucose Awareness Training. , 0, , 169-206.		20