

Shunsuke Kondo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2186372/publications.pdf>

Version: 2024-02-01

104
papers

2,338
citations

257450

24
h-index

265206

42
g-index

108
all docs

108
docs citations

108
times ranked

4175
citing authors

#	ARTICLE	IF	CITATIONS
1	Nivolumab alone or in combination with cisplatin plus gemcitabine in Japanese patients with unresectable or recurrent biliary tract cancer: a non-randomised, multicentre, open-label, phase 1 study. <i>The Lancet Gastroenterology and Hepatology</i> , 2019, 4, 611-621.	8.1	223
2	Cisplatin and Etoposide as First-line Chemotherapy for Poorly Differentiated Neuroendocrine Carcinoma of the Hepatobiliary Tract and Pancreas. <i>Japanese Journal of Clinical Oncology</i> , 2010, 40, 313-318.	1.3	149
3	Phase I study of bintrafusp alfa, a bifunctional fusion protein targeting TGF- β 2 and PD-L1, in patients with pretreated biliary tract cancer. , 2020, 8, e000564.		98
4	Phase 1 Trial of Wilms Tumor 1 (WT1) Peptide Vaccine and Gemcitabine Combination Therapy in Patients With Advanced Pancreatic or Biliary Tract Cancer. <i>Journal of Immunotherapy</i> , 2011, 34, 92-99.	2.4	91
5	Clinical impact of c-Met expression and its gene amplification in hepatocellular carcinoma. <i>International Journal of Clinical Oncology</i> , 2013, 18, 207-213.	2.2	75
6	Utility of Assessing the Number of Mutated KRAS, CDKN2A, TP53, and SMAD4 Genes Using a Targeted Deep Sequencing Assay as a Prognostic Biomarker for Pancreatic Cancer. <i>Pancreas</i> , 2017, 46, 335-340.	1.1	75
7	Clinical impact of pentraxin family expression on prognosis of pancreatic carcinoma. <i>British Journal of Cancer</i> , 2013, 109, 739-746.	6.4	65
8	Homozygous CDA*3 is a major cause of life-threatening toxicities in gemcitabine-treated Japanese cancer patients. <i>British Journal of Cancer</i> , 2009, 100, 870-873.	6.4	56
9	Highly Sensitive Circulating MicroRNA Panel for Accurate Detection of Hepatocellular Carcinoma in Patients With Liver Disease. <i>Hepatology Communications</i> , 2020, 4, 284-297.	4.3	53
10	Clinical Development of Immune Checkpoint Inhibitors. <i>BioMed Research International</i> , 2015, 2015, 1-12.	1.9	51
11	Phase 1 study of abemaciclib, an inhibitor of CDK 4 and 6, as a single agent for Japanese patients with advanced cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 78, 281-288.	2.3	51
12	First-in-Human Phase I Study of an Oral HSP90 Inhibitor, TAS-116, in Patients with Advanced Solid Tumors. <i>Molecular Cancer Therapeutics</i> , 2019, 18, 531-540.	4.1	49
13	Comparison of Chemotherapeutic Treatment Outcomes of Advanced Extrapulmonary Neuroendocrine Carcinomas and Advanced Small-Cell Lung Carcinoma. <i>Neuroendocrinology</i> , 2012, 96, 324-332.	2.5	48
14	Population Pharmacokinetics of Gemcitabine and Its Metabolite in Japanese Cancer Patients. <i>Clinical Pharmacokinetics</i> , 2010, 49, 549-558.	3.5	43
15	Impact of the Integrin Signaling Adaptor Protein NEDD9 on Prognosis and Metastatic Behavior of Human Lung Cancer. <i>Clinical Cancer Research</i> , 2012, 18, 6326-6338.	7.0	43
16	Spontaneous regression of hepatocellular carcinoma. <i>International Journal of Clinical Oncology</i> , 2006, 11, 407-411.	2.2	42
17	Phase II study of sunitinib in Japanese patients with unresectable or metastatic, well-differentiated pancreatic neuroendocrine tumor. <i>Investigational New Drugs</i> , 2013, 31, 1265-1274.	2.6	39
18	C-Reactive Protein Level Is an Indicator of the Aggressiveness of Advanced Pancreatic Cancer. <i>Pancreas</i> , 2016, 45, 110-116.	1.1	37

#	ARTICLE	IF	CITATIONS
19	Phase 1b study of galunisertib in combination with gemcitabine in Japanese patients with metastatic or locally advanced pancreatic cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2017, 79, 1169-1177.	2.3	37
20	Efficacy of sorafenib in patients with hepatocellular carcinoma refractory to transcatheter arterial chemoembolization. <i>Journal of Gastroenterology</i> , 2014, 49, 932-940.	5.1	36
21	Survey of survival among patients with hepatitis C virus-related hepatocellular carcinoma treated with peretinoin, an acyclic retinoid, after the completion of a randomized, placebo-controlled trial. <i>Journal of Gastroenterology</i> , 2015, 50, 667-674.	5.1	36
22	Construction and Validation of a Prognostic Index for Patients With Metastatic Pancreatic Adenocarcinoma. <i>Pancreas</i> , 2011, 40, 415-421.	1.1	35
23	Human papillomavirus infection and immunohistochemical expression of cell cycle proteins pRb, p53, and p16INK4a in sinonasal diseases. <i>Infectious Agents and Cancer</i> , 2015, 10, 23.	2.6	35
24	Do Recurrent and Metastatic Pancreatic Cancer Patients Have the Same Outcomes with Gemcitabine Treatment?. <i>Oncology</i> , 2009, 77, 217-223.	1.9	28
25	Transcatheter Arterial Infusion Chemotherapy with a Fine-powder Formulation of Cisplatin for Advanced Hepatocellular Carcinoma Refractory to Transcatheter Arterial Chemoembolization. <i>Japanese Journal of Clinical Oncology</i> , 2011, 41, 770-775.	1.3	28
26	First-In-Human Phase I Study of a Next-Generation, Oral, TGF β 2 Receptor 1 Inhibitor, LY3200882, in Patients with Advanced Cancer. <i>Clinical Cancer Research</i> , 2021, 27, 6666-6676.	7.0	27
27	Phase Ia/Ib study of the pan-class I PI3K inhibitor pictilisib (GDC-0941) administered as a single agent in Japanese patients with solid tumors and in combination in Japanese patients with non-squamous non-small cell lung cancer. <i>Investigational New Drugs</i> , 2017, 35, 37-46.	2.6	26
28	Pancreatic neuroendocrine tumors: A single-center 20-year experience with 100 patients. <i>Pancreatology</i> , 2016, 16, 99-105.	1.1	25
29	Incidence and risk factors for venous thromboembolism in patients with pretreated advanced pancreatic carcinoma. <i>Oncotarget</i> , 2018, 9, 16883-16890.	1.8	25
30	Trends in the development of MET inhibitors for hepatocellular carcinoma. <i>Future Oncology</i> , 2016, 12, 1275-1286.	2.4	24
31	An Oncogenic <i>ALK</i> Fusion and an <i>RRAS</i> Mutation in <i>KRAS</i> Mutation-Negative Pancreatic Ductal Adenocarcinoma. <i>Oncologist</i> , 2017, 22, 158-164.	3.7	24
32	Safety and Tolerability of Bintrafusp Alfa, a Bifunctional Fusion Protein Targeting TGF β 2 and PD-L1, in Asian Patients with Pretreated Recurrent or Refractory Gastric Cancer. <i>Clinical Cancer Research</i> , 2020, 26, 3202-3210.	7.0	24
33	Treatment Efficacy/Safety and Prognostic Factors in Patients with Advanced Biliary Tract Cancer Receiving Gemcitabine Monotherapy: An Analysis of 100 Cases. <i>Oncology</i> , 2010, 79, 39-45.	1.9	23
34	Complicated paraneoplastic neurological syndromes: a report of two patients with small cell or non-small cell lung cancer. <i>Clinical Neurology and Neurosurgery</i> , 2003, 106, 47-49.	1.4	22
35	Phase 1/2 study assessing the safety and efficacy of dabrafenib and trametinib combination therapy in Japanese patients with <i>BRAF</i> V600 mutation-positive advanced cutaneous melanoma. <i>Journal of Dermatology</i> , 2018, 45, 397-407.	1.2	22
36	Cytotoxic chemotherapy for pancreatic neuroendocrine tumors. <i>Journal of Hepato-Biliary-Pancreatic Sciences</i> , 2015, 22, 628-633.	2.6	20

#	ARTICLE	IF	CITATIONS
37	Lipid profiling of pre-treatment plasma reveals biomarker candidates associated with response rates and hand-foot skin reactions in sorafenib-treated patients. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 82, 677-684.	2.3	20
38	Phase I study of resminostat, an HDAC inhibitor, combined with S-1 in patients with pre-treated biliary tract or pancreatic cancer. <i>Investigational New Drugs</i> , 2019, 37, 109-117.	2.6	20
39	Cabozantinib in Japanese patients with advanced hepatocellular carcinoma: a phase 2 multicenter study. <i>Journal of Gastroenterology</i> , 2021, 56, 181-190.	5.1	20
40	Chemotherapy for advanced poorly differentiated pancreatic neuroendocrine carcinoma. <i>Journal of Hepato-Biliary-Pancreatic Sciences</i> , 2015, 22, 623-627.	2.6	18
41	A phase I and pharmacokinetic study of taladegib, a Smoothed inhibitor, in Japanese patients with advanced solid tumors. <i>Investigational New Drugs</i> , 2018, 36, 647-656.	2.6	17
42	Phase I study of the indoleamine 2,3-dioxygenase 1 inhibitor navoximod (GDC-0919) as monotherapy and in combination with the PD-L1 inhibitor atezolizumab in Japanese patients with advanced solid tumours. <i>Investigational New Drugs</i> , 2020, 38, 468-477.	2.6	17
43	Circulating endothelial cells and other angiogenesis factors in pancreatic carcinoma patients receiving gemcitabine chemotherapy. <i>BMC Cancer</i> , 2012, 12, 268.	2.6	16
44	Assessment of adverse events via a telephone consultation service for cancer patients receiving ambulatory chemotherapy. <i>BMC Research Notes</i> , 2015, 8, 315.	1.4	16
45	Phase I studies of peptide vaccine cocktails derived from GPC3, WDRPUH and NEIL3 for advanced hepatocellular carcinoma. <i>Immunotherapy</i> , 2021, 13, 371-385.	2.0	16
46	Bintrafusp Alfa, a Bifunctional Fusion Protein Targeting TGF β 2 and PD-L1, in Patients with Esophageal Squamous Cell Carcinoma: Results from a Phase 1 Cohort in Asia. <i>Targeted Oncology</i> , 2021, 16, 447-459.	3.6	16
47	First-in-human phase I study of E7090, a novel selective fibroblast growth factor receptor inhibitor, in patients with advanced solid tumors. <i>Cancer Science</i> , 2020, 111, 571-579.	3.9	16
48	Cytogenetic Confirmation of a Gastrointestinal Stromal Tumor and Ewing Sarcoma/Primitive Neuroectodermal Tumor in a Single Patient. <i>Japanese Journal of Clinical Oncology</i> , 2005, 35, 753-756.	1.3	15
49	Evaluating Clinical Genome Sequence Analysis by Watson for Genomics. <i>Frontiers in Medicine</i> , 2018, 5, 305.	2.6	15
50	Immune checkpoint and inflammation as therapeutic targets in pancreatic carcinoma. <i>World Journal of Gastroenterology</i> , 2016, 22, 7440.	3.3	15
51	Multicenter cooperative case survey of hepatitis B virus reactivation by chemotherapeutic agents. <i>Hepatology Research</i> , 2015, 45, 1220-1227.	3.4	14
52	M7824 (MSB0011359C), a bifunctional fusion protein targeting PD-L1 and TGF- β 2, in Asian patients with pretreated biliary tract cancer: Preliminary results from a phase I trial. <i>Annals of Oncology</i> , 2018, 29, viii258-viii259.	1.2	14
53	Oral chemotherapy for the treatment of hepatocellular carcinoma. <i>Expert Opinion on Pharmacotherapy</i> , 2018, 19, 993-1001.	1.8	13
54	Gemcitabine-induced Pleuropericardial Effusion in a Patient with Pancreatic Cancer. <i>Japanese Journal of Clinical Oncology</i> , 2012, 42, 845-850.	1.3	12

#	ARTICLE	IF	CITATIONS
55	Tremelimumab-associated tumor regression following after initial progression: two case reports. <i>Immunotherapy</i> , 2016, 8, 9-15.	2.0	12
56	Salvage chemoradiotherapy after primary chemotherapy for locally advanced pancreatic cancer: a single-institution retrospective analysis. <i>BMC Cancer</i> , 2012, 12, 609.	2.6	11
57	Twenty-six Cases of Advanced Ampullary Adenocarcinoma Treated with Systemic Chemotherapy. <i>Japanese Journal of Clinical Oncology</i> , 2014, 44, 324-330.	1.3	11
58	Transarterial (Chemo)Embolization for Liver Metastases in Patients with Neuroendocrine Tumors. <i>Oncology</i> , 2017, 92, 353-359.	1.9	11
59	Prognostic Factors for Survival in Patients with Advanced Intrahepatic Cholangiocarcinoma Treated with Gemcitabine plus Cisplatin as First-Line Treatment. <i>Oncology</i> , 2018, 94, 72-78.	1.9	11
60	Updated results from a phase I trial of M7824 (MSB0011359C), a bifunctional fusion protein targeting PD-L1 and TGF- β 2, in patients with pretreated recurrent or refractory gastric cancer. <i>Annals of Oncology</i> , 2018, 29, viii222-viii223.	1.2	11
61	Transarterial infusion chemotherapy with cisplatin plus S-1 for hepatocellular carcinoma treatment: a phase I trial. <i>BMC Cancer</i> , 2014, 14, 301.	2.6	10
62	Efficacy of radiotherapy for primary tumor in patients with unresectable pancreatic neuroendocrine tumors. <i>Japanese Journal of Clinical Oncology</i> , 2017, 47, 826-831.	1.3	10
63	Glycemia Control Using A1C Level in Terminal Cancer Patients with Preexisting Type 2 Diabetes. <i>Journal of Palliative Medicine</i> , 2013, 16, 790-793.	1.1	9
64	Phase I study of combination chemotherapy using sorafenib and transcatheter arterial infusion with cisplatin for advanced hepatocellular carcinoma. <i>Cancer Science</i> , 2014, 105, 354-358.	3.9	9
65	Germline mutations in cancer-predisposition genes in patients with biliary tract cancer. <i>Oncotarget</i> , 2019, 10, 5949-5957.	1.8	9
66	Phase I/II study of gemcitabine as a fixed dose rate infusion and S-1 combination therapy (FGS) in gemcitabine-refractory pancreatic cancer patients. <i>Cancer Chemotherapy and Pharmacology</i> , 2012, 69, 957-964.	2.3	8
67	Safety of BI 754111, an anti-LAG-3 monoclonal antibody (mAb), in combination with BI 754091, an anti-PD-1 mAb, in patients with advanced solid tumors.. <i>Journal of Clinical Oncology</i> , 2020, 38, 3063-3063.	1.6	8
68	SHP-2 inhibits tyrosine phosphorylation of Cas-L and regulates cell migration. <i>Biochemical and Biophysical Research Communications</i> , 2009, 382, 210-214.	2.1	7
69	Hepatitis B Virus Reactivation during Treatment with Multi-Tyrosine Kinase Inhibitor for Hepatocellular Carcinoma. <i>Case Reports in Oncology</i> , 2012, 5, 515-519.	0.7	7
70	Phase I clinical trial of oral administration of S-1 in combination with intravenous gemcitabine and cisplatin in patients with advanced biliary tract cancer. <i>Japanese Journal of Clinical Oncology</i> , 2016, 46, hv179.	1.3	7
71	Germline variants in pancreatic cancer patients with a personal or family history of cancer fulfilling the revised Bethesda guidelines. <i>Journal of Gastroenterology</i> , 2018, 53, 1159-1167.	5.1	7
72	Novel endoscopic technique for trisegment drainage in patients with unresectable hilar malignant biliary strictures (with video). <i>Gastrointestinal Endoscopy</i> , 2020, 92, 763-769.	1.0	7

#	ARTICLE	IF	CITATIONS
73	Long-Term Administration of Wilms Tumor-1 Peptide Vaccine in Combination with Gemcitabine Causes Severe Local Skin Inflammation at Injection Sites. <i>Japanese Journal of Clinical Oncology</i> , 2010, 40, 1184-1188.	1.3	6
74	Gemcitabine in Patients With Intraductal Papillary Mucinous Neoplasm With an Associated Invasive Carcinoma of the Pancreas. <i>Pancreas</i> , 2013, 42, 889-892.	1.1	6
75	73P Long-term follow-up of bintrafusp alfa, a bifunctional fusion protein targeting TGF- β 2 and PD-L1, in patients with pretreated biliary tract cancer. <i>Annals of Oncology</i> , 2020, 31, S268-S269.	1.2	6
76	Dose exploration results from Phase 1 study of cemiplimab, a human monoclonal programmed death (PD)-1 antibody, in Japanese patients with advanced malignancies. <i>Cancer Chemotherapy and Pharmacology</i> , 2021, 87, 53-64.	2.3	6
77	First-in-human study of the cancer peptide vaccine TAS0313 in patients with advanced solid tumors. <i>Cancer Science</i> , 2021, 112, 1514-1523.	3.9	6
78	Safety, pharmacokinetics, and efficacy of budigalimab with rovalpituzumab tesirine in patients with small cell lung cancer. <i>Cancer Treatment and Research Communications</i> , 2021, 28, 100405.	1.7	6
79	Phase I/II study of streptozocin monotherapy in Japanese patients with unresectable or metastatic gastroenteropancreatic neuroendocrine tumors. <i>Japanese Journal of Clinical Oncology</i> , 2022, 52, 716-724.	1.3	6
80	Treatment outcome for systemic chemotherapy for recurrent pancreatic cancer after postoperative adjuvant chemotherapy. <i>Pancreatology</i> , 2012, 12, 428-433.	1.1	5
81	A retrospective analysis of factors associated with selection of end-of-life care and actual place of death for patients with cancer. <i>BMJ Open</i> , 2014, 4, e004352.	1.9	5
82	Phase I study on the safety, pharmacokinetic profile, and efficacy of the combination of TSU-68, an oral antiangiogenic agent, and S-1 in patients with advanced hepatocellular carcinoma. <i>Investigational New Drugs</i> , 2014, 32, 928-936.	2.6	5
83	Model Informed Dosing Regimen and Phase I Results of the Anti- PD-1 Antibody Budigalimab (ABBV-181). <i>Clinical and Translational Science</i> , 2021, 14, 277-287.	3.1	5
84	Successful Control of Intractable Hypoglycemia Using Radiopharmaceutical Therapy with Strontium-89 in a Case with Malignant Insulinoma and Bone Metastases. <i>Japanese Journal of Clinical Oncology</i> , 2012, 42, 640-645.	1.3	4
85	Global trends in the distribution of cancer types among patients in oncology phase I trials, 1991-2015. <i>Investigational New Drugs</i> , 2019, 37, 166-174.	2.6	4
86	Impact of the Duration of Diabetes Mellitus on the Outcome of Metastatic Pancreatic Cancer Treated with Gemcitabine: A Retrospective Study. <i>Internal Medicine</i> , 2019, 58, 2435-2441.	0.7	4
87	Phase 1 dose-escalation study of a novel oral PI3K/mTOR dual inhibitor, LY3023414, in patients with cancer. <i>Investigational New Drugs</i> , 2020, 38, 1836-1845.	2.6	4
88	Merestinib monotherapy or in combination for Japanese patients with advanced and/or metastatic cancer: A phase 1 study. <i>Cancer Medicine</i> , 2021, 10, 6579-6589.	2.8	4
89	A multicenter, open-label, phase I study of nivolumab alone or in combination with gemcitabine plus cisplatin in patients with unresectable or recurrent biliary tract cancer. <i>Journal of Clinical Oncology</i> , 2019, 37, 306-306.	1.6	4
90	Targeted-sequencing in rare cancers and the impact on patient treatment. <i>Journal of Clinical Oncology</i> , 2019, 37, e14755-e14755.	1.6	3

#	ARTICLE	IF	CITATIONS
91	Quality evaluation of investigator-initiated trials using post-approval cancer drugs in Japan. <i>Cancer Science</i> , 2017, 108, 995-999.	3.9	2
92	AB053. P-21. M7824 (MSB0011359C), a bifunctional fusion protein targeting transforming growth factor β (TGF- β) and PD-L1, in Asian patients with pretreated biliary tract cancer (BTC): efficacy by BTC subtype. <i>Hepatobiliary Surgery and Nutrition</i> , 2019, 8, AB053-AB053.	1.5	2
93	An open-label phase 1 trial of lenvatinib plus pembrolizumab in patients with advanced selected solid tumors. <i>Annals of Oncology</i> , 2018, 29, vii81.	1.2	1
94	Impact of Hepatitis Virus on the Feasibility and Efficacy of Anticancer Agents in Patients With Hepatocellular Carcinoma in Phase I Clinical Trials. <i>Frontiers in Oncology</i> , 2019, 9, 301.	2.8	1
95	Preliminary Phase 1 Results of the PD-1 Inhibitor ABBV-181 in Japanese vs Western Patients With Advanced Solid Tumors. <i>Annals of Oncology</i> , 2019, 30, vi107.	1.2	1
96	Improved survival among patients enrolled in oncology phase 1 trials in recent decades. <i>Cancer Chemotherapy and Pharmacology</i> , 2020, 85, 449-459.	2.3	1
97	An open-label, phase I trial of BI 754091 alone and in combination with BI 754111 in Asian patients (pts) with advanced solid tumors.. <i>Journal of Clinical Oncology</i> , 2020, 38, 3054-3054.	1.6	1
98	Molecular-targeted Therapies in Gastrointestinal Cancer. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2016, 105, 1051-1060.	0.0	0
99	Do all patients in the phase I oncology trials need to be hospitalized? Domestic but outstanding issues for globalization of drug development in Japan. <i>International Journal of Clinical Oncology</i> , 2017, 22, 780-785.	2.2	0
100	A phase 1 study of niraparib in Japanese patients with advanced solid tumors. <i>Annals of Oncology</i> , 2019, 30, vi127.	1.2	0
101	Phase I study of nivolumab or nivolumab/cisplatin/gemcitabine to treat unresectable/recurrent biliary tract cancer. <i>Annals of Oncology</i> , 2019, 30, vi86-vi87.	1.2	0
102	A Case of Pathological Complete Response after Combination Chemotherapy by Sorafenib and Cisplatin Hepatic Arterial Infusion for an Advanced Hepatocellular Carcinoma. <i>Japanese Journal of Gastroenterological Surgery</i> , 2013, 46, 915-923.	0.1	0
103	Phase I study of cemiplimab, a human monoclonal antibody to programmed death (PD)-1, in Japanese patients (pts) with advanced malignancies: Results from the dose exploration.. <i>Journal of Clinical Oncology</i> , 2019, 37, 33-33.	1.6	0
104	First-in-human study of the cancer peptide vaccine, TAS0313, in patients with advanced solid tumors: Phase I dose-finding part results.. <i>Journal of Clinical Oncology</i> , 2020, 38, 73-73.	1.6	0