

Steinar Hunskaar

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2204240/publications.pdf>

Version: 2024-02-01

57
papers

2,656
citations

279798
23
h-index

182427
51
g-index

57
all docs

57
docs citations

57
times ranked

2261
citing authors

#	ARTICLE	IF	CITATIONS
1	A severity index for epidemiological surveys of female urinary incontinence: Comparison with 48-hour pad-weighting tests. <i>Neurourology and Urodynamics</i> , 2000, 19, 137-145.	1.5	401
2	Are smoking and other lifestyle factors associated with female urinary incontinence? The Norwegian EPINCONT Study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2003, 110, 247-254.	2.3	394
3	Epidemiology and natural history of urinary incontinence in women. <i>Urology</i> , 2003, 62, 16-23.	1.0	278
4	A systematic review of overweight and obesity as risk factors and targets for clinical intervention for urinary incontinence in women. <i>Neurourology and Urodynamics</i> , 2008, 27, 749-757.	1.5	246
5	Help-seeking and associated factors in female urinary incontinence. The Norwegian EPINCONT Study. <i>Scandinavian Journal of Primary Health Care</i> , 2002, 20, 102-107.	1.5	108
6	Continuity in general practice as predictor of mortality, acute hospitalisation, and use of out-of-hours care: a registry-based observational study in Norway. <i>British Journal of General Practice</i> , 2022, 72, e84-e90.	1.4	106
7	Anxiety and depression associated with urinary incontinence. A 10-year follow-up study from the Norwegian HUNT study (EPINCONT). <i>Neurourology and Urodynamics</i> , 2017, 36, 322-328.	1.5	93
8	Iron supplementation in pregnancy: is less enough?. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1997, 76, 822-828.	2.8	78
9	Point-of-care ultrasound in primary care: a systematic review of generalist performed point-of-care ultrasound in unselected populations. <i>Ultrasound Journal</i> , 2019, 11, 31.	3.3	78
10	Female Urinary Incontinence. <i>Scandinavian Journal of Caring Sciences</i> , 1993, 7, 53-56.	2.1	60
11	Why do only a minority of perimenopausal women with urinary incontinence consult a doctor?. <i>Scandinavian Journal of Primary Health Care</i> , 1994, 12, 180-183.	1.5	56
12	Risk of hip fracture among older people using antihypertensive drugs: a nationwide cohort study. <i>BMC Geriatrics</i> , 2015, 15, 153.	2.7	54
13	One hundred and fifty men with urinary incontinence: III. Psychosocial consequences. <i>Scandinavian Journal of Primary Health Care</i> , 1993, 11, 193-196.	1.5	43
14	A Study of Female Urinary Incontinence in General Practice. <i>Scandinavian Journal of Urology and Nephrology</i> , 1996, 30, 465-471.	1.4	41
15	Telephone triage by nurses in primary care out-of-hours services in Norway: an evaluation study based on written case scenarios. <i>BMJ Quality and Safety</i> , 2011, 20, 390-396.	3.7	40
16	Healthcare personnel's attitudes towards patients with urinary incontinence INFORMATION POINT: Regression analysis. <i>Journal of Clinical Nursing</i> , 2001, 10, 455-462.	3.0	37
17	Distance and utilisation of out-of-hours services in a Norwegian urban/rural district: an ecological study. <i>BMC Health Services Research</i> , 2013, 13, 222.	2.2	37
18	Are resistance patterns in uropathogens published by microbiological laboratories valid for general practice?. <i>Apmis</i> , 1999, 107, 676-680.	2.0	32

#	ARTICLE	IF	CITATIONS
19	Development, implementation, and pilot study of a sentinel network ("The Watchtowers") for monitoring emergency primary health care activity in Norway. BMC Health Services Research, 2008, 8, 62.	2.2	31
20	Epidemiology of nocturia. BJU International, 2005, 96, 4-7.	2.5	30
21	Scaling up family medicine training in Gezira, Sudan – a 2-year in-service master programme using modern information and communication technology: a survey study. Human Resources for Health, 2014, 12, 3.	3.1	28
22	Sentinel monitoring of activity of out-of-hours services in Norway in 2007: an observational study. BMC Health Services Research, 2009, 9, 123.	2.2	27
23	Long-term risks of stress and urgency urinary incontinence after different vaginal delivery modes. American Journal of Obstetrics and Gynecology, 2019, 220, 181.e1-181.e8.	1.3	27
24	General practitioners' and out-of-hours doctors' role as gatekeeper in emergency admissions to somatic hospitals in Norway: registry-based observational study. BMC Health Services Research, 2019, 19, 568.	2.2	23
25	Out-of-hours antibiotic prescription after screening with C reactive protein: a randomised controlled study. BMJ Open, 2016, 6, e011231.	1.9	21
26	Validity of a scored urological history in detecting detrusor instability in female urinary incontinence. Acta Obstetrica Et Gynecologica Scandinavica, 1996, 75, 941-945.	2.8	20
27	Comparing unplanned and potentially planned home deaths: a population-based cross-sectional study. BMC Palliative Care, 2018, 17, 69.	1.8	20
28	Allergy to laboratory mice and rats: a review of its prevention, management, and treatment. Laboratory Animals, 1993, 27, 206-221.	1.0	18
29	Understanding of and adherence to advice after telephone counselling by nurse: a survey among callers to a primary emergency out-of-hours service in Norway. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine, 2011, 19, 48.	2.6	18
30	Urinary incontinence associated with anxiety and depression: the impact of psychotropic drugs in a cross-sectional study from the Norwegian HUNT study. BMC Psychiatry, 2020, 20, 521.	2.6	18
31	General Practitioners' Attitudes to Oestrogen Prescription in the Menopause: A National Survey in Norway. Scandinavian Journal of Primary Health Care, 1992, 10, 179-184.	1.5	17
32	One Hundred and Fifty Men with Urinary Incontinence II. Help Seeking and Self Care. Scandinavian Journal of Primary Health Care, 1992, 10, 26-29.	1.5	15
33	Loss of life years due to unavailable helicopter emergency medical service: a single base study from a rural area of Norway. Scandinavian Journal of Primary Health Care, 2019, 37, 233-241.	1.5	14
34	One Hundred and Fifty Men with Urinary Incontinence I. Demography and Medical History. Scandinavian Journal of Primary Health Care, 1992, 10, 21-25.	1.5	12
35	Point-of-care testing with CRP in primary care: a registry-based observational study from Norway. BMC Family Practice, 2015, 16, 170.	2.9	12
36	Dispatch guideline adherence and response interval – a study of emergency medical calls in Norway. BMC Emergency Medicine, 2016, 16, 40.	1.9	12

#	ARTICLE	IF	CITATIONS
37	Direct mailing of consensus recommendations did not alter GPs' knowledge and prescription of oestrogen in the menopause. <i>Scandinavian Journal of Primary Health Care</i> , 1996, 14, 203-208.	1.5	11
38	General practitioners' provision of end-of-life care and associations with dying at home: a registry-based longitudinal study. <i>Family Practice</i> , 2020, 37, 340-347.	1.9	11
39	Telephone counselling by nurses in Norwegian primary care out-of-hours services: a cross-sectional study. <i>BMC Family Practice</i> , 2017, 18, 84.	2.9	9
40	Patient trajectories in a Norwegian unit of municipal emergency beds. <i>Scandinavian Journal of Primary Health Care</i> , 2017, 35, 137-142.	1.5	8
41	Associations between home deaths and end-of-life nursing care trajectories for community-dwelling people: a population-based registry study. <i>BMC Health Services Research</i> , 2019, 19, 698.	2.2	8
42	Construction and validation of a morbidity index based on the International Classification of Primary Care. <i>Scandinavian Journal of Primary Health Care</i> , 2022, 40, 305-312.	1.5	8
43	Continuity of care for patients with chronic disease: a registry-based observational study from Norway. <i>Family Practice</i> , 2022, 39, 570-578.	1.9	7
44	Method Paper "Distance and Travel Time to Casualty Clinics in Norway Based on Crowdsourced Postcode Coordinates: A Comparison with Other Methods. <i>PLoS ONE</i> , 2014, 9, e89287.	2.5	7
45	General practitioners' management of female urinary incontinence: Medical records do not reflect patients' recall. <i>Scandinavian Journal of Primary Health Care</i> , 1995, 13, 168-174.	1.5	6
46	Treatment, transport, and primary care involvement when helicopter emergency medical services are inaccessible: a retrospective study. <i>Scandinavian Journal of Primary Health Care</i> , 2018, 36, 397-405.	1.5	6
47	Primary care utilization among patients with influenza during the 2009 pandemic. Does risk for severe influenza disease or prior contact with the general practitioner have any influence?. <i>Family Practice</i> , 2015, 32, 56-61.	1.9	4
48	Confidence in Procedural Skills before and after a Two-Year Master's Programme in Family Medicine in Gezira State, Sudan. <i>Advances in Medicine</i> , 2017, 2017, 1-11.	0.8	4
49	Female urinary incontinence - the role of the general practitioner. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2000, 79, 1046-1051.	2.8	3
50	Medical consequences of violence: a two-year prospective study from a primary care accident and emergency department in Norway. <i>Scandinavian Journal of Primary Health Care</i> , 2002, 20, 108-112.	1.5	3
51	The market lifecycle of duloxetine for urinary incontinence in Norway. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2010, 89, 217-222.	2.8	3
52	Impact on core values of family medicine from a 2-year Master's programme in Gezira, Sudan: observational study. <i>BMC Family Practice</i> , 2019, 20, 145.	2.9	3
53	Helicopter emergency medical service (HEMS) activity after increased distance to out-of-hours services: an observational study from Norway. <i>BMC Emergency Medicine</i> , 2020, 20, 88.	1.9	3
54	Are smoking and other lifestyle factors associated with female urinary incontinence? The Norwegian EPINCONT Study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2003, 110, 247-254.	2.3	3

#	ARTICLE	IF	CITATIONS
55	Focus groups as a path to clinical knowledge about the acutely and severely ill child. Scandinavian Journal of Primary Health Care, 1997, 15, 26-29.	1.5	2
56	From direct attendance to telephone triage in an emergency primary healthcare service: an observational study. BMJ Open, 2022, 12, e054046.	1.9	2
57	Increased acceptance of HRT and improved level of information: a change in Norwegian women's opinion from 1990 to 1997. Acta Obstetricia Et Gynecologica Scandinavica, 2001, 80, 623-627.	2.8	0