


List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2211138/publications.pdf Version: 2024-02-01



Ιτινι Υλινι

#	Article	IF	CITATIONS
1	Reduced lκB-α Protein Levels in Peripheral Blood Cells of Patients with Multiple Sclerosis—A Possible Cause of Constitutive NF-κB Activation. Journal of Clinical Medicine, 2020, 9, 2534.	2.4	2
2	Increased constitutive activation of NF-κB p65 (RelA) in peripheral blood cells of patients with progressive multiple sclerosis. Journal of Neuroimmunology, 2018, 320, 111-116.	2.3	13
3	NF-κB Pathways in the Pathogenesis of Multiple Sclerosis and the Therapeutic Implications. Frontiers in Molecular Neuroscience, 2016, 9, 84.	2.9	88
4	Circulating brain derived neurotrophic factor (BDNF) and frequency of BDNF positive T cells in peripheral blood in human ischemic stroke: Effect on outcome. Journal of Neuroimmunology, 2015, 286, 42-47.	2.3	47
5	Correlation of Adrenomedullin gene expression in peripheral blood leukocytes with severity of ischemic stroke. International Journal of Neuroscience, 2014, 124, 271-280.	1.6	10
6	Increased expression of the hypoxiaâ€related genes in peripheral blood leukocytes of human subjects with acute ischemic stroke. Clinical and Experimental Neuroimmunology, 2014, 5, 216-226.	1.0	2
7	A Simple and Reliable Immunohistochemical Method for Colocalization of 2 Antigens in the Same Cells of Paraffin-embedded Tissues. Applied Immunohistochemistry and Molecular Morphology, 2013, 21, 471-477.	1.2	1
8	Levels of interleukinÂ33 and soluble suppression of tumorigenicityÂ2 in acute ischemic stroke. Clinical and Experimental Neuroimmunology, 2013, 4, 339-347.	1.0	1
9	Interleukin-6 Gene Promoter-572 C Allele May Play a Role in Rate of Disease Progression in Multiple Sclerosis. International Journal of Molecular Sciences, 2012, 13, 13667-13679.	4.1	17
10	Frequency and function of regulatory T cells after ischaemic stroke in humans. Journal of Neuroimmunology, 2012, 243, 89-94.	2.3	70
11	Prolonged elevation of cytokine levels after human acute ischaemic stroke with evidence of individual variability. Journal of Neuroimmunology, 2012, 246, 78-84.	2.3	22
12	Levels of phosphorylated axonal neurofilament subunit H (pNfH) are increased in acute ischemic stroke. Journal of the Neurological Sciences, 2011, 304, 117-121.	0.6	58
13	The effect of ageing on human lymphocyte subsets: comparison of males and females. Immunity and Ageing, 2010, 7, 4.	4.2	133
14	Immune activation in the peripheral blood of patients with acute ischemic stroke. Journal of Neuroimmunology, 2009, 206, 112-117.	2.3	98
15	The Linker Domain of the Ha-Ras Hypervariable Region Regulates Interactions with Exchange Factors, Raf-1 and Phosphoinositide 3-Kinase. Journal of Biological Chemistry, 2002, 277, 272-278.	3.4	76
16	GTP-dependent segregation of H-ras from lipid rafts is required for biological activity. Nature Cell Biology, 2001, 3, 368-375.	10.3	492
17	Escape from Apoptosis after Prolonged Serum Deprivation Is Associated with the Regulation of the Mitochondrial Death Pathway by Bcl Biochemical and Biophysical Research Communications, 2000, 277, 487-493.	2.1	18
18	Ras Isoforms Vary in Their Ability to Activate Raf-1 and Phosphoinositide 3-Kinase. Journal of Biological Chemistry, 1998, 273, 24052-24056.	3.4	393

Jun Yan

#	Article	lF	CITATIONS
19	14-3-3 Facilitates Ras-Dependent Raf-1 Activation In Vitro and In Vivo. Molecular and Cellular Biology, 1998, 18, 3947-3955.	2.3	124
20	Activity of Plasma Membrane-recruited Raf-1 Is Regulated by Ras via the Raf Zinc Finger. Journal of Biological Chemistry, 1997, 272, 20139-20145.	3.4	97
21	Defective Signaling through the B Cell Antigen Receptor in Epstein-Barr Virus-transformed Ataxia-Telangiectasia Cells. Journal of Biological Chemistry, 1997, 272, 9489-9495.	3.4	38
22	rRNA genes from the lower chordateHerdmania momus: structural similarity with higher eukaryotes. Nucleic Acids Research, 1990, 18, 7063-7070.	14.5	25