Dorte Herzke

List of Publications by Citations

Source: https://exaly.com/author-pdf/2212687/dorte-herzke-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

127
papers6,738
citations39
h-index79
g-index136
ext. papers8,003
ext. citations7.6
avg, IF5.95
L-index

#	Paper	IF	Citations
127	Novel brominated flame retardants: a review of their analysis, environmental fate and behaviour. <i>Environment International</i> , 2011 , 37, 532-56	12.9	1030
126	Levels and trends of brominated flame retardants in the European environment. <i>Chemosphere</i> , 2006 , 64, 187-208	8.4	641
125	Brominated flame retardants in the Arctic environmenttrends and new candidates. <i>Science of the Total Environment</i> , 2010 , 408, 2885-918	10.2	570
124	Levels and trends of PBDEs and HBCDs in the global environment: status at the end of 2012. <i>Environment International</i> , 2014 , 65, 147-58	12.9	304
123	Levels and trends of HBCD and BDEs in the European and Asian environments, with some information for other BFRs. <i>Chemosphere</i> , 2008 , 73, 223-41	8.4	213
122	Brominated flame retardants in the European chemicals policy of REACH-Regulation and determination in materials. <i>Journal of Chromatography A</i> , 2009 , 1216, 320-33	4.5	169
121	Perfluoroalkyl and polyfluoroalkyl substances (PFASs) in consumer products in Norway - a pilot study. <i>Chemosphere</i> , 2012 , 88, 980-7	8.4	159
120	Negligible Impact of Ingested Microplastics on Tissue Concentrations of Persistent Organic Pollutants in Northern Fulmars off Coastal Norway. <i>Environmental Science & Environmental &</i>	10.3	157
119	An overview of the uses of per- and polyfluoroalkyl substances (PFAS). <i>Environmental Sciences: Processes and Impacts</i> , 2020 , 22, 2345-2373	4.3	146
118	Levels, Isomer Profiles, and Estimated Riverine Mass Discharges of Perfluoroalkyl Acids and Fluorinated Alternatives at the Mouths of Chinese Rivers. <i>Environmental Science & Environmental Science &</i>	10.3	138
117	Tracking pan-continental trends in environmental contamination using sentinel raptors-what types of samples should we use?. <i>Ecotoxicology</i> , 2016 , 25, 777-801	2.9	102
116	Two trace analytical methods for determination of hydroxylated PCBs and other halogenated phenolic compounds in eggs from Norwegian birds of prey. <i>Analytical Chemistry</i> , 2004 , 76, 441-52	7.8	90
115	The structure of the fire fighting foam surfactant Forafac 1157 and its biological and photolytic transformation products. <i>Chemosphere</i> , 2012 , 89, 869-75	8.4	87
114	Perfluorinated and other persistent halogenated organic compounds in European shag (Phalacrocorax aristotelis) and common eider (Somateria mollissima) from Norway: a suburban to remote pollutant gradient. <i>Science of the Total Environment</i> , 2009 , 408, 340-8	10.2	79
113	The concept of essential use for determining when uses of PFASs can be phased out. <i>Environmental Sciences: Processes and Impacts</i> , 2019 , 21, 1803-1815	4.3	71
112	What is the effect of phasing out long-chain per- and polyfluoroalkyl substances on the concentrations of perfluoroalkyl acids and their precursors in the environment? A systematic review. <i>Environmental Evidence</i> , 2018 , 7,	3.3	71
111	A first evaluation of the usefulness of feathers of nestling predatory birds for non-destructive biomonitoring of persistent organic pollutants. <i>Environment International</i> , 2011 , 37, 622-30	12.9	67

(2016-2003)

110	Polychlorinated camphenes (toxaphenes), polybrominated diphenylethers and other halogenated organic pollutants in glaucous gull (Larus hyperboreus) from Svalbard and BjĒnŪa (Bear Island). <i>Environmental Pollution</i> , 2003 , 121, 293-300	9.3	64
109	Levels and trends of poly- and perfluoroalkyl substances in the Arctic environment IAn update. <i>Emerging Contaminants</i> , 2019 , 5, 240-271	5.8	63
108	Perfluorinated alkylated substances in vegetables collected in four European countries; occurrence and human exposure estimations. <i>Environmental Science and Pollution Research</i> , 2013 , 20, 7930-9	5.1	62
107	Organochlorines in egg samples from Norwegian birds of prey: congener-, isomer- and enantiomer specific considerations. <i>Science of the Total Environment</i> , 2002 , 291, 59-71	10.2	62
106	Seasonality in contaminant accumulation in Arctic marine pelagic food webs using trophic magnification factor as a measure of bioaccumulation. <i>Environmental Toxicology and Chemistry</i> , 2011 , 30, 1026-35	3.8	59
105	Zfich Statement on Future Actions on Per- and Polyfluoroalkyl Substances (PFASs). <i>Environmental Health Perspectives</i> , 2018 , 126, 84502	8.4	58
104	Survival rate and breeding outputs in a high Arctic seabird exposed to legacy persistent organic pollutants and mercury. <i>Environmental Pollution</i> , 2015 , 200, 1-9	9.3	55
103	Temporal trends and pattern of polyfluoroalkyl compounds in Tawny Owl (Strix aluco) eggs from Norway, 1986-2009. <i>Environmental Science & Environmental Science & Environmenta</i>	10.3	54
102	Plastic litter in the European Arctic: What do we know?. Emerging Contaminants, 2019, 5, 308-318	5.8	54
101	Occurrence of perfluoroalkyl substances (PFASs) in various food items of animal origin collected in four European countries. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment,</i> 2013 , 30, 1918-32	3.2	53
100	Endocrine and fitness correlates of long-chain perfluorinated carboxylates exposure in Arctic breeding black-legged kittiwakes. <i>Environmental Science & Environmental Science</i>	10.3	52
99	Strategies for grouping per- and polyfluoroalkyl substances (PFAS) to protect human and environmental health. <i>Environmental Sciences: Processes and Impacts</i> , 2020 , 22, 1444-1460	4.3	51
98	Strongly increasing blood concentrations of lipid-soluble organochlorines in high arctic common eiders during incubation fast. <i>Chemosphere</i> , 2010 , 79, 320-5	8.4	49
97	Organohalogenated contaminants in white-tailed eagle (Haliaeetus albicilla) nestlings: An assessment of relationships to immunoglobulin levels, telomeres and oxidative stress. <i>Science of the Total Environment</i> , 2016 , 539, 337-349	10.2	47
96	BFR-governmental testing programme. Environment International, 2003, 29, 781-92	12.9	47
95	Dietary exposure to selected perfluoroalkyl acids (PFAAs) in four European regions. <i>Food Additives</i> and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment, 2013 , 30, 2141-51	3.2	46
94	Relationships between organohalogen contaminants and blood plasma clinical-chemical parameters in chicks of three raptor species from Northern Norway. <i>Ecotoxicology and Environmental Safety</i> , 2010 , 73, 7-17	7	45
93	Exposure to oxychlordane is associated with shorter telomeres in arctic breeding kittiwakes. <i>Science of the Total Environment</i> , 2016 , 563-564, 125-30	10.2	42

92	Perfluorinated substances and telomeres in an Arctic seabird: Cross-sectional and longitudinal approaches. <i>Environmental Pollution</i> , 2017 , 230, 360-367	9.3	42
91	Perfluorinated and chlorinated pollutants as predictors of demographic parameters in an endangered seabird. <i>Environmental Pollution</i> , 2008 , 156, 417-24	9.3	42
90	Blood plasma clinical-chemical parameters as biomarker endpoints for organohalogen contaminant exposure in Norwegian raptor nestlings. <i>Ecotoxicology and Environmental Safety</i> , 2012 , 80, 76-83	7	40
89	Differences between Arctic and Atlantic fjord systems on bioaccumulation of persistent organic pollutants in zooplankton from Svalbard. <i>Science of the Total Environment</i> , 2011 , 409, 2783-95	10.2	40
88	Bird feathers as a biomonitor for environmental pollutants: Prospects and pitfalls. <i>TrAC - Trends in Analytical Chemistry</i> , 2019 , 118, 223-226	14.6	39
87	Are imported consumer products an important diffuse source of PFASs to the Norwegian environment?. <i>Environmental Pollution</i> , 2015 , 198, 223-30	9.3	38
86	Influence of season, location, and feeding strategy on bioaccumulation of halogenated organic contaminants in Arctic marine zooplankton. <i>Environmental Toxicology and Chemistry</i> , 2011 , 30, 77-87	3.8	38
85	Perfluoroalkyl substances in soft tissues and tail feathers of Belgian barn owls (Tyto alba) using statistical methods for left-censored data to handle non-detects. <i>Environment International</i> , 2013 , 52, 9-16	12.9	37
84	The high persistence of PFAS is sufficient for their management as a chemical class. <i>Environmental Sciences: Processes and Impacts</i> , 2020 , 22, 2307-2312	4.3	37
83	Occurrence of perfluorinated alkylated substances in cereals, salt, sweets and fruit items collected in four European countries. <i>Chemosphere</i> , 2015 , 129, 179-85	8.4	36
82	The search for alternative aqueous film forming foams (AFFF) with a low environmental impact: physiological and transcriptomic effects of two Forafac(()) fluorosurfactants in turbot. <i>Aquatic Toxicology</i> , 2011 , 104, 168-76	5.1	36
81	Natural and man-made organobromine compounds in marine biota from Central Norway. <i>Environment International</i> , 2007 , 33, 17-26	12.9	36
80	Are Fluoropolymers Really of Low Concern for Human and Environmental Health and Separate from Other PFAS?. <i>Environmental Science & Environmental Scie</i>	10.3	35
79	Developmental Toxicity of Perfluorooctanesulfonate (PFOS) and Its Chlorinated Polyfluoroalkyl Ether Sulfonate Alternative F-53B in the Domestic Chicken. <i>Environmental Science & Emp; Technology</i> , 2018 , 52, 12859-12867	10.3	35
78	What is the effect of phasing out long-chain per- and polyfluoroalkyl substances on the concentrations of perfluoroalkyl acids and their precursors in the environment? A systematic review protocol. <i>Environmental Evidence</i> , 2015 , 4, 3	3.3	34
77	Perfluorinated, brominated, and chlorinated contaminants in a population of lesser black-backed gulls (Larus fuscus). <i>Environmental Toxicology and Chemistry</i> , 2008 , 27, 1383	3.8	34
76	Higher plasma oxidative damage and lower plasma antioxidant defences in an Arctic seabird exposed to longer perfluoroalkyl acids. <i>Environmental Research</i> , 2019 , 168, 278-285	7.9	33
75	Contaminants and energy expenditure in an Arctic seabird: Organochlorine pesticides and perfluoroalkyl substances are associated with metabolic rate in a contrasted manner. <i>Environmental Research</i> , 2017 , 157, 118-126	7.9	31

(2010-2016)

74	Spatial and temporal trends in perfluoroalkyl substances (PFASs) in ringed seals (Pusa hispida) from Svalbard. <i>Environmental Pollution</i> , 2016 , 214, 230-238	9.3	31
73	Effect of body condition on tissue distribution of perfluoroalkyl substances (PFASs) in Arctic fox (Vulpes lagopus). <i>Environmental Science & Environmental & Environm</i>	10.3	29
72	The stress of being contaminated? Adrenocortical function and reproduction in relation to persistent organic pollutants in female black legged kittiwakes. <i>Science of the Total Environment</i> , 2014 , 476-477, 553-60	10.2	29
71	Geographical Differences in Dietary Exposure to Perfluoroalkyl Acids between Manufacturing and Application Regions in China. <i>Environmental Science & Environmental Science & </i>	10.3	28
70	Relationships between POPs and baseline corticosterone levels in black-legged kittiwakes (Rissa tridactyla) across their breeding cycle. <i>Environmental Pollution</i> , 2012 , 164, 219-26	9.3	28
69	White-Tailed Eagle () Body Feathers Document Spatiotemporal Trends of Perfluoroalkyl Substances in the Northern Environment. <i>Environmental Science & Environmental Science & </i>	10.3	27
68	Temporal dynamics of circulating persistent organic pollutants in a fasting seabird under different environmental conditions. <i>Environmental Science & Environmental Science &</i>	10.3	27
67	Environmental pollutants in the Swedish marine ecosystem, with special emphasis on polybrominated diphenyl ethers (PBDE). <i>Chemosphere</i> , 2011 , 82, 1286-92	8.4	27
66	Fluctuating wing asymmetry and hepatic concentrations of persistent organic pollutants are associated in European shag (Phalacrocorax aristotelis) chicks. <i>Science of the Total Environment</i> , 2010 , 408, 578-85	10.2	26
65	Exposure to per- and polyfluoroalkyl substances through the consumption of fish from lakes affected by aqueous film-forming foam emissions - A combined epidemiological and exposure modeling approach. The SAMINOR 2 Clinical Study. <i>Environment International</i> , 2016 , 94, 272-282	12.9	25
64	Temporal trends and spatial differences of perfluoroalkylated substances in livers of harbor porpoise (Phocoena phocoena) populations from Northern Europe, 1991-2008. <i>Science of the Total Environment</i> , 2012 , 419, 216-24	10.2	25
63	Car Tire Crumb Rubber: Does Leaching Produce a Toxic Chemical Cocktail in Coastal Marine Systems?. <i>Frontiers in Environmental Science</i> , 2020 , 8,	4.8	25
62	White-tailed eagle (Haliaeetus albicilla) feathers from Norway are suitable for monitoring of legacy, but not emerging contaminants. <i>Science of the Total Environment</i> , 2019 , 647, 525-533	10.2	24
61	Ecological and spatial factors drive intra- and interspecific variation in exposure of subarctic predatory bird nestlings to persistent organic pollutants. <i>Environment International</i> , 2013 , 57-58, 25-33	12.9	24
60	Plasma concentrations of organohalogenated pollutants in predatory bird nestlings: associations to growth rate and dietary tracers. <i>Environmental Toxicology and Chemistry</i> , 2013 , 32, 2520-7	3.8	24
59	The BEEP Stavanger Workshop: Mesocosm exposures. <i>Aquatic Toxicology</i> , 2006 , 78 Suppl 1, S5-12	5.1	24
58	Polychlorinated biphenyls (PCBs), polybrominated diphenyl ethers (PBDEs) and perfluorinated alkylated substances (PFASs) in traditional seafood items from western Greenland. <i>Environmental Science and Pollution Research</i> , 2014 , 21, 4741-50	5.1	23
57	Salmon farms as a source of organohalogenated contaminants in wild fish. <i>Environmental Science & Environmental & Environmenta</i>	10.3	22

56	Dissimilar effects of organohalogenated compounds on thyroid hormones in glaucous gulls. <i>Environmental Research</i> , 2017 , 158, 350-357	7.9	21
55	Detailed analysis of polybrominated biphenyl congeners in bird eggs from Norway. <i>Environmental Pollution</i> , 2008 , 156, 1204-10	9.3	21
54	Plasma concentrations of organohalogenated contaminants in white-tailed eagle nestlings - The role of age and diet. <i>Environmental Pollution</i> , 2019 , 246, 527-534	9.3	21
53	Increased adrenal responsiveness and delayed hatching date in relation to polychlorinated biphenyl exposure in Arctic-breeding black-legged kittiwakes (Rissa tridactyla). <i>General and Comparative Endocrinology</i> , 2015 , 219, 165-72	3	19
52	Perfluoroalkyl substances detected in the world's southernmost marine mammal, the Weddell seal (Leptonychotes weddellii). <i>Environmental Pollution</i> , 2015 , 197, 62-67	9.3	19
51	Antiparasite treatments reduce humoral immunity and impact oxidative status in raptor nestlings. <i>Ecology and Evolution</i> , 2013 , 3, 5157-66	2.8	19
50	Estimating human exposure to perfluoroalkyl acids via solid food and drinks: Implementation and comparison of different dietary assessment methods. <i>Environmental Research</i> , 2017 , 158, 269-276	7.9	18
49	Per- and polyfluoroalkyl substances in plasma and feathers of nestling birds of prey from northern Norway. <i>Environmental Research</i> , 2017 , 158, 277-285	7.9	18
48	Persistent organic pollutants and organophosphate esters in feathers and blood plasma of adult kittiwakes (Rissa tridactyla) from Svalbard - associations with body condition and thyroid hormones. <i>Environmental Research</i> , 2018 , 164, 158-164	7.9	16
47	Impacts of climate and feeding conditions on the annual accumulation (1986-2009) of persistent organic pollutants in a terrestrial raptor. <i>Environmental Science & Environmental Science & Environmen</i>	10.3	16
46	Exposure to PFAS is Associated with Telomere Length Dynamics and Demographic Responses of an Arctic Top Predator. <i>Environmental Science & Environmental Science & Environment</i>	10.3	16
45	DNA double-strand breaks in relation to persistent organic pollutants in a fasting seabird. <i>Ecotoxicology and Environmental Safety</i> , 2014 , 106, 68-75	7	15
44	Monitoring of raptors and their contamination levels in Norway. <i>Ambio</i> , 2008 , 37, 420-4	6.5	15
43	Analytical strategies for successful enantioselective separation of atropisomeric polybrominated biphenyls 132 and 149 in environmental samples. <i>Journal of Chromatography A</i> , 2005 , 1063, 193-9	4.5	15
42	Spatial and temporal distribution of chiral pesticides in Calanus spp. from three Arctic fjords. <i>Environmental Pollution</i> , 2014 , 192, 154-61	9.3	14
41	Perfluoroalkyl substance concentrations in a terrestrial raptor: relationships to environmental conditions and individual traits. <i>Environmental Toxicology and Chemistry</i> , 2015 , 34, 184-91	3.8	14
40	Integrated exposure assessment of northern goshawk (Accipiter gentilis) nestlings to legacy and emerging organic pollutants using non-destructive samples. <i>Environmental Research</i> , 2019 , 178, 108678	7.9	13
39	Integument colouration in relation to persistent organic pollutants and body condition in arctic breeding black-legged kittiwakes (Rissa tridactyla). <i>Science of the Total Environment</i> , 2014 , 470-471, 248-	102 -54	13

(2020-2017)

38	Temporal variation in circulating concentrations of organochlorine pollutants in a pelagic seabird breeding in the high Arctic. <i>Environmental Toxicology and Chemistry</i> , 2017 , 36, 442-448	3.8	13
37	Snow buntings (Plectrophenax nivealis) as bio-indicators for exposure differences to legacy and emerging persistent organic pollutants from the Arctic terrestrial environment on Svalbard. <i>Science of the Total Environment</i> , 2019 , 667, 638-647	10.2	12
36	The NORMAN Association and the European Partnership for Chemicals Risk Assessment (PARC): let cooperate!. <i>Environmental Sciences Europe</i> , 2020 , 32,	5	12
35	Enantiomer-selective and quantitative trace analysis of selected persistent organic pollutants (POP) in traditional food from western Greenland. <i>Journal of Toxicology and Environmental Health - Part A: Current Issues</i> , 2014 , 77, 616-27	3.2	12
34	Kinetics and organotropy of some polyfluorinated dibenzo-p-dioxins and dibenzofurans (PFDD/PFDF) in rats. <i>Life Sciences</i> , 2002 , 71, 1475-86	6.8	11
33	Potential Effect of Migration Strategy on Pollutant Occurrence in Eggs of Arctic Breeding Barnacle Geese (Branta leucopsis). <i>Environmental Science & Description (Company)</i> 2019, 53, 5427-5435	10.3	10
32	Determination of the enantiomer fraction of PBB 149 by gas chromatography/electron capture negative ionization tandem mass spectrometry in the selected reaction monitoring mode. <i>Rapid Communications in Mass Spectrometry</i> , 2005 , 19, 3719-23	2.2	10
31	Persistent organic pollutant levels and the importance of source proximity in Baltic and Svalbard breeding common eiders. <i>Environmental Toxicology and Chemistry</i> , 2016 , 35, 1526-33	3.8	10
30	DNA double-strand breaks in incubating female common eiders (Somateria mollissima): Comparison between a low and a high polluted area. <i>Environmental Research</i> , 2016 , 151, 297-303	7.9	10
29	A schematic sampling protocol for contaminant monitoring in raptors. <i>Ambio</i> , 2021 , 50, 95-100	6.5	10
28	Individual variability in contaminants and physiological status in a resident Arctic seabird species. <i>Environmental Pollution</i> , 2019 , 249, 191-199	9.3	9
27	Pelagic vs Coastal-Key Drivers of Pollutant Levels in Barents Sea Polar Bears with Contrasted Space-Use Strategies. <i>Environmental Science & Environmental Science & Environme</i>	10.3	9
26	Antioxidant Responses in Relation to Persistent Organic Pollutants and Metals in a Low- and a High-Exposure Population of Seabirds. <i>Environmental Science & Environmental Sci</i>	10.3	9
25	DNA damage in Arctic seabirds: Baseline, sensitivity to a genotoxic stressor, and association with organohalogen contaminants. <i>Environmental Toxicology and Chemistry</i> , 2018 , 37, 1084-1091	3.8	9
24	Contamination of ivory gulls (Pagophila eburnea) at four colonies in Svalbard in relation to their trophic behaviour. <i>Polar Biology</i> , 2017 , 40, 917-929	2	8
23	Per- and polyfluorinated substances in the Nordic Countries		8
22	Ingested plastics in northern fulmars (Fulmarus glacialis): A pathway for polybrominated diphenyl ether (PBDE) exposure?. <i>Science of the Total Environment</i> , 2021 , 778, 146313	10.2	8
21	Spatiotemporal Analysis of Perfluoroalkyl Substances in White-Tailed Eagle (Nestlings from	10.3	

20	Contaminants, prolactin and parental care in an Arctic seabird: Contrasted associations of perfluoroalkyl substances and organochlorine compounds with egg-turning behavior. <i>General and Comparative Endocrinology</i> , 2020 , 291, 113420	3	7
19	Oceanic long-range transport of organic additives present in plastic products: an overview. <i>Environmental Sciences Europe</i> , 2021 , 33,	5	7
18	Organochlorines, perfluoroalkyl substances, mercury, and egg incubation temperature in an Arctic seabird: Insights from data loggers. <i>Environmental Toxicology and Chemistry</i> , 2018 , 37, 2881-2894	3.8	5
17	Schadstoff-Ferntransport in die Arktis. <i>Environmental Sciences Europe</i> , 2001 , 13, 216-226		5
16	A Bad Start in Life? Maternal Transfer of Legacy and Emerging Poly- and Perfluoroalkyl Substances to Eggs in an Arctic Seabird. <i>Environmental Science & Emp; Technology</i> , 2021 ,	10.3	5
15	Seabird-Transported Contaminants Are Reflected in the Arctic Tundra, But Not in Its Soil-Dwelling Springtails (Collembola). <i>Environmental Science & Environmental Science & E</i>	10.3	4
14	Identification of toxaphene congeners in bird eggs by combining quadrupole NICI-mS and ion trap EI-MS/MS. <i>Journal of Separation Science</i> , 2002 , 25, 453-461	3.4	4
13	Moving forward in microplastic research: A Norwegian perspective. <i>Environment International</i> , 2021 , 157, 106794	12.9	4
12	Current Levels and Trends of Brominated Flame Retardants in the Environment. <i>Handbook of Environmental Chemistry</i> , 2010 , 123-140	0.8	3
11	Bioaccumulation of Brominated Flame Retardants. Handbook of Environmental Chemistry, 2010, 141-18	8 5 0.8	3
10	Maternal-Child Exposures to Persistent Organic Pollutants in Dhaka, Bangladesh. <i>Exposure and Health</i> , 2020 , 12, 79-87	8.8	3
9	Finding essentiality feasible: common questions and misinterpretations concerning the "essential-use" concept. <i>Environmental Sciences: Processes and Impacts</i> , 2021 , 23, 1079-1087	4.3	3
8	Blood clinical-chemical parameters and feeding history in growing Japanese quail (Coturnix japonica) chicks exposed to Tris(1,3-dichloro-2-propyl) phosphate and Dechlorane Plus in ovo. <i>Toxicological and Environmental Chemistry</i> , 2017 , 99, 938-952	1.4	2
7	A novel use of the leukocyte coping capacity assay to assess the immunomodulatory effects of organohalogenated contaminants in avian wildlife. <i>Environment International</i> , 2020 , 142, 105861	12.9	2
6	Trophic and fitness correlates of mercury and organochlorine compound residues in egg-laying Antarctic petrels. <i>Environmental Research</i> , 2021 , 193, 110518	7.9	2
5	Addressing Urgent Questions for PFAS in the 21st Century. <i>Environmental Science & Eamp; Technology</i> , 2021 , 55, 12755-12765	10.3	2
4	Perfluorinated, brominated, and chlorinated contaminants in a population of lesser black-backed gulls (Larus fuscus). <i>Environmental Toxicology and Chemistry</i> , 2008 , 27, 1383-92	3.8	1
3	First documentation of plastic ingestion in the arctic glaucous gull (Larus hyperboreus) <i>Science of the Total Environment</i> , 2022 , 155340	10.2	0

LIST OF PUBLICATIONS

Ecosystem specific accumulation of organohalogenated compounds: A comparison between adjacent freshwater and terrestrial avian predators.. *Environmental Research*, **2022**, 113455

7.9 0

Anti-parasite treatment and blood biochemistry in raptor nestlings. *Canadian Journal of Zoology*, **2017**, 95, 685-693

1.5