

Mitsuro Kanda

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/221960/publications.pdf>

Version: 2024-02-01

298
papers

7,927
citations

57719

44
h-index

95218

68
g-index

303
all docs

303
docs citations

303
times ranked

9289
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Presence of Somatic Mutations in Most Early-Stage Pancreatic Intraepithelial Neoplasia. <i>Gastroenterology</i> , 2012, 142, 730-733.e9. | 0.6 | 568 |
| 2 | Mutant <i>GNAS</i> detected in duodenal collections of secretin-stimulated pancreatic juice indicates the presence or emergence of pancreatic cysts. <i>Gut</i> , 2013, 62, 1024-1033. | 6.1 | 160 |
| 3 | Mutant TP53 in Duodenal Samples of Pancreatic Juice From Patients With Pancreatic Cancer or High-Grade Dysplasia. <i>Clinical Gastroenterology and Hepatology</i> , 2013, 11, 719-730.e5. | 2.4 | 154 |
| 4 | Modified Blumgart Anastomosis for Pancreaticojejunostomy: Technical Improvement in Matched Historical Control Study. <i>Journal of Gastrointestinal Surgery</i> , 2014, 18, 1108-1115. | 0.9 | 145 |
| 5 | Molecular mechanisms of peritoneal dissemination in gastric cancer. <i>World Journal of Gastroenterology</i> , 2016, 22, 6829. | 1.4 | 121 |
| 6 | Nutritional predictors for postoperative short-term and long-term outcomes of patients with gastric cancer. <i>Medicine (United States)</i> , 2016, 95, e3781. | 0.4 | 105 |
| 7 | Effectiveness of plasma treatment on pancreatic cancer cells. <i>International Journal of Oncology</i> , 2015, 47, 1655-1662. | 1.4 | 98 |
| 8 | Recent advances in the molecular diagnostics of gastric cancer. <i>World Journal of Gastroenterology</i> , 2015, 21, 9838. | 1.4 | 92 |
| 9 | Clinical Implication of Inflammation-Based Prognostic Score in Pancreatic Cancer. <i>Medicine (United States)</i> 107:1431-1439 (2014) | 0.4 | 90 |
| 10 | Pattern of Lymph Node Metastasis Spread in Pancreatic Cancer. <i>Pancreas</i> , 2011, 40, 951-955. | 0.5 | 89 |
| 11 | Promoter hypermethylation of fibulin 1 gene is associated with tumor progression in hepatocellular carcinoma. <i>Molecular Carcinogenesis</i> , 2011, 50, 571-579. | 1.3 | 86 |
| 12 | Effectiveness of plasma treatment on gastric cancer cells. <i>Gastric Cancer</i> , 2015, 18, 635-643. | 2.7 | 83 |
| 13 | Estrogen receptor 1 gene as a tumor suppressor gene in hepatocellular carcinoma detected by triple-combination array analysis. <i>International Journal of Oncology</i> , 2013, 43, 88-94. | 1.4 | 81 |
| 14 | Significance of SYT8 For the Detection, Prediction, and Treatment of Peritoneal Metastasis From Gastric Cancer. <i>Annals of Surgery</i> , 2018, 267, 495-503. | 2.1 | 81 |
| 15 | KRAS and Guanine Nucleotide-Binding Protein Mutations in Pancreatic Juice Collected From the Duodenum of Patients at High Risk for Neoplasia Undergoing Endoscopic Ultrasound. <i>Clinical Gastroenterology and Hepatology</i> , 2015, 13, 963-969.e4. | 2.4 | 74 |
| 16 | Clinical impact of sarcopenia on prognosis in pancreatic ductal adenocarcinoma: A retrospective cohort study. <i>International Journal of Surgery</i> , 2017, 39, 45-51. | 1.1 | 74 |
| 17 | Intraperitoneal Administration of Plasma-Activated Medium: Proposal of a Novel Treatment Option for Peritoneal Metastasis From Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2017, 24, 1188-1194. | 0.7 | 74 |
| 18 | Comparison of inflammation-based prognostic scores as predictors of tumor recurrence in patients with hepatocellular carcinoma after curative resection. <i>Journal of Hepato-Biliary-Pancreatic Sciences</i> , 2014, 21, 682-688. | 1.4 | 72 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Prognostic impact of pancreatic margin status in the intraductal papillary mucinous neoplasms of the pancreas. <i>Surgery</i> , 2010, 148, 285-290. | 1.0 | 71 |
| 20 | Identification of the collagen type 1 alpha 1 gene (COL1A1) as a candidate survival-related factor associated with hepatocellular carcinoma. <i>BMC Cancer</i> , 2014, 14, 108. | 1.1 | 71 |
| 21 | Adverse prognostic impact of perioperative allogeneic transfusion on patients with stage II/III gastric cancer. <i>Gastric Cancer</i> , 2016, 19, 255-263. | 2.7 | 70 |
| 22 | Epidermal Growth Factor-Containing Fibulin-Like Extracellular Matrix Protein 1, EFEMP1, a Novel Tumor-Suppressor Gene Detected in Hepatocellular Carcinoma Using Double Combination Array Analysis. <i>Annals of Surgical Oncology</i> , 2010, 17, 923-932. | 0.7 | 69 |
| 23 | Genetic and epigenetic aspects of initiation and progression of hepatocellular carcinoma. <i>World Journal of Gastroenterology</i> , 2015, 21, 10584. | 1.4 | 66 |
| 24 | Invasion of the Splenic Artery Is a Crucial Prognostic Factor in Carcinoma of the Body and Tail of the Pancreas. <i>Annals of Surgery</i> , 2010, 251, 483-487. | 2.1 | 65 |
| 25 | Impact of Operative Blood Loss on Survival in Invasive Ductal Adenocarcinoma of the Pancreas. <i>Pancreas</i> , 2011, 40, 3-9. | 0.5 | 63 |
| 26 | The Controlling Nutritional Status Score Serves as a Predictor of Short- and Long-Term Outcomes for Patients with Stage 2 or 3 Gastric Cancer: Analysis of a Multi-institutional Data Set. <i>Annals of Surgical Oncology</i> , 2019, 26, 456-464. | 0.7 | 61 |
| 27 | Correlations of the expression of vascular endothelial growth factor B and its isoforms in hepatocellular carcinoma with clinicopathological parameters. <i>Journal of Surgical Oncology</i> , 2008, 98, 190-196. | 0.8 | 59 |
| 28 | Preservation of the Pyloric Ring Has Little Value in Surgery for Pancreatic Head Cancer: A Comparative Study Comparing Three Surgical Procedures. <i>Annals of Surgical Oncology</i> , 2012, 19, 176-183. | 0.7 | 58 |
| 29 | Preoperative Internal Biliary Drainage Increases the Risk of Bile Juice Infection and Pancreatic Fistula After Pancreatoduodenectomy. <i>Pancreas</i> , 2015, 44, 465-470. | 0.5 | 58 |
| 30 | Comprehensive Genomic Profiling of Neuroendocrine Carcinomas of the Gastrointestinal System. <i>Cancer Discovery</i> , 2022, 12, 692-711. | 7.7 | 58 |
| 31 | SMAD4 Expression Predicts Local Spread and Treatment Failure in Resected Pancreatic Cancer. <i>Pancreas</i> , 2015, 44, 660-664. | 0.5 | 57 |
| 32 | Recurrence Pattern and Prognosis of Pancreatic Cancer After Pancreatic Fistula. <i>Annals of Surgical Oncology</i> , 2011, 18, 2329-2337. | 0.7 | 56 |
| 33 | Inverse Probability of Treatment Weighting Analysis of Upfront Surgery Versus Neoadjuvant Chemoradiotherapy Followed by Surgery for Pancreatic Adenocarcinoma with Arterial Abutment. <i>Medicine (United States)</i> , 2015, 94, e1647. | 0.4 | 55 |
| 34 | Function and diagnostic value of Anosmin in gastric cancer progression. <i>International Journal of Cancer</i> , 2016, 138, 721-730. | 2.3 | 55 |
| 35 | SYT7 acts as a driver of hepatic metastasis formation of gastric cancer cells. <i>Oncogene</i> , 2018, 37, 5355-5366. | 2.6 | 55 |
| 36 | Adverse impact of low skeletal muscle index on the prognosis of hepatocellular carcinoma after hepatic resection. <i>International Journal of Surgery</i> , 2016, 30, 136-142. | 1.1 | 54 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Detection of metallothionein 1G as a methylated tumor suppressor gene in human hepatocellular carcinoma using a novel method of double combination array analysis. <i>International Journal of Oncology</i> , 2009, 35, 477-83. | 1.4 | 53 |
| 38 | Postoperative adjuvant chemotherapy with S-1 alters recurrence patterns and prognostic factors among patients with stage II/III gastric cancer: A propensity score matching analysis. <i>Surgery</i> , 2015, 158, 1573-1580. | 1.0 | 53 |
| 39 | Comparison of the international consensus guidelines for predicting malignancy in intraductal papillary mucinous neoplasms. <i>Surgery</i> , 2016, 159, 878-884. | 1.0 | 53 |
| 40 | Epithelial to mesenchymal transition correlates with tumor budding and predicts prognosis in esophageal squamous cell carcinoma. <i>Journal of Surgical Oncology</i> , 2014, 110, 764-769. | 0.8 | 51 |
| 41 | Vein resections >3Âcm during pancreatectomy are associated withÂpoor 1-year patency rates. <i>Surgery</i> , 2015, 157, 708-715. | 1.0 | 51 |
| 42 | Clinical benefits of neoadjuvant chemoradiotherapy for adenocarcinoma of the pancreatic head: an observational study using inverse probability of treatment weighting. <i>Journal of Gastroenterology</i> , 2017, 52, 81-93. | 2.3 | 51 |
| 43 | Clinical Implications of Naples Prognostic Score in Patients with Resected Pancreatic Cancer. <i>Annals of Surgical Oncology</i> , 2020, 27, 887-895. | 0.7 | 50 |
| 44 | Leukemia inhibitory factor receptor (LIFR) is detected as a novel suppressor gene of hepatocellular carcinoma using double-combination array. <i>Cancer Letters</i> , 2010, 289, 170-177. | 3.2 | 49 |
| 45 | Reduced Expression of Reelin (RELN) Gene Is Associated With High Recurrence Rate of Hepatocellular Carcinoma. <i>Annals of Surgical Oncology</i> , 2011, 18, 572-579. | 0.7 | 49 |
| 46 | Dihydropyrimidinase-like 3 facilitates malignant behavior of gastric cancer. <i>Journal of Experimental and Clinical Cancer Research</i> , 2014, 33, 66. | 3.5 | 49 |
| 47 | Preoperative predictors of postoperative complications after gastric cancer resection. <i>Surgery Today</i> , 2020, 50, 3-11. | 0.7 | 48 |
| 48 | Therapeutic monoclonal antibody targeting of neuronal pentraxin receptor to control metastasis in gastric cancer. <i>Molecular Cancer</i> , 2020, 19, 131. | 7.9 | 48 |
| 49 | Anti-thyroid antibodies and thyroid echo pattern at baseline as risk factors for thyroid dysfunction induced by anti-programmed cell death-1 antibodies: a prospective study. <i>British Journal of Cancer</i> , 2020, 122, 771-777. | 2.9 | 48 |
| 50 | Updated evidence on adjuvant treatments for gastric cancer. <i>Expert Review of Gastroenterology and Hepatology</i> , 2015, 9, 1549-1560. | 1.4 | 47 |
| 51 | Proposal of the Coagulation Score as a Predictor for Short-Term and Long-Term Outcomes of Patients with Resectable Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2017, 24, 502-509. | 0.7 | 46 |
| 52 | Metastatic pathway-specific transcriptome analysis identifies <i>MFSD4</i> as a putative tumor suppressor and biomarker for hepatic metastasis in patients with gastric cancer. <i>Oncotarget</i> , 2016, 7, 13667-13679. | 0.8 | 46 |
| 53 | The impact of dose/time modification in irinotecan- and oxaliplatin-based chemotherapies on outcomes in metastatic colorectal cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2014, 73, 847-855. | 1.1 | 45 |
| 54 | Prognostic impact of expression and methylation status of DENN/MADD domain-containing protein 2D in gastric cancer. <i>Gastric Cancer</i> , 2015, 18, 288-296. | 2.7 | 45 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Synaptotagmin XIII expression and peritoneal metastasis in gastric cancer. <i>British Journal of Surgery</i> , 2018, 105, 1349-1358. | 0.1 | 44 |
| 56 | Genome-Wide Somatic Copy Number Alterations in Low-Grade PanINs and IPMNs from Individuals with a Family History of Pancreatic Cancer. <i>Clinical Cancer Research</i> , 2012, 18, 4303-4312. | 3.2 | 43 |
| 57 | Adverse Effects of Intraoperative Blood Loss on Long-Term Outcomes after Curative Gastrectomy of Patients with Stage II/III Gastric Cancer. <i>Digestive Surgery</i> , 2016, 33, 121-128. | 0.6 | 43 |
| 58 | Clinical utility of the platelet-lymphocyte ratio as a predictor of postoperative complications after radical gastrectomy for clinical T2-4 gastric cancer. <i>World Journal of Gastroenterology</i> , 2017, 23, 2519. | 1.4 | 43 |
| 59 | Diagnostic efficacy of circular RNAs as noninvasive, liquid biopsy biomarkers for early detection of gastric cancer. <i>Molecular Cancer</i> , 2022, 21, 42. | 7.9 | 43 |
| 60 | Epithelial to mesenchymal transition might be induced via CD44 isoform switching in colorectal cancer. <i>Journal of Surgical Oncology</i> , 2014, 110, 745-751. | 0.8 | 42 |
| 61 | Clinical Implication of Morphological Subtypes in Management of Intraductal Papillary Mucinous Neoplasm. <i>Annals of Surgical Oncology</i> , 2014, 21, 2444-2452. | 0.7 | 41 |
| 62 | The combination of the serum carbohydrate antigen 19-9 and carcinoembryonic antigen is a simple and accurate predictor of mortality in pancreatic cancer patients. <i>Surgery Today</i> , 2014, 44, 1692-1701. | 0.7 | 41 |
| 63 | Downregulation of DENND2D by promoter hypermethylation is associated with early recurrence of hepatocellular carcinoma. <i>International Journal of Oncology</i> , 2014, 44, 44-52. | 1.4 | 41 |
| 64 | The Expression of Melanoma-Associated Antigen D2 Both in Surgically Resected and Serum Samples Serves as Clinically Relevant Biomarker of Gastric Cancer Progression. <i>Annals of Surgical Oncology</i> , 2016, 23, 214-221. | 0.7 | 41 |
| 65 | Novel diagnostics for aggravating pancreatic fistulas at the acute phase after pancreatectomy. <i>World Journal of Gastroenterology</i> , 2014, 20, 8535. | 1.4 | 41 |
| 66 | Diversity of Clinical Implication of B-Cell Translocation Gene 1 Expression by Histopathologic and Anatomic Subtypes of Gastric Cancer. <i>Digestive Diseases and Sciences</i> , 2015, 60, 1256-1264. | 1.1 | 40 |
| 67 | Protein arginine methyltransferase 5 is associated with malignant phenotype and peritoneal metastasis in gastric cancer. <i>International Journal of Oncology</i> , 2016, 49, 1195-1202. | 1.4 | 40 |
| 68 | Preoperative Albumin-Bilirubin Grade Predicts Recurrences After Radical Gastrectomy in Patients with pT2-4 Gastric Cancer. <i>World Journal of Surgery</i> , 2018, 42, 773-781. | 0.8 | 40 |
| 69 | B-cell translocation gene 1 serves as a novel prognostic indicator of hepatocellular carcinoma. <i>International Journal of Oncology</i> , 2015, 46, 641-648. | 1.4 | 39 |
| 70 | Epigenetic suppression of the immunoregulator MZB1 is associated with the malignant phenotype of gastric cancer. <i>International Journal of Cancer</i> , 2016, 139, 2290-2298. | 2.3 | 39 |
| 71 | FAM46C Serves as a Predictor of Hepatic Recurrence in Patients with Resectable Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2017, 24, 3438-3445. | 0.7 | 39 |
| 72 | Delay in initiation of postoperative adjuvant chemotherapy with S-1 monotherapy and prognosis for gastric cancer patients: analysis of a multi-institutional dataset. <i>Gastric Cancer</i> , 2019, 22, 1215-1225. | 2.7 | 39 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Estimated pancreatic parenchymal remnant volume accurately predicts clinically relevant pancreatic fistula after pancreatoduodenectomy. <i>Surgery</i> , 2014, 156, 601-610. | 1.0 | 38 |
| 74 | Tumor Infiltrative Pattern Predicts Sites of Recurrence After Curative Gastrectomy for Stages 2 and 3 Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2016, 23, 1934-1940. | 0.7 | 38 |
| 75 | Review of recent efforts to discover biomarkers for early detection, monitoring, prognosis, and prediction of treatment responses of patients with gastric cancer. <i>Expert Review of Gastroenterology and Hepatology</i> , 2018, 12, 657-670. | 1.4 | 38 |
| 76 | Review of recent molecular landscape knowledge of gastric cancer. <i>Histology and Histopathology</i> , 2018, 33, 11-26. | 0.5 | 38 |
| 77 | Long-lasting discussion: Adverse effects of intraoperative blood loss and allogeneic transfusion on prognosis of patients with gastric cancer. <i>World Journal of Gastroenterology</i> , 2019, 25, 2743-2751. | 1.4 | 38 |
| 78 | Impact of the Controlling Nutritional Status Score on the Prognosis After Curative Resection of Pancreatic Ductal Adenocarcinoma. <i>Pancreas</i> , 2018, 47, 823-829. | 0.5 | 36 |
| 79 | The Preoperative Prognostic Nutritional Index Predicts Short-Term and Long-Term Outcomes of Patients with Stage II/III Gastric Cancer: Analysis of a Multi-Institution Dataset. <i>Digestive Surgery</i> , 2020, 37, 135-144. | 0.6 | 36 |
| 80 | Prognostic Implications of Lymph Node Metastases in Carcinoma of the Body and Tail of the Pancreas. <i>Pancreas</i> , 2011, 40, 1029-1033. | 0.5 | 35 |
| 81 | Influence of Food Intake on the Healing Process of Postoperative Pancreatic Fistula After Pancreatoduodenectomy: A Multi-institutional Randomized Controlled Trial. <i>Annals of Surgical Oncology</i> , 2015, 22, 3905-3912. | 0.7 | 34 |
| 82 | A randomized phase II multicenter trial to explore efficacy of weekly intraperitoneal in comparison with intravenous paclitaxel administered immediately after gastrectomy to the patients with high risk of peritoneal recurrence: final results of the INPACT trial. <i>Gastric Cancer</i> , 2018, 21, 1014-1023. | 2.7 | 34 |
| 83 | Dynamin 3: a new candidate tumor suppressor gene in hepatocellular carcinoma detected by triple combination array analysis. <i>OncoTargets and Therapy</i> , 2013, 6, 1417. | 1.0 | 32 |
| 84 | Troponin I2 as a Specific Biomarker for Prediction of Peritoneal Metastasis in Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2018, 25, 2083-2090. | 0.7 | 32 |
| 85 | Multi-institutional analysis of the prognostic significance of postoperative complications after curative resection for gastric cancer. <i>Cancer Medicine</i> , 2019, 8, 5194-5201. | 1.3 | 32 |
| 86 | Intraoperative Blood Loss is Associated with Shortened Postoperative Survival of Patients with Stage II/III Gastric Cancer: Analysis of a Multi-institutional Dataset. <i>World Journal of Surgery</i> , 2019, 43, 870-877. | 0.8 | 32 |
| 87 | Number of retrieved lymph nodes is an independent prognostic factor after total gastrectomy for patients with stage III gastric cancer: propensity score matching analysis of a multi-institution dataset. <i>Gastric Cancer</i> , 2019, 22, 853-863. | 2.7 | 32 |
| 88 | Dihydropyrimidinase-like 3 is a putative hepatocellular carcinoma tumor suppressor. <i>Journal of Gastroenterology</i> , 2015, 50, 590-600. | 2.3 | 31 |
| 89 | The levels of SYT13 and CEA mRNAs in peritoneal lavages predict the peritoneal recurrence of gastric cancer. <i>Gastric Cancer</i> , 2019, 22, 1143-1152. | 2.7 | 31 |
| 90 | Clinical significance of expression and epigenetic profiling of TUSC1 in gastric cancer. <i>Journal of Surgical Oncology</i> , 2014, 110, 136-144. | 0.8 | 30 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Amido-Bridged Nucleic Acid-Modified Antisense Oligonucleotides Targeting SYT13 to Treat Peritoneal Metastasis of Gastric Cancer. <i>Molecular Therapy - Nucleic Acids</i> , 2020, 22, 791-802. | 2.3 | 30 |
| 92 | Significance of Preoperative Systemic Inflammation Score in Short-Term and Long-Term Outcomes of Patients with Pathological T2-4 Gastric Cancer After Radical Gastrectomy. <i>World Journal of Surgery</i> , 2018, 42, 3277-3285. | 0.8 | 29 |
| 93 | Clinical Implications of Lysyl Oxidase-Like Protein 2 Expression in Pancreatic Cancer. <i>Scientific Reports</i> , 2018, 8, 9846. | 1.6 | 29 |
| 94 | Serum levels of ANOS1 serve as a diagnostic biomarker of gastric cancer: a prospective multicenter observational study. <i>Gastric Cancer</i> , 2020, 23, 203-211. | 2.7 | 29 |
| 95 | A microRNA-based liquid biopsy signature for the early detection of esophageal squamous cell carcinoma: a retrospective, prospective and multicenter study. <i>Molecular Cancer</i> , 2022, 21, 44. | 7.9 | 29 |
| 96 | Comparison of Pancreatic Head Resection With Segmental Duodenectomy and Pylorus-Preserving Pancreatoduodenectomy for Benign and Low-Grade Malignant Neoplasms of the Pancreatic Head. <i>Pancreas</i> , 2011, 40, 1258-1263. | 0.5 | 28 |
| 97 | Mutant KRAS and GNAS DNA Concentrations in Secretin-Stimulated Pancreatic Fluid Collected from the Pancreatic Duct and the Duodenal Lumen. <i>Clinical and Translational Gastroenterology</i> , 2014, 5, e62. | 1.3 | 28 |
| 98 | Lymph node ratio as parameter of regional lymph node involvement in pancreatic cancer. <i>Langenbeck's Archives of Surgery</i> , 2016, 401, 1143-1152. | 0.8 | 28 |
| 99 | The significance of relative dose intensity in adjuvant chemotherapy of pancreatic ductal adenocarcinoma—including the analysis of clinicopathological factors influencing relative dose intensity. <i>Medicine (United States)</i> , 2016, 95, e4282. | 0.4 | 28 |
| 100 | Perioperative Serum Carcinoembryonic Antigen Levels Predict Recurrence and Survival of Patients with Pathological T2-4 Gastric Cancer Treated with Curative Gastrectomy. <i>Digestive Surgery</i> , 2018, 35, 55-63. | 0.6 | 28 |
| 101 | Decreased expression of prenyl diphosphate synthase subunit 2 correlates with reduced survival of patients with gastric cancer. <i>Journal of Experimental and Clinical Cancer Research</i> , 2014, 33, 88. | 3.5 | 27 |
| 102 | Preoperative Identification of a Prognostic Factor for Pancreatic Neuroendocrine Tumors Using Multiphase Contrast-Enhanced Computed Tomography. <i>Pancreas</i> , 2016, 45, 198-203. | 0.5 | 27 |
| 103 | Evaluation and proposal of novel resectability criteria for pancreatic cancer established by the Japan Pancreas Society. <i>Surgery</i> , 2017, 162, 784-791. | 1.0 | 27 |
| 104 | Risk Prediction of Postoperative Pneumonia After Subtotal Esophagectomy Based on Preoperative Serum Cholinesterase Concentrations. <i>Annals of Surgical Oncology</i> , 2019, 26, 3718-3726. | 0.7 | 27 |
| 105 | Pancreatic Fat and Body Composition Measurements by Computed Tomography are Associated with Pancreatic Fistula After Pancreatectomy. <i>Annals of Surgical Oncology</i> , 2021, 28, 530-538. | 0.7 | 27 |
| 106 | Significance of the Splenic Vein and Its Branches in Pancreatoduodenectomy with Resection of the Portal Vein System. <i>Digestive Surgery</i> , 2015, 32, 382-388. | 0.6 | 26 |
| 107 | NRAGE promotes the malignant phenotype of hepatocellular carcinoma. <i>Oncology Letters</i> , 2016, 11, 1847-1854. | 0.8 | 26 |
| 108 | The protein arginine methyltransferase 5 promotes malignant phenotype of hepatocellular carcinoma cells and is associated with adverse patient outcomes after curative hepatectomy. <i>International Journal of Oncology</i> , 2017, 50, 381-386. | 1.4 | 26 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Pancreatoduodenectomy With Portal Vein Resection Is Feasible and Potentially Beneficial for Elderly Patients With Pancreatic Cancer. <i>Pancreas</i> , 2014, 43, 951-958. | 0.5 | 25 |
| 110 | Reduced Expression of Adherens Junctions Associated Protein 1 Predicts Recurrence of Hepatocellular Carcinoma After Curative Hepatectomy. <i>Annals of Surgical Oncology</i> , 2015, 22, 1499-1507. | 0.7 | 25 |
| 111 | Overexpression of Derlin 3 is associated with malignant phenotype of breast cancer cells. <i>Oncology Reports</i> , 2017, 38, 1760-1766. | 1.2 | 25 |
| 112 | Reduced expression of DENND2D through promoter hypermethylation is an adverse prognostic factor in squamous cell carcinoma of the esophagus. <i>Oncology Reports</i> , 2014, 31, 693-700. | 1.2 | 24 |
| 113 | Excess Weight Adversely Influences Treatment Length of Postoperative Pancreatic Fistula. <i>Pancreas</i> , 2015, 44, 971-976. | 0.5 | 24 |
| 114 | Adherens junctions associated protein 1 serves as a predictor of recurrence of squamous cell carcinoma of the esophagus. <i>International Journal of Oncology</i> , 2015, 47, 1811-1818. | 1.4 | 24 |
| 115 | GPR155 Serves as a Predictive Biomarker for Hematogenous Metastasis in Patients with Gastric Cancer. <i>Scientific Reports</i> , 2017, 7, 42089. | 1.6 | 24 |
| 116 | Feasibility of subtotal esophagectomy with systematic lymphadenectomy in selected elderly patients with esophageal cancer; a propensity score matching analysis. <i>BMC Surgery</i> , 2019, 19, 143. | 0.6 | 24 |
| 117 | Pattern of first recurrent lesions in pancreatic cancer: hepatic relapse is associated with dismal prognosis and portal vein invasion. <i>Hepato-Gastroenterology</i> , 2014, 61, 1756-61. | 0.5 | 24 |
| 118 | Combination Treatment of Human Pancreatic Cancer Xenograft Models with the Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor Erlotinib and Oncolytic Herpes Simplex Virus HF10. <i>Annals of Surgical Oncology</i> , 2014, 21, 691-698. | 0.7 | 23 |
| 119 | Factors related to occurrence and aggravation of pancreatic fistula after radical gastrectomy for gastric cancer. <i>Journal of Surgical Oncology</i> , 2015, 112, 381-386. | 0.8 | 23 |
| 120 | Comparison of the Survival Outcomes of Pancreatic Cancer and Intraductal Papillary Mucinous Neoplasms. <i>Pancreas</i> , 2018, 47, 974-979. | 0.5 | 23 |
| 121 | Evaluation of MAGE4 expression in hepatocellular carcinoma in Japanese patients. <i>Journal of Surgical Oncology</i> , 2013, 108, 557-562. | 0.8 | 22 |
| 122 | Feeding Duodenostomy Decreases the Incidence of Mechanical Obstruction After Radical Esophageal Cancer Surgery. <i>World Journal of Surgery</i> , 2015, 39, 1105-1110. | 0.8 | 22 |
| 123 | Increased Expression of DNAJC12 is Associated with Aggressive Phenotype of Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2019, 26, 836-844. | 0.7 | 22 |
| 124 | Prognostic significance of perioperative tumor marker levels in stage II/III gastric cancer. <i>World Journal of Gastrointestinal Oncology</i> , 2019, 11, 17-27. | 0.8 | 22 |
| 125 | Long-term quality of life and nutrition status of the aboral pouch reconstruction after total gastrectomy for gastric cancer: a prospective multicenter observational study (CCOG1505). <i>Gastric Cancer</i> , 2019, 22, 607-616. | 2.7 | 21 |
| 126 | Expression Analysis of THOP1 in Background Liver, a Prognostic Predictive Factor in Hepatocellular Carcinoma, Extracted by Multiarray Analysis. <i>Annals of Surgical Oncology</i> , 2014, 21, 443-450. | 0.7 | 20 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Predictive value of drain amylase content for peripancreatic inflammatory fluid collections after laparoscopic (assisted) distal gastrectomy. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2016, 30, 4353-4362. | 1.3 | 20 |
| 128 | Identification of intragenic methylation in the TUSC1 gene as a novel prognostic marker of hepatocellular carcinoma. <i>Oncology Reports</i> , 2014, 31, 1305-1313. | 1.2 | 19 |
| 129 | The Prognostic Relevance of Subcarinal Lymph Node Dissection in Esophageal Squamous Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2016, 23, 611-618. | 0.7 | 19 |
| 130 | Usefulness of preoperative estimated glomerular filtration rate to predict complications after curative gastrectomy in patients with clinical T2â€“4 gastric cancer. <i>Gastric Cancer</i> , 2017, 20, 736-743. | 2.7 | 19 |
| 131 | FBXO50 Enhances the Malignant Behavior of Gastric Cancer Cells. <i>Annals of Surgical Oncology</i> , 2017, 24, 3771-3779. | 0.7 | 19 |
| 132 | Expression of sushi domain containing two reflects the malignant potential of gastric cancer. <i>Cancer Medicine</i> , 2018, 7, 5194-5204. | 1.3 | 19 |
| 133 | Modified Systemic Inflammation Score is Useful for Risk Stratification After Radical Resection of Squamous Cell Carcinoma of the Esophagus. <i>Annals of Surgical Oncology</i> , 2019, 26, 4773-4781. | 0.7 | 19 |
| 134 | Biological and conditional factors should be included when defining criteria for resectability for patients with pancreatic cancer. <i>Hpb</i> , 2019, 21, 1211-1218. | 0.1 | 19 |
| 135 | A functional polymorphism in the epidermal growth factor gene predicts hepatocellular carcinoma risk in Japanese hepatitis C patients. <i>OncoTargets and Therapy</i> , 2013, 6, 1805. | 1.0 | 18 |
| 136 | Aberrant expression of melanoma-associated antigen-D2 serves as a prognostic indicator of hepatocellular carcinoma outcome following curative hepatectomy. <i>Oncology Letters</i> , 2015, 9, 1201-1206. | 0.8 | 18 |
| 137 | Emerging evidence of the molecular landscape specific for hematogenous metastasis from gastric cancer. <i>World Journal of Gastrointestinal Oncology</i> , 2018, 10, 124-136. | 0.8 | 18 |
| 138 | Establishment of Peritoneal and Hepatic Metastasis Mouse Xenograft Models Using Gastric Cancer Cell Lines. <i>In Vivo</i> , 2019, 33, 1785-1792. | 0.6 | 18 |
| 139 | Fraser extracellular matrix complex subunit 1 promotes liver metastasis of gastric cancer. <i>International Journal of Cancer</i> , 2020, 146, 2865-2876. | 2.3 | 18 |
| 140 | Propensity-score-matched analysis of a multi-institutional dataset to compare postoperative complications between Billroth I and Roux-en-Y reconstructions after distal gastrectomy. <i>Gastric Cancer</i> , 2020, 23, 734-745. | 2.7 | 18 |
| 141 | Overexpression of <i>ankyrin1</i> promotes pancreatic cancer cell growth. <i>Oncotarget</i> , 2016, 7, 34977-34987. | 0.8 | 18 |
| 142 | Association of Inflammasome Components in Background Liver with Poor Prognosis After Curatively-resected Hepatocellular Carcinoma. <i>Anticancer Research</i> , 2017, 37, 293-300. | 0.5 | 18 |
| 143 | Clinical utility of PDSS2 expression to stratify patients at risk for recurrence of hepatocellular carcinoma. <i>International Journal of Oncology</i> , 2014, 45, 2005-2012. | 1.4 | 17 |
| 144 | Suppression of SAMS1 Expression is Associated with the Malignant Phenotype of Hepatocellular Carcinoma. <i>Annals of Surgical Oncology</i> , 2015, 22, 1453-1460. | 0.7 | 17 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | Neurotrophin Receptor-Interacting Melanoma Antigen-Encoding Gene Homolog is Associated with Malignant Phenotype of Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2016, 23, 532-539. | 0.7 | 17 |
| 146 | Prognostic significance of AKR1B10 gene expression in hepatocellular carcinoma and surrounding non-tumorous liver tissue. <i>Oncology Letters</i> , 2016, 12, 4821-4828. | 0.8 | 17 |
| 147 | Modified Blumgart Suturing Technique for Remnant Closure After Distal Pancreatectomy: a Propensity Score-Matched Analysis. <i>Journal of Gastrointestinal Surgery</i> , 2016, 20, 374-384. | 0.9 | 17 |
| 148 | Transcriptomic Profiling Identifies a Risk Stratification Signature for Predicting Peritoneal Recurrence and Micrometastasis in Gastric Cancer. <i>Clinical Cancer Research</i> , 2021, 27, 2292-2300. | 3.2 | 17 |
| 149 | Hepatic metastasis of gastric cancer is associated with enhanced expression of ethanolamine kinase 2 via the p53-Bcl-2 intrinsic apoptosis pathway. <i>British Journal of Cancer</i> , 2021, 124, 1449-1460. | 2.9 | 17 |
| 150 | Detection of doublecortin domain-containing 2 (DCDC2), a new candidate tumor suppressor gene of hepatocellular carcinoma, by triple combination array analysis. <i>Journal of Experimental and Clinical Cancer Research</i> , 2013, 32, 65. | 3.5 | 16 |
| 151 | Prognostic relevance of SAMSIN1 expression in gastric cancer. <i>Oncology Letters</i> , 2016, 12, 4708-4716. | 0.8 | 16 |
| 152 | Homeobox C10 Influences on the Malignant Phenotype of Gastric Cancer Cell Lines and its Elevated Expression Positively Correlates with Recurrence and Poor Survival. <i>Annals of Surgical Oncology</i> , 2019, 26, 1535-1543. | 0.7 | 16 |
| 153 | Oral Food Intake Versus Fasting on Postoperative Pancreatic Fistula After Distal Pancreatectomy. <i>Medicine (United States)</i> , 2015, 94, e2398. | 0.4 | 15 |
| 154 | Overexpression of melanoma-associated antigen D4 is an independent prognostic factor in squamous cell carcinoma of the esophagus. <i>Ecological Management and Restoration</i> , 2015, 28, 188-195. | 0.2 | 15 |
| 155 | Identification of NCCRP1 as an epigenetically regulated tumor suppressor and biomarker for malignant phenotypes of squamous cell carcinoma of the esophagus. <i>Oncology Letters</i> , 2017, 14, 4822-4828. | 0.8 | 15 |
| 156 | Downregulation of GPR155 as a prognostic factor after curative resection of hepatocellular carcinoma. <i>BMC Cancer</i> , 2017, 17, 610. | 1.1 | 15 |
| 157 | Recent advances in molecular biomarkers for patients with hepatocellular carcinoma. <i>Expert Review of Molecular Diagnostics</i> , 2019, 19, 725-738. | 1.5 | 15 |
| 158 | Level of Melanotransferrin in Tissue and Sera Serves as a Prognostic Marker of Gastric Cancer. <i>Anticancer Research</i> , 2019, 39, 6125-6133. | 0.5 | 15 |
| 159 | Detection of indocyanine green fluorescence to determine tumor location during laparoscopic gastrectomy for gastric cancer: Results of a prospective study. <i>Asian Journal of Endoscopic Surgery</i> , 2020, 13, 160-167. | 0.4 | 15 |
| 160 | CD44 single nucleotide polymorphism and isoform switching may predict gastric cancer recurrence. <i>Journal of Surgical Oncology</i> , 2015, 112, 622-628. | 0.8 | 14 |
| 161 | CCNJ detected by triple combination array analysis as a tumor-related gene of hepatocellular carcinoma. <i>International Journal of Oncology</i> , 2015, 46, 1963-1970. | 1.4 | 14 |
| 162 | S-1 plus nab-paclitaxel is a promising regimen for pancreatic cancer in a preclinical model. <i>Journal of Surgical Oncology</i> , 2016, 113, 413-419. | 0.8 | 14 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 163 | Salvage pharyngolaryngectomy with total esophagectomy following definitive chemoradiotherapy. Ecological Management and Restoration, 2016, 29, 598-602. | 0.2 | 14 |
| 164 | Molecular alterations in the carcinogenesis and progression of hepatocellular carcinoma: Tumor factors and background liver factors. Oncology Letters, 2016, 12, 3662-3668. | 0.8 | 14 |
| 165 | Pathological tumor infiltrative pattern and sites of initial recurrence in stage II/III gastric cancer: Propensity score matching analysis of a multi-institutional dataset. Cancer Medicine, 2018, 7, 6020-6029. | 1.3 | 14 |
| 166 | Preoperative six-minute walk distance as a predictor of postoperative complication in patients with esophageal cancer. Ecological Management and Restoration, 2020, 33, . | 0.2 | 14 |
| 167 | Optimized Cutoff Value of Serum Squamous Cell Carcinoma Antigen Concentration Accurately Predicts Recurrence After Curative Resection of Squamous Cell Carcinoma of the Esophagus. Annals of Surgical Oncology, 2020, 27, 1233-1240. | 0.7 | 14 |
| 168 | Emerging evidence of molecular biomarkers in hepatocellular carcinoma. Histology and Histopathology, 2018, 33, 343-355. | 0.5 | 14 |
| 169 | MZB1 expression indicates poor prognosis in estrogen receptor-positive breast cancer. Oncology Letters, 2020, 20, 1-1. | 0.8 | 14 |
| 170 | Lysosomal-associated membrane protein family member 5 promotes the metastatic potential of gastric cancer cells. Gastric Cancer, 2022, 25, 558-572. | 2.7 | 14 |
| 171 | Capecitabine and oxaliplatin combined with bevacizumab are feasible for treating selected Japanese patients at least 75 years of age with metastatic colorectal cancer. BMC Cancer, 2015, 15, 786. | 1.1 | 13 |
| 172 | The efficacy and safety of CapeOX plus bevacizumab therapy followed by capecitabine plus bevacizumab maintenance therapy in patients with metastatic colorectal cancer: a multi-center, single-arm, phase II study (CCOG-0902). BMC Cancer, 2017, 17, 243. | 1.1 | 13 |
| 173 | PRAME as a Potential Biomarker for Liver Metastasis of Gastric Cancer. Annals of Surgical Oncology, 2020, 27, 2071-2080. | 0.7 | 13 |
| 174 | G-protein subunit gamma-4 expression has potential for detection, prediction and therapeutic targeting in liver metastasis of gastric cancer. British Journal of Cancer, 2021, 125, 220-228. | 2.9 | 13 |
| 175 | Hepatectomy for hepatocellular carcinoma in patients with hemophilia. Journal of Hepato-Biliary-Pancreatic Sciences, 2014, 21, 824-828. | 1.4 | 12 |
| 176 | A resected case of symptomatic acinar cell cystadenoma of the pancreas displacing the main pancreatic duct. Surgical Case Reports, 2016, 2, 39. | 0.2 | 12 |
| 177 | Clinical Signatures of Mucinous and Poorly Differentiated Subtypes of Colorectal Adenocarcinomas by a Propensity Score Analysis of an Independent Patient Database from Three Phase III Trials. Diseases of the Colon and Rectum, 2018, 61, 461-471. | 0.7 | 12 |
| 178 | Pattern-Specific Transcriptomics Identifies <i>ASGR2</i> as a Predictor of Hematogenous Recurrence of Gastric Cancer. Molecular Cancer Research, 2018, 16, 1420-1429. | 1.5 | 12 |
| 179 | Comparison of non-invasive liver reserve and fibrosis models: Implications for surgery and prognosis for hepatocellular carcinoma. Hepatology Research, 2019, 49, 1305-1315. | 1.8 | 12 |
| 180 | Perioperative and prognostic implication of albumin-bilirubin-TNM score in Child-Pugh class A hepatocellular carcinoma. Annals of Gastroenterological Surgery, 2019, 3, 65-74. | 1.2 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 181 | Novel Prognostic Implications of DUPAN-2 in the Era of Initial Systemic Therapy for Pancreatic Cancer. <i>Annals of Surgical Oncology</i> , 2020, 27, 2081-2089. | 0.7 | 12 |
| 182 | Systemic Inflammation Score as a Predictor of Pneumonia after Radical Resection of Gastric Cancer: Analysis of a Multi-Institutional Dataset. <i>Digestive Surgery</i> , 2020, 37, 401-410. | 0.6 | 12 |
| 183 | Exploration of Exosomal Micro RNA Biomarkers Related to Epithelial-to-Mesenchymal Transition in Pancreatic Cancer. <i>Anticancer Research</i> , 2020, 40, 1843-1853. | 0.5 | 12 |
| 184 | Blockade of CHRN2 signaling with a therapeutic monoclonal antibody attenuates the aggressiveness of gastric cancer cells. <i>Oncogene</i> , 2021, 40, 5495-5504. | 2.6 | 12 |
| 185 | Optimal Preoperative Multidisciplinary Treatment in Borderline Resectable Pancreatic Cancer. <i>Cancers</i> , 2021, 13, 36. | 1.7 | 12 |
| 186 | Albumin-Bilirubin Score Predicts Tolerability to Adjuvant S-1 Monotherapy after Curative Gastrectomy. <i>Journal of Gastric Cancer</i> , 2019, 19, 183. | 0.9 | 12 |
| 187 | Translational implication of Kallmann syndrome-1 gene expression in hepatocellular carcinoma. <i>International Journal of Oncology</i> , 2015, 46, 2546-2554. | 1.4 | 11 |
| 188 | Serosal Invasion Strongly Associated With Recurrence After Curative Hepatic Resection of Hepatocellular Carcinoma. <i>Medicine (United States)</i> , 2015, 94, e602. | 0.4 | 11 |
| 189 | Pharmacokinetic dose adjustment of 5-FU in modified FOLFOX7 plus bevacizumab for metastatic colorectal cancer in Japanese patients: a-JUST phase II clinical trial. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 78, 1253-1261. | 1.1 | 11 |
| 190 | Copine β 1/25 expression predicts prognosis following curative resection of esophageal squamous cell carcinoma. <i>Oncology Reports</i> , 2018, 40, 3772-3780. | 1.2 | 11 |
| 191 | Incorporating molecular biomarkers into clinical practice for gastric cancer. <i>Expert Review of Anticancer Therapy</i> , 2019, 19, 757-771. | 1.1 | 11 |
| 192 | KCNJ15 Expression and Malignant Behavior of Esophageal Squamous Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2020, 27, 2559-2568. | 0.7 | 11 |
| 193 | Characteristics Associated with Nodal and Distant Recurrence After Radical Esophagectomy for Squamous Cell Carcinoma of the Thoracic Esophagus. <i>Annals of Surgical Oncology</i> , 2020, 27, 3195-3205. | 0.7 | 11 |
| 194 | miR-23b-3p Plays an Oncogenic Role in Hepatocellular Carcinoma. <i>Annals of Surgical Oncology</i> , 2021, 28, 3416-3426. | 0.7 | 11 |
| 195 | Peritoneal Lavage Tumor DNA as a Novel Biomarker for Predicting Peritoneal Recurrence in Pancreatic Ductal Adenocarcinoma. <i>Annals of Surgical Oncology</i> , 2021, 28, 2277-2286. | 0.7 | 11 |
| 196 | Prognostic Implications of Intraoperative Radiotherapy for Unresectable Pancreatic Cancer. <i>Pancreatology</i> , 2011, 11, 68-75. | 0.5 | 10 |
| 197 | Protein tyrosine kinase 7: a hepatocellular carcinoma-related gene detected by triple-combination array. <i>Journal of Surgical Research</i> , 2015, 195, 444-453. | 0.8 | 10 |
| 198 | Combination of continuous paravertebral block and epidural anesthesia in postoperative pain control after esophagectomy. <i>Esophagus</i> , 2016, 13, 42-47. | 1.0 | 10 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 199 | ASO Author Reflections: Troponin I2â€”A Specific Biomarker for Detection and Prediction of Peritoneal Metastasis in Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2018, 25, 709-710. | 0.7 | 10 |
| 200 | A novel dualâ€”marker expression panel for easy and accurate risk stratification of patients with gastric cancer. <i>Cancer Medicine</i> , 2018, 7, 2463-2471. | 1.3 | 10 |
| 201 | Short-term outcomes after conventional transthoracic esophagectomy. <i>Nagoya Journal of Medical Science</i> , 2016, 78, 69-78. | 0.6 | 10 |
| 202 | Operative Treatment of Thrombotic Occlusion of the Portal Vein Immediately After Pancreatectomy With Portal Vein Resection. <i>Pancreas</i> , 2010, 39, 265-266. | 0.5 | 9 |
| 203 | <i>PRAME</i> Expression as a Potential Biomarker for Hematogenous Recurrence of Esophageal Squamous Cell Carcinoma. <i>Anticancer Research</i> , 2019, 39, 5943-5951. | 0.5 | 9 |
| 204 | Tissue Expression of Melanoma-associated Antigen A6 and Clinical Characteristics of Gastric Cancer. <i>Anticancer Research</i> , 2019, 39, 5903-5910. | 0.5 | 9 |
| 205 | STRA6 Expression Serves as a Prognostic Biomarker of Gastric Cancer. <i>Cancer Genomics and Proteomics</i> , 2020, 17, 509-516. | 1.0 | 9 |
| 206 | AMIGO2 Expression as a Potential Prognostic Biomarker for Gastric Cancer. <i>Anticancer Research</i> , 2020, 40, 6713-6721. | 0.5 | 9 |
| 207 | Expression and Malignant Potential of B4GALNT4 in Esophageal Squamous Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2020, 27, 3247-3256. | 0.7 | 9 |
| 208 | Association between Lymphovascular Invasion and Recurrence in Patients with pT1N+ or pT2â€”3N0 Gastric Cancer: a Multi-institutional Dataset Analysis. <i>Journal of Gastric Cancer</i> , 2020, 20, 41. | 0.9 | 9 |
| 209 | D2 lymph node dissection confers little benefit on the overall survival of older patients with resectable gastric cancer: a propensity score-matching analysis of a multi-institutional dataset. <i>Surgery Today</i> , 2020, 50, 1434-1442. | 0.7 | 9 |
| 210 | Platelet isoform of phosphofructokinase accelerates malignant features in breast cancer. <i>Oncology Reports</i> , 2021, 47, . | 1.2 | 9 |
| 211 | Preservation of the Pyloric Ring Confers Little Benefit in Patients Undergoing Total Pancreatectomy. <i>World Journal of Surgery</i> , 2014, 38, 1807-1813. | 0.8 | 8 |
| 212 | High expression of Janus kinase 2 in background normal liver tissue of resected hepatocellular carcinoma is associated with worse prognosis. <i>Oncology Reports</i> , 2015, 33, 767-773. | 1.2 | 8 |
| 213 | Prognostic Impact of Portal System Invasion in Pancreatic Cancer Based on Image Classification. <i>Pancreas</i> , 2018, 47, 1350-1356. | 0.5 | 8 |
| 214 | Accurate Risk Stratification of Patients with Nodeâ€”Positive Gastric Cancer by Lymph Node Ratio. <i>World Journal of Surgery</i> , 2020, 44, 4184-4192. | 0.8 | 8 |
| 215 | Characteristics of Lung Metastasis as an Initial Recurrence Pattern After Curative Resection of Pancreatic Cancer. <i>Pancreas</i> , 2020, 49, 699-705. | 0.5 | 8 |
| 216 | Integrated multigene expression panel to prognosticate patients with gastric cancer. <i>Oncotarget</i> , 2018, 9, 18775-18785. | 0.8 | 8 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 217 | Intraperitoneal Chemotherapy as Adjuvant or Perioperative Chemotherapy for Patients with Type 4 Scirrhous Gastric Cancer: PHOENIX-GC2 Trial. <i>Journal of Clinical Medicine</i> , 2021, 10, 5666. | 1.0 | 8 |
| 218 | The COMET Open-label Phase II Study of Neoadjuvant FOLFOX or XELOX Treatment Combined with Molecular Targeting Monoclonal Antibodies in Patients with Resectable Liver Metastasis of Colorectal Cancer. <i>Annals of Surgical Oncology</i> , 2017, 24, 546-553. | 0.7 | 7 |
| 219 | Nutritional Recovery after Open and Laparoscopic Distal Gastrectomy for Early Gastric Cancer: A Prospective Multicenter Comparative Trial (CCOG1204). <i>Digestive Surgery</i> , 2018, 35, 11-18. | 0.6 | 7 |
| 220 | Optical trocar access for initial trocar placement in laparoscopic gastrointestinal surgery: propensity score matching analysis. <i>Asian Journal of Endoscopic Surgery</i> , 2019, 12, 37-42. | 0.4 | 7 |
| 221 | An Open-Label Single-Arm Phase II Study of Treatment with Neoadjuvant S-1 Plus Cisplatin for Clinical Stage III Squamous Cell Carcinoma of the Esophagus. <i>Oncologist</i> , 2020, 25, e1650-e1654. | 1.9 | 7 |
| 222 | Correlation Between Poor Prognosis and Lower TPPP Gene Expression in Hepatocellular Carcinoma. <i>Anticancer Research</i> , 2016, 36, 4639-4646. | 0.5 | 7 |
| 223 | Is the measurement of drain amylase content useful for predicting pancreas-related complications after gastrectomy with systematic lymphadenectomy?. <i>World Journal of Gastroenterology</i> , 2020, 26, 1594-1600. | 1.4 | 7 |
| 224 | Prognostic Value of a Modified Albumin-Bilirubin Score Designed for Patients with Esophageal Squamous Cell Carcinoma After Radical Resection. <i>Annals of Surgical Oncology</i> , 2022, 29, 4889-4896. | 0.7 | 7 |
| 225 | Expression of regulatory factor X1 can predict the prognosis of breast cancer. <i>Oncology Letters</i> , 2017, 13, 4334-4340. | 0.8 | 6 |
| 226 | Proposal of a Scoring Scale to Estimate Risk of the Discontinuation of S-1 Adjuvant Monotherapy in Patients with Stage II to III Gastric Cancer: A Multi-Institutional Dataset Analysis. <i>World Journal of Surgery</i> , 2019, 43, 2016-2024. | 0.8 | 6 |
| 227 | Phase II study of capecitabine plus oxaliplatin (CapOX) as adjuvant chemotherapy for locally advanced rectal cancer (CORONA II). <i>International Journal of Clinical Oncology</i> , 2020, 25, 118-125. | 1.0 | 6 |
| 228 | Survival times are similar among patients with peritoneal, hematogenous, and nodal recurrences after curative resections for gastric cancer. <i>Cancer Medicine</i> , 2020, 9, 5392-5399. | 1.3 | 6 |
| 229 | Chromobox 2 Expression Predicts Prognosis After Curative Resection of Oesophageal Squamous Cell Carcinoma. <i>Cancer Genomics and Proteomics</i> , 2020, 17, 391-400. | 1.0 | 6 |
| 230 | High Serum Uric Acid Levels Could Be a Risk Factor of Hepatocellular Carcinoma Recurrences. <i>Nutrition and Cancer</i> , 2021, 73, 996-1003. | 0.9 | 6 |
| 231 | Accurate Prediction of Prognosis After Radical Resection of Gastric Cancer by the Modified Systemic Inflammation Score; a Multicenter Dataset Analysis. <i>World Journal of Surgery</i> , 2021, 45, 2513-2520. | 0.8 | 6 |
| 232 | Preoperative docetaxel, cisplatin, and fluorouracil treatment with pegfilgrastim on day 7 for patients with esophageal cancer: A phase II study. <i>Asia-Pacific Journal of Clinical Oncology</i> , 2022, 18, 578-585. | 0.7 | 6 |
| 233 | Method of Bilateral Pleural Drainage by Single Blake Drain After Esophagectomy. <i>World Journal of Surgery</i> , 2015, 39, 727-731. | 0.8 | 5 |
| 234 | Modified two-dimensional response as surrogate marker of overall survival in patients with metastatic colorectal cancer. <i>Cancer Science</i> , 2016, 107, 1492-1498. | 1.7 | 5 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 235 | Randomized phase II study of daily and alternate-day administration of S-1 for advanced gastric cancer (JFMC43-1003). <i>International Journal of Clinical Oncology</i> , 2017, 22, 1052-1059. | 1.0 | 5 |
| 236 | Expression, Function, and Prognostic Value of MAGE-D4 Protein in Esophageal Squamous Cell Carcinoma. <i>Anticancer Research</i> , 2019, 39, 6015-6023. | 0.5 | 5 |
| 237 | Phase I Study of Intraperitoneal Administration of Paclitaxel Combined with S-1 Plus Cisplatin for Gastric Cancer with Peritoneal Metastasis. <i>Oncology</i> , 2020, 98, 48-52. | 0.9 | 5 |
| 238 | Tissue <i>RNF2</i> Expression Levels Are Associated With Peritoneal Recurrence and Poor Prognosis in Gastric Cancer. <i>Anticancer Research</i> , 2021, 41, 609-617. | 0.5 | 5 |
| 239 | Neoadjuvant chemoradiotherapy with S-1 in patients with borderline resectable pancreatic cancer.. <i>Journal of Clinical Oncology</i> , 2014, 32, 302-302. | 0.8 | 5 |
| 240 | A phase II trial to evaluate the efficacy of panitumumab combined with fluorouracil-based chemotherapy for metastatic colorectal cancer: the PF trial. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 81, 829-838. | 1.1 | 4 |
| 241 | Tumor size ≤ 50 mm as an Independent Prognostic Factor for Patients with Stage II or III Gastric Cancer After Postoperative ≤ 1 Monotherapy: Analysis of a Multi-institution Dataset. <i>World Journal of Surgery</i> , 2020, 44, 194-201. | 0.8 | 4 |
| 242 | Short-term outcomes of gastrectomy after neoadjuvant chemotherapy for clinical stage III gastric cancer: propensity score-matched analysis of a multi-institutional database. <i>Surgery Today</i> , 2021, 51, 821-828. | 0.7 | 4 |
| 243 | An integrated multigene expression panel to predict long-term survival after curative hepatectomy in patients with hepatocellular carcinoma. <i>Oncotarget</i> , 2017, 8, 71070-71079. | 0.8 | 4 |
| 244 | Surveillance of Esophageal Cancer in the Republic of Uzbekistan from 2000 to 2018. <i>Asian Pacific Journal of Cancer Prevention</i> , 2020, 21, 2281-2285. | 0.5 | 4 |
| 245 | Expression of cellular retinoic acid binding protein 1 predicts peritoneal recurrence of gastric cancer. <i>International Journal of Oncology</i> , 2022, 60, . | 1.4 | 4 |
| 246 | Detection of serum melanoma-associated antigen D4 in patients with squamous cell carcinoma of the esophagus. <i>Ecological Management and Restoration</i> , 2016, 29, 663-669. | 0.2 | 3 |
| 247 | RASEF expression correlates with hormone receptor status in breast cancer. <i>Oncology Letters</i> , 2018, 16, 7223-7230. | 0.8 | 3 |
| 248 | Cutting-edge evidence of adjuvant treatments for gastric cancer. <i>Expert Review of Gastroenterology and Hepatology</i> , 2018, 12, 1109-1122. | 1.4 | 3 |
| 249 | Randomised phase II trial of capecitabine plus oxaliplatin with continuous versus intermittent use of oxaliplatin as adjuvant chemotherapy for stage II/III colon cancer (CCOG-1302 study). <i>European Journal of Cancer</i> , 2021, 144, 61-71. | 1.3 | 3 |
| 250 | Update on molecular biomarkers for diagnosis and prediction of prognosis and treatment responses in gastric cancer. <i>Histology and Histopathology</i> , 2021, 36, 817-832. | 0.5 | 3 |
| 251 | Synaptotagmin 13 Is Highly Expressed in Estrogen Receptor-Positive Breast Cancer. <i>Current Oncology</i> , 2021, 28, 4080-4092. | 0.9 | 3 |
| 252 | Editors' Choice Efficacy of enteral nutrients containing β -hydroxy- β -methylbutyrate, glutamine, and arginine for the patients with anastomotic leakage after gastrectomy: study protocol of a multicenter phase II clinical trial. <i>Nagoya Journal of Medical Science</i> , 2018, 80, 351-355. | 0.6 | 3 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 253 | SLC7A9 as a Potential Biomarker for Lymph Node Metastasis of Esophageal Squamous Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2022, 29, 2699-2709. | 0.7 | 3 |
| 254 | A Vascular Endothelial Growth Factor Gene Polymorphism Predicts Malignant Potential in Intraductal Papillary Mucinous Neoplasm. <i>Pancreas</i> , 2015, 44, 608-614. | 0.5 | 2 |
| 255 | Prognosis After Laparoscopic Gastrectomy in Patients with Pathological Stage II or III Gastric Cancer Who Were Preoperatively Diagnosed with Clinical Stage I: Propensity Score Matching Analysis of a Multicenter Dataset. <i>Annals of Surgical Oncology</i> , 2020, 27, 268-275. | 0.7 | 2 |
| 256 | Newly developed primary malignancies in long-term survivors who underwent curative esophagectomy for squamous cell carcinoma of the esophagus. <i>Surgery Today</i> , 2021, 51, 153-158. | 0.7 | 2 |
| 257 | Efficacy of Splenectomy for Proximal Gastric Cancer with Greater Curvature Invasion or Type 4 Tumor: a Propensity Score Analysis of a Multi-institutional Dataset. <i>World Journal of Surgery</i> , 2021, 45, 2840-2848. | 0.8 | 2 |
| 258 | Prognostic impact of a microscopic positive margin in patients undergoing gastrectomy for gastric cancer: a propensity score-matched analysis of a multi-institutional dataset. <i>Surgery Today</i> , 2022, 52, 559-566. | 0.7 | 2 |
| 259 | E-PASS scoring system serves as a predictor of short- and long-term outcomes in gastric cancer surgery. <i>Surgery Today</i> , 2022, 52, 914-922. | 0.7 | 2 |
| 260 | Intraductal Papillary Mucinous Carcinoma with Portal Annular Pancreas Anomaly Treated by Middle-preserving Pancreatectomy. <i>Japanese Journal of Gastroenterological Surgery</i> , 2015, 48, 706-714. | 0.0 | 2 |
| 261 | Transcriptomic profiling on localized gastric cancer identified CPLX1 as a gene promoting malignant phenotype of gastric cancer and a predictor of recurrence after surgery and subsequent chemotherapy. <i>Journal of Gastroenterology</i> , 2022, 57, 640-653. | 2.3 | 2 |
| 262 | Preoperative neutrophil-to-platelet ratio as a potential prognostic factor for gastric cancer with positive peritoneal lavage cytology in the absence of other non-curative factors: a multi-institutional dataset analysis. <i>Surgery Today</i> , 2023, 53, 198-206. | 0.7 | 2 |
| 263 | Operative Treatment of Pancreatic Ductal Adenocarcinoma With Extensive Portal Venous Tumor Embolism. <i>Pancreas</i> , 2010, 39, 268-269. | 0.5 | 1 |
| 264 | Stapling an extracorporeal Billroth-II anastomosis by the complete double stapling technique after laparoscopy-assisted distal gastrectomy. <i>Asian Journal of Endoscopic Surgery</i> , 2017, 10, 137-142. | 0.4 | 1 |
| 265 | ASO Author Reflections: Homeobox C10 Influences on the Malignant Phenotype of Gastric Cancer Cell Lines and its Elevated Expression Positively Correlates with Recurrence and Poor Survival. <i>Annals of Surgical Oncology</i> , 2019, 26, 596-597. | 0.7 | 1 |
| 266 | Clinical impact of additional therapy for residual pancreatic cancer. <i>Surgery Today</i> , 2020, 50, 440-448. | 0.7 | 1 |
| 267 | Impact of molecular surgical margin analysis on the prediction of pancreatic cancer recurrences after pancreaticoduodenectomy. <i>Clinical Epigenetics</i> , 2021, 13, 172. | 1.8 | 1 |
| 268 | Clinical features of colorectal mucinous and poorly differentiated adenocarcinomas; study concept of a propensity score analysis in a pooled data of 5530 patients. <i>Annals of Cancer Research and Therapy</i> , 2016, 24, 52-53. | 0.1 | 1 |
| 269 | Abstract 3831: Clinical significance of SMAD4 expression in resectable pancreatic cancer: correlation with tumor progression and recurrence pattern. , 2014, , . | | 1 |
| 270 | Abstract 1148: Epithelial to mesenchymal transition might be induced via CD44 isoform switch in colorectal cancer. , 2014, , . | | 1 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 271 | ASO Visual Abstract: Prognostic Value of a Modified Albuminâ€“Bilirubin Grade Designed for Patients with Esophageal Squamous Cell Carcinoma after Radical Resection. <i>Annals of Surgical Oncology</i> , 2022, , 1. | 0.7 | 1 |
| 272 | ASO Author Reflections: Increased Expression of DNAJC12 is Associated with Aggressive Phenotype of Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2019, 26, 592-593. | 0.7 | 0 |
| 273 | ASO Author Reflections: Characteristics Associated with Nodal and Distant Recurrence After Radical Esophagectomy for Squamous Cell Carcinoma of the Thoracic Esophagus. <i>Annals of Surgical Oncology</i> , 2020, 27, 3206-3207. | 0.7 | 0 |
| 274 | ASO Author Reflections: KCNJ15 Expression and Malignant Behavior of Esophageal Squamous Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2020, 27, 2569-2570. | 0.7 | 0 |
| 275 | ASO Author Reflections: Expression and Malignant Potential of B4GALNT4 in Esophageal Squamous Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2020, 27, 3257-3258. | 0.7 | 0 |
| 276 | Age-Related Differences in the Prognosis of Pancreatic Cancer According to Perioperative Systemic Therapy. <i>Pancreas</i> , 2021, 50, 37-46. | 0.5 | 0 |
| 277 | Abstract 1733: Fibulin 1 is a novel tumor suppressor gene detected in hepatocellular carcinoma using a double combination array analysis. , 2010, , . | | 0 |
| 278 | Abstract 4660: Double combination array analysis detected A kinase anchor protein 12 (AKAP12) gene as a new tumor suppressor gene of hepatocellular carcinoma. , 2010, , . | | 0 |
| 279 | Abstract 1750: Reduced xpression of Reelin (RELN) gene is associated with high recurrence rate of hepatocellular carcinoma. , 2010, , . | | 0 |
| 280 | Abstract 3539: A study ofTHOP1, a predictive factor of prognosis in HCC, by multi-array analysis of background liver.. , 2013, , . | | 0 |
| 281 | Overexpression of melanoma-associated antigen D4 as an independent prognostic factor in squamous cell carcinoma of the esophagus.. <i>Journal of Clinical Oncology</i> , 2014, 32, 71-71. | 0.8 | 0 |
| 282 | Risk factors that affect continuation of S-1 chemotherapy as an adjuvant setting after radical gastrectomy for gastric cancer.. <i>Journal of Clinical Oncology</i> , 2014, 32, 151-151. | 0.8 | 0 |
| 283 | Response to the first-line FOLFOX plus bevacizumab (BEV) therapy to predict responses to the subsequent therapies and survival in the BEV beyond progression (BBP) strategy for metastatic colorectal cancer: A retrospective analysis of CCOG-0801 study.. <i>Journal of Clinical Oncology</i> , 2014, 32, 428-428. | 0.8 | 0 |
| 284 | Reduced expression of DENND2D through promoter hypermethylation as an adverse prognostic factor in squamous cell carcinoma of the esophagus.. <i>Journal of Clinical Oncology</i> , 2014, 32, 58-58. | 0.8 | 0 |
| 285 | The impact of dose reduction and time delay in irinotecan- and oxaliplatin-based chemotherapies on outcomes in metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2014, 32, 631-631. | 0.8 | 0 |
| 286 | Abstract 2870: ZGPAT gene expression in non-tumor hepatocellular carcinoma tissue is a likely biomarker for survival risk. , 2014, , . | | 0 |
| 287 | Abstract 3822: Correlation between worse prognosis and lower expression of theTPPPgene in patients with hepatocellular carcinoma, detected by multiarray analysis. , 2014, , . | | 0 |
| 288 | Abstract 4717: Detection of the Cyclin J (CCNJ) as a new cancer-related gene in human hepatocellular carcinoma by using a method of triple combination array analysis. , 2014, , . | | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 289 | Abstract 3816: Correlation between worse prognosis and higher expression of the JAK2 gene in corresponding non-neoplastic tissue in patients with hepatocellular carcinoma, extracted by multiarray analysis. , 2014, , . | | 0 |
| 290 | Abstract 3423: Expression level of inflammasomes components NLRP3, NLRC4, and CASP1 in background non tumorous tissue were associated with worse prognosis for curatively resected hepatocellular carcinoma. , 2015, , . | | 0 |
| 291 | Abstract 4101: Clinical significance of ZEB1 mRNA levels in peritoneal washing for gastric cancer. , 2015, , . | | 0 |
| 292 | Abstract 3422: Alteration of aldo-keto reductase family 1, member B10 (AKR1B10) expression among tumor and background non-tumorous tissue of curatively resected hepatocellular carcinoma is associated with worse prognosis. , 2015, , . | | 0 |
| 293 | Abstract 4253: A vascular endothelial growth factor gene polymorphism predicts malignant potential in intraductal papillary mucinous neoplasm. , 2015, , . | | 0 |
| 294 | A prospective trial to evaluate treatment effects of a β -hydroxy- β -methylbutyrate containing nutrient for leakage at the anastomotic site after esophagectomy. Nagoya Journal of Medical Science, 2020, 82, 33-37. | 0.6 | 0 |
| 295 | Neoadjuvant docetaxel, oxaliplatin plus S-1 for treating clinical stage III squamous cell carcinoma of the esophagus: Study protocol of an open-label phase II trial. Contemporary Clinical Trials Communications, 2021, 24, 100853. | 0.5 | 0 |
| 296 | ASO Visual Abstract: SLC7A9 as a Potential Biomarker for Lymph Node Metastasis of Esophageal Squamous Cell Carcinoma. Annals of Surgical Oncology, 2022, 29, 2710. | 0.7 | 0 |
| 297 | Drain Amylase Concentrations at 3 h After Gastrectomy Enhance Early Prediction of Postoperative Peripancreatic Inflammatory Fluid Collection. World Journal of Surgery, 2022, 46, 648-655. | 0.8 | 0 |
| 298 | ASO Author Reflections: Optimized Cutoff Value of Albumin-Bilirubin Score to Predict Prognosis of Patients with Esophageal Squamous Cell Carcinoma After Radical Resection. Annals of Surgical Oncology, 2022, , 1. | 0.7 | 0 |