

# Kazuo Kubota

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2220814/publications.pdf>

Version: 2024-02-01

34  
papers

1,717  
citations

304743

22  
h-index

454955

30  
g-index

35  
all docs

35  
docs citations

35  
times ranked

1859  
citing authors

#	ARTICLE	IF	CITATIONS
1	[18F]FDG uptake in axillary lymph nodes and deltoid muscle after COVID-19 mRNA vaccination: a cohort study to determine incidence and contributing factors using a multivariate analysis. <i>Annals of Nuclear Medicine</i> , 2022, 36, 340-350.	2.2	11
2	Comparison of 18F-FDG PET/CT and 67Ga-SPECT for the diagnosis of fever of unknown origin: a multicenter prospective study in Japan. <i>Annals of Nuclear Medicine</i> , 2021, 35, 31-46.	2.2	12
3	FDG-PET/CT for Large-Vessel Vasculitis. , 2020, , 115-146.		0
4	FDG-PET/CT in Patients with Inflammation or Fever of Unknown Origin (IUO and FUO). , 2020, , 43-55.		0
5	Clinical Value of FDG-PET/CT for the Evaluation of Rheumatic Diseases: Rheumatoid Arthritis, Polymyalgia Rheumatica, and Relapsing Polychondritis. <i>Seminars in Nuclear Medicine</i> , 2017, 47, 408-424.	4.6	66
6	Impact of FDG-PET findings on decisions regarding patient management strategies: a multicenter trial in patients with lung cancer and other types of cancer. <i>Annals of Nuclear Medicine</i> , 2015, 29, 431-441.	2.2	28
7	Differences in fluorodeoxyglucose positron emission tomography/computed tomography findings between elderly onset rheumatoid arthritis and polymyalgia rheumatica. <i>Modern Rheumatology</i> , 2015, 25, 546-551.	1.8	53
8	Utility of fluorodeoxyglucose positron emission tomography/computed tomography for early diagnosis and evaluation of disease activity of relapsing polychondritis: a case series and literature review. <i>Rheumatology</i> , 2014, 53, 1482-1490.	1.9	64
9	Lesion-based analysis of 18F-FDG uptake and 111In-Pentetreotide uptake by neuroendocrine tumors. <i>Annals of Nuclear Medicine</i> , 2014, 28, 1004-1010.	2.2	42
10	Large vessel vasculitis in elderly patients: early diagnosis and steroid-response evaluation with FDG-PET/CT and contrast-enhanced CT. <i>Rheumatology International</i> , 2014, 34, 1545-1554.	3.0	43
11	Clinical value of <sup>18</sup> F-fluoro-dexoxyglucose positron emission tomography/computed tomography in patients with adult-onset Still's disease: A seven-case series and review of the literature. <i>Modern Rheumatology</i> , 2014, 24, 645-650.	1.8	47
12	Clinical value of whole-body PET/CT in patients with active rheumatic diseases. <i>Arthritis Research and Therapy</i> , 2014, 16, 423.	3.5	81
13	Whole-body fluorodeoxyglucose positron emission tomography/computed tomography in patients with active polymyalgia rheumatica: evidence for distinctive bursitis and large-vessel vasculitis. <i>Modern Rheumatology</i> , 2012, 22, 705-711.	1.8	84
14	Validation for performing 11C-methionine and 18F-FDG-PET studies on the same day. <i>Nuclear Medicine Communications</i> , 2012, 33, 297-304.	1.1	2
15	Value of Carotid Artery Tenderness for the Early Diagnosis of Takayasu Arteritis. <i>Internal Medicine</i> , 2012, 51, 3431-3434.	0.7	3
16	Effects of blood glucose level on FDG uptake by liver: a FDG-PET/CT study. <i>Nuclear Medicine and Biology</i> , 2011, 38, 347-351.	0.6	63
17	Clinical Role of FDG PET/CT for Methotrexate-Related Malignant Lymphoma. <i>Clinical Nuclear Medicine</i> , 2011, 36, 533-537.	1.3	11
18	FDG PET for rheumatoid arthritis: basic considerations and whole-body PET/CT. <i>Annals of the New York Academy of Sciences</i> , 2011, 1228, 29-38.	3.8	65

#	ARTICLE	IF	CITATIONS
19	FDG-PET for the diagnosis of fever of unknown origin: a Japanese multi-center study. <i>Annals of Nuclear Medicine</i> , 2011, 25, 355-364.	2.2	68
20	Clinical impact of 18F-FDG PET/CT on the management and diagnosis of infectious spondylitis. <i>Nuclear Medicine Communications</i> , 2010, 31, 691-698.	1.1	32
21	Constrictive tuberculous pericarditis diagnosed using 18F-fluorodeoxyglucose positron emission tomography: a report of two cases. <i>Annals of Nuclear Medicine</i> , 2010, 24, 421-425.	2.2	19
22	Whole-body FDG-PET/CT on rheumatoid arthritis of large joints. <i>Annals of Nuclear Medicine</i> , 2009, 23, 783-791.	2.2	111
23	Advantage of delayed whole-body FDG-PET imaging for tumour detection. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2001, 28, 696-703.	2.1	221
24	From tumor biology to clinical PET: A review of positron emission tomography (PET) in oncology. <i>Annals of Nuclear Medicine</i> , 2001, 15, 471-486.	2.2	234
25	Relationship between trait anxiety, brain activity and natural killer cell activity in cancer patients: a preliminary PET study. <i>Psycho-Oncology</i> , 2001, 10, 541-546.	2.3	28
26	Reproducibility of PET brain mapping of cancer patients. , 2000, 9, 157-163.		15
27	Glucose uptake by individual skeletal muscles during running using whole-body positron emission tomography. <i>European Journal of Applied Physiology</i> , 2000, 83, 297-302.	2.5	51
28	Hypometabolism in the limbic system of cancer patients observed by positron emission tomography. , 1999, 8, 283-286.		26
29	Effects of smoking on regional cerebral blood flow in cerebral vascular disease patients and normal subjects.. <i>Tohoku Journal of Experimental Medicine</i> , 1987, 151, 261-268.	1.2	7
30	Experimental studies on myocardial glucose metabolism of rats with 18F-2-fluoro-2-deoxy-d-glucose. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 1985, 10, 341-5.	2.1	25
31	Tumor detection with carbon-11-labelled amino acids. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 1984, 9, 136-140.	2.1	76
32	Studies on 18F-labeled pyrimidines. Tumor uptakes of 18F-5-fluorouracil, 18F-5-fluorouridine, and 18F-5-fluorodeoxyuridine in animals. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 1983, 8, 258-261.	2.1	56
33	A Quantitative Approach of Abdominal Aortic Atherosclerosis with X-ray Computed Tomography. <i>The Journal of Japan Atherosclerosis Society</i> , 1983, 11, 463-466.	0.0	2
34	Experimental study for cancer diagnosis with positron-labeled fluorinated glucose analogs: [18F]-2-fluoro-2-deoxy-D-mannose: A new tracer for cancer detection. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 1982, 7, 294-297.	2.1	71