
DonajÃ- VerÃ³nica GÃ³mez-Velasco

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2226200/publications.pdf

Version: 2024-02-01



DonajÃ-VerÃ³nica

#	Article	IF	CITATIONS
1	METS-IR, a novel score to evaluate insulin sensitivity, is predictive of visceral adiposity and incident type 2 diabetes. European Journal of Endocrinology, 2018, 178, 533-544.	1.9	173
2	A Loss-of-Function Splice Acceptor Variant in <i>IGF2</i> Is Protective for Type 2 Diabetes. Diabetes, 2017, 66, 2903-2914.	0.3	52
3	Metabolic Score for Visceral Fat (METS-VF), a novel estimator of intra-abdominal fat content and cardio-metabolic health. Clinical Nutrition, 2020, 39, 1613-1621.	2.3	37
4	Natural course of metabolically healthy phenotype and risk of developing Cardiometabolic diseases: a three years follow-up study. BMC Endocrine Disorders, 2021, 21, 85.	0.9	28
5	Clinical and metabolomic predictors of regression to normoglycemia in a population at intermediate cardiometabolic risk. Cardiovascular Diabetology, 2021, 20, 56.	2.7	10
6	Familial hypertriglyceridemia: an entity with distinguishable features from other causes of hypertriglyceridemia. Lipids in Health and Disease, 2021, 20, 14.	1.2	9
7	Elevated serum uric acid is a facilitating mechanism for insulin resistance mediated accumulation of visceral adipose tissue. Clinical Endocrinology, 2022, 96, 707-718.	1.2	8
8	Contribution of Known Genetic Risk Variants to Dyslipidemias and Type 2 Diabetes in Mexico: A Population-Based Nationwide Study. Genes, 2020, 11, 114.	1.0	7
9	Primary Barriers of Adherence to a Structured Nutritional Intervention in Patients with Dyslipidemia. Nutrients, 2021, 13, 1744.	1.7	7
10	Evaluation of a Web Platform to Record Lifestyle Habits in Subjects at Risk of Developing Type 2 Diabetes in a Middle-Income Population: Prospective Interventional Study. JMIR Diabetes, 2022, 7, e25105.	0.9	4