

Marisa Borenstein

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2235930/publications.pdf>

Version: 2024-02-01

11
papers

337
citations

1162367

8
h-index

1473754

9
g-index

12
all docs

12
docs citations

12
times ranked

290
citing authors

#	ARTICLE	IF	CITATIONS
1	Fetal optic chiasm: three steps for visualization and measurement on routine transabdominal ultrasound. <i>Ultrasound in Obstetrics and Gynecology</i> , 2019, 54, 135-136.	0.9	4
2	Visualizaci3n de la anatom3a fetal por ecograf3a prenatal entre las semanas 11-13+6. <i>Diagnostico Prenatal</i> , 2011, 22, 68-68.	0.1	0
3	Depth of brain fissures in normal fetuses by prenatal ultrasound between 19 and 30 weeks of gestation. <i>Ultrasound in Obstetrics and Gynecology</i> , 2010, 36, 693-699.	0.9	48
4	Aberrant right subclavian artery at 16 to 23 + 6 weeks of gestation: a marker for chromosomal abnormality. <i>Ultrasound in Obstetrics and Gynecology</i> , 2010, 36, 548-552.	0.9	60
5	P04.03: Frontomaxillary facial angle at 11 + 0 to 13 + 6 weeks in Chinese population. <i>Ultrasound in Obstetrics and Gynecology</i> , 2010, 36, 180-180.	0.9	0
6	Gestational sac volume: comparison between SonoAVC and VOCAL measurements at 11 + 0 to 13 + 6 weeks of gestation. <i>Ultrasound in Obstetrics and Gynecology</i> , 2009, 34, 510-514.	0.9	10
7	Aberrant right subclavian artery at 11 + 0 to 13 + 6 weeks of gestation in chromosomally normal and abnormal fetuses. <i>Ultrasound in Obstetrics and Gynecology</i> , 2008, 31, 20-24.	0.9	55
8	Frontomaxillary facial angle in screening for trisomy 21 at 11 + 0 to 13 + 6 weeks. <i>Ultrasound in Obstetrics and Gynecology</i> , 2008, 32, 5-11.	0.9	63
9	Frontomaxillary facial angle in chromosomally normal fetuses at 11 + 0 to 13 + 6 weeks. <i>Ultrasound in Obstetrics and Gynecology</i> , 2007, 30, 737-741.	0.9	39
10	Frontomaxillary facial angle in fetuses with trisomy 13 at 11 + 0 to 13 + 6 weeks. <i>Ultrasound in Obstetrics and Gynecology</i> , 2007, 30, 819-823.	0.9	22
11	Frontomaxillary and mandibulomaxillary facial angles at 11 + 0 to 13 + 6 weeks in fetuses with trisomy 18. <i>Ultrasound in Obstetrics and Gynecology</i> , 2007, 30, 928-933.	0.9	36