

Koji Oba

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/224072/publications.pdf>

Version: 2024-02-01

251
papers

6,038
citations

87888
38
h-index

95266
68
g-index

254
all docs

254
docs citations

254
times ranked

8870
citing authors

#	ARTICLE	IF	CITATIONS
1	Benefit of Adjuvant Chemotherapy for Resectable Gastric Cancer. JAMA - Journal of the American Medical Association, 2010, 303, 1729.	7.4	711
2	Clinical significance of serum tumor markers for gastric cancer: a systematic review of literature by the Task Force of the Japanese Gastric Cancer Association. Gastric Cancer, 2014, 17, 26-33.	5.3	395
3	Effects of Candesartan Compared With Amlodipine in Hypertensive Patients With High Cardiovascular Risks. Hypertension, 2008, 51, 393-398.	2.7	227
4	Comparison of Targeted vs Random Biopsies for Surveillance of Ulcerative Colitis-Associated Colorectal Cancer. Gastroenterology, 2016, 151, 1122-1130.	1.3	171
5	Role of chemotherapy for advanced/recurrent gastric cancer: An individual-patient-data meta-analysis. European Journal of Cancer, 2013, 49, 1565-1577.	2.8	136
6	Disease-Free Survival as a Surrogate for Overall Survival in Adjuvant Trials of Gastric Cancer: A Meta-Analysis. Journal of the National Cancer Institute, 2013, 105, 1600-1607.	6.3	133
7	Safety and efficacy of rituximab in systemic sclerosis (DESIREs): a double-blind, investigator-initiated, randomised, placebo-controlled trial. Lancet Rheumatology, The, 2021, 3, e489-e497.	3.9	105
8	Sequential paclitaxel followed by tegafur and uracil (UFT) or S-1 versus UFT or S-1 monotherapy as adjuvant chemotherapy for T4a/b gastric cancer (SAMIT): a phase 3 factorial randomised controlled trial. Lancet Oncology, The, 2014, 15, 886-893.	10.7	104
9	The efficacy of tacrolimus in patients with interstitial lung diseases complicated with polymyositis or dermatomyositis. Rheumatology, 2015, 54, 39-44.	1.9	104
10	Diamine oxidase as a marker of intestinal mucosal injury and the effect of soluble dietary fiber on gastrointestinal tract toxicity after intravenous 5-fluorouracil treatment in rats. Medical Molecular Morphology, 2014, 47, 100-107.	1.0	98
11	Statistical evaluation of surrogate endpoints with examples from cancer clinical trials. Biometrical Journal, 2016, 58, 104-132.	1.0	93
12	Efficacy of adjuvant immunochemotherapy with polysaccharide K for patients with curative resections of gastric cancer. Cancer Immunology, Immunotherapy, 2007, 56, 905-911.	4.2	92
13	Vasoactive-inotropic score as a predictor of morbidity and mortality in adults after cardiac surgery with cardiopulmonary bypass. Journal of Anesthesia, 2018, 32, 167-173.	1.7	86
14	Does postoperative morbidity worsen the oncological outcome after radical surgery for gastrointestinal cancers? A systematic review of the literature. Annals of Gastroenterological Surgery, 2017, 1, 11-23.	2.4	85
15	High and low negative pressure suction techniques in EUS-guided fine-needle tissue acquisition by using 25-gauge needles: a multicenter, prospective, randomized, controlled trial. Gastrointestinal Endoscopy, 2014, 80, 1030-1037.e1.	1.0	83
16	Progression-Free Survival as a Surrogate for Overall Survival in Advanced/Recurrent Gastric Cancer Trials: A Meta-Analysis. Journal of the National Cancer Institute, 2013, 105, 1667-1670.	6.3	78
17	Efficacy of adjuvant Immunochemotherapy with polysaccharide K for patients with curatively resected colorectal cancer: a meta-analysis of centrally randomized controlled clinical trials. Cancer Immunology, Immunotherapy, 2006, 55, 404-411.	4.2	76
18	Impact of postoperative complications on the colorectal cancer survival and recurrence: analyses of pooled individual patients' data from three large phase III randomized trials. Cancer Medicine, 2017, 6, 1573-1580.	2.8	73

#	ARTICLE	IF	CITATIONS
19	The influence of childhood abuse, adult stressful life events and temperaments on depressive symptoms in the nonclinical general adult population. <i>Journal of Affective Disorders</i> , 2014, 158, 101-107.	4.1	68
20	Detection of early gastric cancer by a real-time autofluorescence imaging system. <i>Cancer Letters</i> , 2001, 165, 155-159.	7.2	67
21	Matched Pair Analysis to Examine the Effects of a Planned Preoperative Exercise Program in Early Gastric Cancer Patients with Metabolic Syndrome to Reduce Operative Risk: The Adjuvant Exercise for General Elective Surgery (AEGES) Study Group. <i>Annals of Surgical Oncology</i> , 2014, 21, 2044-2050.	1.5	63
22	Randomized controlled trial on the skin toxicity of panitumumab in Japanese patients with metastatic colorectal cancer: HGCSG1001 study; J-STEPP. <i>Future Oncology</i> , 2015, 11, 617-627.	2.4	61
23	An individual patient data meta-analysis of adjuvant therapy with uracil+tegafur (UFT) in patients with curatively resected rectal cancer. <i>British Journal of Cancer</i> , 2007, 96, 1170-1177.	6.4	59
24	Prognostic significance of epithelial-mesenchymal transition-related markers in extrahepatic cholangiocarcinoma: comprehensive immunohistochemical study using a tissue microarray. <i>British Journal of Cancer</i> , 2014, 111, 1363-1372.	6.4	59
25	Is Pulse Pressure a Predictor of New-Onset Diabetes in High-Risk Hypertensive Patients?. <i>Diabetes Care</i> , 2010, 33, 1122-1127.	8.6	58
26	Meta-analyses of randomized trials assessing the interest chemotherapy in advanced gastric cancer. <i>Journal of Clinical Oncology</i> , 2008, 26, 4563-4563.	1.6	56
27	Multicenter randomized controlled study to assess the effect of prophylactic clipping on post-polypectomy delayed bleeding. <i>Digestive Endoscopy</i> , 2016, 28, 570-576.	2.3	55
28	Pattern of programmed cell death-ligand 1 expression and CD8-positive T-cell infiltration before and after chemoradiotherapy in rectal cancer. <i>European Journal of Cancer</i> , 2018, 91, 11-20.	2.8	55
29	The Optimal Target Blood Pressure for Antihypertensive Treatment in Japanese Elderly Patients with High-Risk Hypertension: A Subanalysis of the Candesartan Antihypertensive Survival Evaluation in Japan (CASE-J) Trial. <i>Hypertension Research</i> , 2008, 31, 1595-1601.	2.7	47
30	Prospective Phase II Study of FOLFIRI for mCRC in Japan, Including the Analysis of UGT1A1*28/*6 Polymorphisms. <i>Japanese Journal of Clinical Oncology</i> , 2011, 41, 477-482.	1.3	47
31	Serum myostatin levels are independently associated with skeletal muscle wasting in patients with heart failure. <i>International Journal of Cardiology</i> , 2016, 220, 483-487.	1.7	47
32	FOLFIRI plus bevacizumab as second-line therapy in patients with metastatic colorectal cancer after first-line bevacizumab plus oxaliplatin-based therapy: the randomized phase III EAGLE study. <i>Annals of Oncology</i> , 2015, 26, 1427-1433.	1.2	45
33	Does remnant gastric cancer really differ from primary gastric cancer? A systematic review of the literature by the Task Force of Japanese Gastric Cancer Association. <i>Gastric Cancer</i> , 2016, 19, 339-349.	5.3	44
34	Phase II trial of paclitaxel and cisplatin as neoadjuvant chemotherapy for locally advanced gastric cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2013, 71, 1309-1314.	2.3	43
35	Therapeutic efficacy of azithromycin, clarithromycin, minocycline and tosufloxacin against macrolide-resistant and macrolide-sensitive <i>Mycoplasma pneumoniae</i> pneumonia in pediatric patients. <i>PLoS ONE</i> , 2017, 12, e0173635.	2.5	43
36	Long-term outcome in Japanese patients with lupus nephritis. <i>Lupus</i> , 2014, 23, 1124-1132.	1.6	41

#	ARTICLE	IF	CITATIONS
37	Effect of a stylet on a histological specimen in EUS-guided fine-needle tissue acquisition by using 22-gauge needles: a multicenter, prospective, randomized, controlled trial. <i>Gastrointestinal Endoscopy</i> , 2015, 82, 837-844.e1.	1.0	41
38	Acute Hemodynamic Effects of Adaptive Servo-Ventilation in Patients With Heart Failure. <i>Circulation Journal</i> , 2013, 77, 1214-1220.	1.6	40
39	Immunogenomic profiles associated with response to neoadjuvant chemoradiotherapy in patients with rectal cancer. <i>British Journal of Surgery</i> , 2019, 106, 1381-1392.	0.3	40
40	Radiomics Approach Outperforms Diameter Criteria for Predicting Pathological Lateral Lymph Node Metastasis After Neoadjuvant (Chemo)Radiotherapy in Advanced Low Rectal Cancer. <i>Annals of Surgical Oncology</i> , 2020, 27, 4273-4283.	1.5	40
41	A Comparison of the Effects of the GLP-1 Analogue Liraglutide and Insulin Glargine on Endothelial Function and Metabolic Parameters: A Randomized, Controlled Trial Sapporo Athero-Incretin Study 2 (SAIS2). <i>PLoS ONE</i> , 2015, 10, e0135854.	2.5	40
42	Open-label, randomized, comparative, phase III study on effects of reducing steroid use in combination with Palonosetron. <i>Cancer Science</i> , 2015, 106, 891-895.	3.9	39
43	Role of diabetes and obesity in outcomes of the candesartan antihypertensive survival evaluation in Japan (CASE-J) trial. <i>Hypertension Research</i> , 2010, 33, 600-606.	2.7	37
44	Effects of KIR ligand incompatibility on clinical outcomes of umbilical cord blood transplantation without ATG for acute leukemia in complete remission. <i>Blood Cancer Journal</i> , 2013, 3, e164-e164.	6.2	37
45	Risk Factors for Peritoneal Recurrence in Stage II to III Colon Cancer. <i>Diseases of the Colon and Rectum</i> , 2018, 61, 803-808.	1.3	36
46	Association of blood mercury levels during pregnancy with infant birth size by blood selenium levels in the Japan Environment and Children's Study: A prospective birth cohort. <i>Environment International</i> , 2019, 125, 418-429.	10.0	36
47	Comprehensive Comparative Analysis of Prognostic Value of Systemic Inflammatory Biomarkers for Patients with Stage II/III Colon Cancer. <i>Annals of Surgical Oncology</i> , 2020, 27, 844-852.	1.5	36
48	Patterns of Drug Use and HIV-Related Risk Behaviors among Incarcerated People in a Prison in Iran. <i>Journal of Urban Health</i> , 2010, 87, 603-616.	3.6	35
49	Decreased serum brain-derived neurotrophic factor levels are correlated with exercise intolerance in patients with heart failure. <i>International Journal of Cardiology</i> , 2013, 168, e142-e144.	1.7	35
50	Serum Brain-Derived Neurotrophic Factor Level Predicts Adverse Clinical Outcomes in Patients With Heart Failure. <i>Journal of Cardiac Failure</i> , 2015, 21, 300-306.	1.7	34
51	Assessing clinical benefit response in the treatment of gastric malignant ascites with non-measurable lesions: a multicenter phase II trial of paclitaxel for malignant ascites secondary to advanced/recurrent gastric cancer. <i>Gastric Cancer</i> , 2011, 14, 81-90.	5.3	32
52	A new prognostic classification for overall survival in Asian patients with previously untreated metastatic renal cell carcinoma. <i>Cancer Science</i> , 2012, 103, 1695-1700.	3.9	32
53	Efficacy of Adjuvant Chemotherapy Using Oral Fluorinated Pyrimidines for Curatively Resected Gastric Cancer: A Meta-Analysis of Centrally Randomized Controlled Clinical Trials in Japan. <i>Journal of Chemotherapy</i> , 2006, 18, 311-317.	1.5	30
54	Reduced-intensity vs myeloablative conditioning allogeneic hematopoietic SCT for patients aged over 45 years with ALL in remission: a study from the Adult ALL Working Group of the Japan Society for Hematopoietic Cell Transplantation (JSHCT). <i>Bone Marrow Transplantation</i> , 2013, 48, 1389-1394.	2.4	30

#	ARTICLE	IF	CITATIONS
55	Clinical effectiveness of four neuraminidase inhibitors (oseltamivir, zanamivir, laninamivir, and) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Japan. Journal of Infection and Chemotherapy, 2018, 24, 449-457.	1.7	30
56	Relationship between the achieved blood pressure and the incidence of cardiovascular events in Japanese hypertensive patients with complications: a sub-analysis of the CASE-J trial. Hypertension Research, 2009, 32, 248-254.	2.7	29
57	<scp><i>UGT1A1*6</i></scp>, <scp><i>1A7*3</i></scp>, and <scp><i>1A9*22</i></scp> genotypes predict severe neutropenia in <scp>FOLFIRI</scp>â€treated metastatic colorectal cancer in two prospective studies in Japan. Cancer Science, 2013, 104, 1662-1669.	3.9	29
58	A Randomized Controlled Trial Comparing the Effects of Sitagliptin and Glimepiride on Endothelial Function and Metabolic Parameters: Sapporo Athero-Incretin Study 1 (SAIS1). PLoS ONE, 2016, 11, e0164255.	2.5	29
59	Initial safety report on the tolerability of modified FOLFOX6 as adjuvant therapy in patients with curatively resected stage II or III colon cancer (JFMC41-1001-C2: JOIN trial). Cancer Chemotherapy and Pharmacology, 2015, 76, 75-84.	2.3	26
60	Clinical significance of left ventricular hypertrophy and changes in left ventricular mass in high-risk hypertensive patients: a subanalysis of the Candesartan Antihypertensive Survival Evaluation in Japan trial. Journal of Hypertension, 2009, 27, 1705-1712.	0.5	25
61	The role of ultrasound imaging in adult patients with testicular torsion: a systematic review and meta-analysis. Journal of Medical Ultrasonics (2001), 2019, 46, 325-334.	1.3	25
62	Comparison of the clinical effectiveness of zanamivir and laninamivir octanoate for children with influenza <scp>A</scp> (<scp>H</scp>3<scp>N</scp>2) and <scp>B</scp> in the 2011â€2012 season. Influenza and Other Respiratory Viruses, 2014, 8, 151-158.	3.4	24
63	Prognosis of Japanese patients with previously untreated metastatic renal cell carcinoma in the era of molecularâ€targeted therapy. Cancer Science, 2015, 106, 618-626.	3.9	24
64	Relative increase in lymphocytes from as early as 1Âmonth predicts improved response to dasatinib in chronic-phase chronic myelogenous leukemia. International Journal of Hematology, 2014, 99, 41-52.	1.6	23
65	Differences in UGT1A1, UGT1A7, and UGT1A9 Polymorphisms between Uzbek and Japanese Populations. Molecular Diagnosis and Therapy, 2014, 18, 333-342.	3.8	23
66	Utility of a Three-Dimensional Printed Pelvic Model for Lateral Pelvic Lymph Node Dissection Education: A Randomized Controlled Trial. Journal of the American College of Surgeons, 2019, 229, 552-559.e3.	0.5	23
67	Pharmacokinetic study of weekly administration dose of paclitaxel in patients with advanced or recurrent gastric cancer in Japan. Gastric Cancer, 2007, 10, 52-57.	5.3	22
68	Multicenter Phase II study of FOLFOX or biweekly XELOX and Erbitux (cetuximab) as first-line therapy in patients with wild-type KRAS/BRAF metastatic colorectal cancer: The FLEET study. BMC Cancer, 2015, 15, 695.	2.6	22
69	Open-label clinical trial of bezafibrate treatment in patients with fatty acid oxidation disorders in Japan. Molecular Genetics and Metabolism Reports, 2018, 15, 55-63.	1.1	22
70	Rate of Reoperation Decreased Significantly After Year 2002 in Patients With Crohn's Disease. Clinical Gastroenterology and Hepatology, 2020, 18, 898-907.e5.	4.4	22
71	Regional Differences in Prevalence of Macrolide Resistance among Pediatric <i>Mycoplasma pneumoniae</i> Infections in Hokkaido, Japan. Japanese Journal of Infectious Diseases, 2016, 69, 186-190.	1.2	21
72	Safety of laparoscopic surgery for colorectal cancer in patients over 80Âyears old: a propensity score matching study. Surgery Today, 2017, 47, 951-958.	1.5	21

#	ARTICLE	IF	CITATIONS
73	Clinical impact of tumor location on the colon cancer survival and recurrence: analyses of pooled data from three large phase <sc>III</sc> randomized clinical trials. Cancer Medicine, 2017, 6, 2523-2530.	2.8	21
74	One-Day Versus Three-Day Dexamethasone in Combination with Palonosetron for the Prevention of Chemotherapy-Induced Nausea and Vomiting: A Systematic Review and Individual Patient Data-Based Meta-Analysis. Oncologist, 2019, 24, 1593-1600.	3.7	21
75	Cardiac Magnetic Resonance Performs Better in the Detection of Functionally Significant Coronary Artery Stenosis Compared to Single-Photon Emission Computed Tomography and Dobutamine Stress Echocardiography. Circulation Journal, 2014, 78, 2468-2476.	1.6	20
76	Prognostic Significance of Peritoneal Metastasis in Stage IV Colorectal Cancer Patients With R0 Resection: A Multicenter, Retrospective Study. Diseases of the Colon and Rectum, 2017, 60, 1041-1049.	1.3	20
77	Is Memorial Sloan-Kettering Cancer Center risk classification appropriate for Japanese patients with metastatic renal cell carcinoma in the cytokine era?. Urologic Oncology: Seminars and Original Investigations, 2013, 31, 1276-1282.	1.6	19
78	SAPPHIRE: a randomised phase II study of planned discontinuation or continuous treatment of oxaliplatin after six cycles of modified FOLFOX6 plus panitumumab in patients with colorectal cancer. European Journal of Cancer, 2019, 119, 158-167.	2.8	19
79	Risk factors for outlet obstruction after laparoscopic surgery and diverting ileostomy for rectal cancer. Surgery Today, 2021, 51, 366-373.	1.5	19
80	Temperament and character profiles of Japanese university students with depressive episodes and ideas of suicide or self-harm: A PHQ-9 screening study. Comprehensive Psychiatry, 2013, 54, 1215-1221.	3.1	18
81	Predictors of rituximab effect on modified Rodnan skin score in systemic sclerosis: a machine-learning analysis of the DesiReSâ€trial. Rheumatology, 2022, 61, 4364-4373.	1.9	18
82	The domain structure of the EORTC QLQâ€STO22 supported by Japanese validation data. Psycho-Oncology, 2008, 17, 474-479.	2.3	17
83	A prospective, multicenter survey on the validity of shorter periendoscopic cessation of antithrombotic agents in Japan. Journal of Gastroenterology, 2017, 52, 50-60.	5.1	17
84	Adaptive Servo-Ventilation Has More Favorable Acute Effects on Hemodynamics Than Continuous Positive Airway Pressure in Patients With Heart Failure. International Heart Journal, 2015, 56, 527-532.	1.0	16
85	Prediction of the preoperative chemoradiotherapy response for rectal cancer by peripheral blood lymphocyte subsets. World Journal of Surgical Oncology, 2015, 13, 30.	1.9	16
86	A Phase II Study of XELOX and Cetuximab as First-Line Therapy in Patients With KRAS Wild Type Metastatic Colorectal Cancer (FLEET2 Study). Clinical Colorectal Cancer, 2016, 15, 329-336.	2.3	16
87	Longitudinal comparison of quality of life after real-time tumor-tracking intensity-modulated radiation therapy and radical prostatectomy in patients with localized prostate cancer. Journal of Radiation Research, 2013, 54, 1095-1101.	1.6	15
88	A novel system for predicting the toxicity of irinotecan based on statistical pattern recognition with UGT1A genotypes. International Journal of Oncology, 2014, 45, 1381-1390.	3.3	15
89	Phase <sc>II</sc> study on hepatic arterial infusion chemotherapy using percutaneous catheter placement techniques for liver metastases from colorectal cancer (JFMC28 study). Asia-Pacific Journal of Clinical Oncology, 2015, 11, 41-48.	1.1	15
90	Hazard rate of tumor recurrence over time in patients with colon cancer: implications for postoperative surveillance from three Japanese Foundation for Multidisciplinary Treatment of Cancer (JFMC) clinical trials. Journal of Cancer, 2017, 8, 4057-4064.	2.5	15

#	ARTICLE	IF	CITATIONS
91	JOIN trial: treatment outcome and recovery status of peripheral sensory neuropathy during a 3-year follow-up in patients receiving modified FOLFOX6 as adjuvant treatment for stage II/III colon cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2019, 84, 1269-1277.	2.3	15
92	Randomized phase 2 study of perampanel for sporadic amyotrophic lateral sclerosis. <i>Journal of Neurology</i> , 2022, 269, 885-896.	3.6	15
93	Randomized controlled Phase III study comparing hepatic arterial infusion with systemic chemotherapy after curative resection for liver metastasis of colorectal carcinoma: JFMC 29â€œ0003. <i>Journal of Cancer Research and Therapeutics</i> , 2017, 13, 84.	0.9	15
94	Multicenter phase II study of second-line cetuximab plus folinic acid/5-fluorouracil/irinotecan (FOLFIRI) in KRAS wild-type metastatic colorectal cancer: the FLIER study. <i>Anticancer Research</i> , 2014, 34, 1967-73.	1.1	15
95	Multicenter randomized phase II clinical trial of oxaliplatin reintroduction as a third- or later-line therapy for metastatic colorectal cancerâ€”biweekly versus standard triweekly XELOX (The ORION) Tj ETQq1 1 0.784314 rgBT/Overl	1.1	15
96	Does a laparoscopic approach attenuate the body weight loss and lean body mass loss observed in open distal gastrectomy for gastric cancer? a single-institution exploratory analysis of the JCOG 0912 phase III trial. <i>Gastric Cancer</i> , 2018, 21, 345-352.	5.3	14
97	Efficacy of Oral Anticancer Agents for Colorectal Cancer. <i>Diseases of the Colon and Rectum</i> , 2006, 49, S82-S91.	1.3	13
98	A feasibility study of sequential paclitaxel and S-1 (PTX/S-1) chemotherapy as postoperative adjuvant chemotherapy for advanced gastric cancer. <i>Gastric Cancer</i> , 2006, 9, 114-119.	5.3	13
99	Long-term effects of candesartan and amlodipine on cardiovascular morbidity and mortality in Japanese high-risk hypertensive patients: the Candesartan Antihypertensive Survival Evaluation in Japan Extension Study (CASE-J Ex). <i>Hypertension Research</i> , 2011, 34, 1295-1301.	2.7	13
100	Optimal predictive value of preoperative serum carcinoembryonic antigen for surgical outcomes in stage I nonâ€œsmall cell lung cancer: Differences according to histology and smoking status. <i>Journal of Surgical Oncology</i> , 2013, 107, 619-624.	1.7	13
101	Efficacy of corticosteroids for cancer-related fatigue: A pilot randomized placebo-controlled trial of advanced cancer patients. <i>Palliative and Supportive Care</i> , 2015, 13, 1301-1308.	1.0	13
102	Capecitabine and oxaliplatin combined with bevacizumab are feasible for treating selected Japanese patients at least 75Â¥years of age with metastatic colorectal cancer. <i>BMC Cancer</i> , 2015, 15, 786.	2.6	13
103	Transcatheter Arterial Embolization with Ethanol Injection in Symptomatic Patients with Enlarged Polycystic Kidneys. <i>Radiology</i> , 2015, 277, 277-285.	7.3	13
104	Statistical challenges for central monitoring in clinical trials: a review. <i>International Journal of Clinical Oncology</i> , 2016, 21, 28-37.	2.2	13
105	Multicenter phase II study of infusional 5-fluorouracil (5-FU), leucovorin, and oxaliplatin, plus biweekly cetuximab as first-line treatment in patients with metastatic colorectal cancer (CELINE trial). <i>Oncology Letters</i> , 2017, 13, 747-753.	1.8	13
106	Detection of atypical data in multicenter clinical trials using unsupervised statistical monitoring. <i>Clinical Trials</i> , 2019, 16, 512-522.	1.6	13
107	Sustained discontinuation of infliximab with a raising-dose strategy after obtaining remission in patients with rheumatoid arthritis: the RRRR study, a randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2020, 79, 94-102.	0.9	13
108	A retrospective review of 10-year trends in general anesthesia for cesarean delivery at a university hospital: the impact of a newly launched team on obstetric anesthesia practice. <i>BMC Health Services Research</i> , 2020, 20, 421.	2.2	13

#	ARTICLE	IF	CITATIONS
109	Oxidized albumin in blood reflects the severity of multiple vascular complications in diabetes mellitus. <i>Metabolism Open</i> , 2020, 6, 100032.	2.9	13
110	Efficacy of Molecular Response at 1 or 3 Months after the Initiation of Dasatinib Treatment Can Predict an Improved Response to Dasatinib in Imatinib-Resistant or Imatinib-Intolerant Japanese Patients with Chronic Myelogenous Leukemia during the Chronic Phase. <i>Journal of Clinical and Experimental Hematopathology</i> : JCEH, 2014, 54, 197-204.	0.8	12
111	Comparison of Different Symptom Assessment Scales for Multiple System Atrophy. <i>Cerebellum</i> , 2016, 15, 190-200.	2.5	12
112	Clinical Signatures of Mucinous and Poorly Differentiated Subtypes of Colorectal Adenocarcinomas by a Propensity Score Analysis of an Independent Patient Database from Three Phase III Trials. <i>Diseases of the Colon and Rectum</i> , 2018, 61, 461-471.	1.3	12
113	Incidence rate and characteristics of symptomatic vitamin D deficiency in children: a nationwide survey in Japan. <i>Endocrine Journal</i> , 2018, 65, 593-599.	1.6	12
114	Influence of Coronary Risk Factors on Coronary Events in Japanese High-Risk Hypertensive Patients - Primary and Secondary Prevention of Ischemic Heart Disease in a Subanalysis of the Candesartan Antihypertensive Survival Evaluation in Japan (CASE-J) Trial -. <i>Circulation Journal</i> , 2011, 75, 2411-2416.	1.6	11
115	Pharmacokinetic dose adjustment of 5-FU in modified FOLFOX7 plus bevacizumab for metastatic colorectal cancer in Japanese patients: a-JUST phase II clinical trial. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 78, 1253-1261.	2.3	11
116	CD133 Expression at the Metastatic Site Predicts Patients' Outcome in Colorectal Cancer with Synchronous Liver Metastasis. <i>Annals of Surgical Oncology</i> , 2016, 23, 1916-1923.	1.5	11
117	Randomized phase II study of cetuximab versus irinotecan and cetuximab in patients with chemo-refractory KRAS codon G13D metastatic colorectal cancer (G13D-study). <i>Cancer Chemotherapy and Pharmacology</i> , 2017, 79, 29-36.	2.3	11
118	Development and Validation of a Prediction Model for Postoperative Peritoneal Metastasis After Curative Resection of Colon Cancer. <i>Annals of Surgical Oncology</i> , 2018, 25, 1366-1373.	1.5	11
119	Meta-analysis of randomized clinical trials in the era of individual patient data sharing. <i>International Journal of Clinical Oncology</i> , 2018, 23, 403-409.	2.2	11
120	Randomized phase III trial comparing surgery alone to UFT+PSK for stage II rectal cancer (JFMC38 trial). <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 81, 65-71.	2.3	11
121	Clinical TNM staging for esophageal, gastric, and colorectal cancers in the era of neoadjuvant therapy: A systematic review of the literature. <i>Annals of Gastroenterological Surgery</i> , 2021, 5, 404-418.	2.4	11
122	Preoperative sarcopenia is a poor prognostic factor in lower rectal cancer patients undergoing neoadjuvant chemoradiotherapy: a retrospective study. <i>International Journal of Clinical Oncology</i> , 2022, 27, 141-153.	2.2	11
123	Multicenter Phase II Trial of Combination Chemotherapy with Weekly Paclitaxel and 5-Fluorouracil for the Treatment of Advanced or Recurrent Gastric Carcinoma. <i>Journal of Chemotherapy</i> , 2007, 19, 444-450.	1.5	10
124	Efficacy of adjuvant chemotherapy using tegafur-based regimen for curatively resected gastric cancer: update of a meta-analysis. <i>International Journal of Clinical Oncology</i> , 2009, 14, 85-89.	2.2	10
125	Long-term effects of candesartan and amlodipine on cardiovascular mortality and morbidity in Japanese high-risk hypertensive patients: Rationale, design, and characteristics of candesartan antihypertensive survival evaluation in Japan extension (CASE-J Ex). <i>Contemporary Clinical Trials</i> , 2009, 30, 97-101.	1.8	10
126	Association between working overtime and psychological stress reactions in elementary and junior high school teachers in Japan: a large-scale cross-sectional study. <i>Industrial Health</i> , 2021, 60, 133-145.	1.0	10

#	ARTICLE	IF	CITATIONS
127	A Randomized Phase II Trial of Preoperative Exercise to Reduce Operative Risk in Gastric Cancer Patients with Metabolic Syndrome: Adjuvant Exercise for General Elective Surgery (AEGES) Study Group. Japanese Journal of Clinical Oncology, 2008, 38, 71-73.	1.3	9
128	A phase I study of palliative chemoradiation therapy with paclitaxel and cisplatin for local symptoms due to an unresectable primary advanced or locally recurrent gastric adenocarcinoma. Cancer Chemotherapy and Pharmacology, 2009, 64, 1071-1077.	2.3	9
129	FOLFIRI Plus Bevacizumab 5 mg/kg Versus 10 mg/kg as Second-line Therapy in Patients with Metastatic Colorectal Cancer Who Have Failed First-line Bevacizumab Plus Oxaliplatin-based Therapy: A Randomized Phase III Study (EAGLE Study). Japanese Journal of Clinical Oncology, 2012, 42, 134-138.	1.3	9
130	Immunohistochemical Validation and Expression Profiling of NKG2D Ligands in a Wide Spectrum of Human Epithelial Neoplasms. Journal of Histochemistry and Cytochemistry, 2015, 63, 217-227.	2.5	9
131	Rapid reduction in <i>BCR-ABL</i> transcript predicts deep molecular response in dasatinib-treated chronic phase chronic myeloid leukaemia patients. European Journal of Haematology, 2018, 100, 27-35.	2.2	9
132	Prognostic value of metastatic lymph node regression grade after neoadjuvant chemoradiotherapy in patients with locally advanced rectal cancer. Surgery, 2019, 166, 1061-1067.	1.9	9
133	Is there diversity among <i>UGT1A1</i> polymorphism in Japan. World Journal of Gastrointestinal Oncology, 2012, 4, 170.	2.0	9
134	Effects of Cardiac Complications on Cardiovascular Events in Japanese High-Risk Hypertensive Patients Subanalysis of the CASE-J Trial. Circulation Journal, 2009, 73, 1080-1085.	1.6	8
135	Relationship Between the Waiting Times for Surgery and Survival in Patients with Gastric Cancer. World Journal of Surgery, 2020, 44, 1209-1215.	1.6	8
136	Intraperitoneal Chemotherapy as Adjuvant or Perioperative Chemotherapy for Patients with Type 4 Scirrhous Gastric Cancer: PHOENIX-GC2 Trial. Journal of Clinical Medicine, 2021, 10, 5666.	2.4	8
137	Prospective phase II trial of second-line FOLFIRI in patients with advanced colorectal cancer including analysis of <i>UGT1A1</i> polymorphisms: FLIGHT 2 study. Anticancer Research, 2014, 34, 195-201.	1.1	8
138	Sleep-Disordered Breathing Is an Independent Risk Factor of Aborted Sudden Cardiac Arrest in Patients With Coronary Artery Spasm. Circulation Journal, 2012, 76, 2204-2210.	1.6	7
139	The COMET Open-label Phase II Study of Neoadjuvant FOLFOX or XELOX Treatment Combined with Molecular Targeting Monoclonal Antibodies in Patients with Resectable Liver Metastasis of Colorectal Cancer. Annals of Surgical Oncology, 2017, 24, 546-553.	1.5	7
140	Clinical Efficacy and Safety of First-Line Dasatinib Therapy and the Relevance of Velocity of <i>BCR-ABL1</i> Transcript Decline for Achievement of Molecular Responses in Newly Diagnosed Chronic-Phase Chronic Myeloid Leukemia: Report from the Juntendo Yamanashi Cooperative Study Group. Oncology, 2018, 94, 85-91.	1.9	7
141	Open-label clinical trial of bezafibrate treatment in patients with fatty acid oxidation disorders in Japan; 2nd report QOL survey. Molecular Genetics and Metabolism Reports, 2019, 20, 100496.	1.1	7
142	Rationale and design of the EMPYREAN study. ESC Heart Failure, 2020, 7, 3134-3141.	3.1	7
143	Treatment using oxaliplatin and S-1 adjuvant chemotherapy for pathological stage III gastric cancer: a multicenter phase II study (TOSA trial) protocol. BMC Cancer, 2018, 18, 186.	2.6	6
144	Risk-Stratified Incidence of Renal Replacement Therapy Initiation: A Longitudinal Analysis Using Medical Claims and Health Checkup Data. Tohoku Journal of Experimental Medicine, 2019, 248, 125-135.	1.2	6

#	ARTICLE	IF	CITATIONS
145	Associations between visceral obesity and renal impairment in health checkup participants: a retrospective cohort study. <i>Clinical and Experimental Nephrology</i> , 2020, 24, 935-945.	1.6	6
146	Sites of blood collection and topical antiseptics associated with contaminated cultures: prospective observational study. <i>Scientific Reports</i> , 2021, 11, 6211.	3.3	6
147	A randomized phase III study of hepatic arterial infusion chemotherapy with 5-fluorouracil and subsequent systemic chemotherapy versus systemic chemotherapy alone for colorectal cancer patients with curatively resected liver metastases (Japanese Foundation for Multidisciplinary) Tj ETQq1 1 0.784314 0.978 BT / Overflock 10	0.9	6
148	Percentage of residual B cells after 2Âweeks of rituximab treatment predicts the improvement of systemic sclerosisâ€associated interstitial lung disease. <i>Journal of Dermatology</i> , 2022, 49, 179-183.	1.2	6
149	Multicenter phase II study of FOLFOX for metastatic colorectal cancer (mCRC) in Japan; SWIFT-1 and 2 study. <i>Hepato-Gastroenterology</i> , 2009, 56, 1346-53.	0.5	6
150	Phase II study of weekly paclitaxel, cisplatin, and 5-fluorouracil for advanced gastric cancer. <i>Gastric Cancer</i> , 2011, 14, 332-338.	5.3	5
151	How to use marginal structural models in randomized trials to estimate the natural direct and indirect effects of therapies mediated by causal intermediates. <i>Clinical Trials</i> , 2011, 8, 277-287.	1.6	5
152	Low dose capecitabine plus weekly paclitaxel in patients with metastatic breast cancer: a multicenter phase II study KBCSG-0609. <i>Cancer Chemotherapy and Pharmacology</i> , 2013, 71, 741-747.	2.3	5
153	Relationship between pre-anesthetic and intra-anesthetic airway resistance in patients undergoing general anesthesia: A prospective observational study. <i>PLoS ONE</i> , 2017, 12, e0172421.	2.5	5
154	Development and validation of a prognostic nomogram for colorectal cancer after radical resection based on individual patient data from three large-scale phase III trials. <i>Oncotarget</i> , 2017, 8, 99150-99160.	1.8	5
155	Pretreatment evaluation of fluorescence resonance energy transferâ€based drug sensitivity test for patients with chronic myelogenous leukemia treated with dasatinib. <i>Cancer Science</i> , 2018, 109, 2256-2265.	3.9	5
156	Development and Validation of a Prediction Model for Organ-Specific Recurrences After Curative Resection of Colon Cancer. <i>Diseases of the Colon and Rectum</i> , 2019, 62, 1043-1054.	1.3	5
157	Efficacy of bezafibrate for preventing myopathic attacks in patients with very long-chain acyl-CoA dehydrogenase deficiency. <i>Brain and Development</i> , 2021, 43, 214-219.	1.1	5
158	Adjuvant chemotherapy with tegafur/uracil (UFT) for gastric cancer. A meta-analysis of centrally randomized clinical trials. <i>Journal of Clinical Oncology</i> , 2006, 24, 4033-4033.	1.6	5
159	Relative Increase of Lymphocytes As Early As 1 Month After Initiation of Dasatinib Is a Reliable Predictor for Achieving Complete Molecular Response At 12 Months in Chronic Phase CML Patients Treated with Dasatinib. <i>Blood</i> , 2012, 120, 691-691.	1.4	5
160	Higher Hemoglobin A1c After Discharge Is an Independent Predictor of Adverse Outcomes in Patients With Acute Coronary Syndromeâ€â€ Findings From the PACIFIC Registry â€. <i>Circulation Journal</i> , 2016, 80, 1607-1614.	1.6	4
161	Two-by-two cross-over study to evaluate agreement between versions of a quantitative coronary analysis system (QAngio XA). <i>International Journal of Cardiovascular Imaging</i> , 2017, 33, 779-787.	1.5	4
162	mFOLFOX6 Plus Panitumumab Versus 5-FU/LV Plus Panitumumab After Six Cycles of Frontline mFOLFOX6 Plus Panitumumab: A Randomized Phase II Study of Patients With Unresectable or Advanced/Recurrent, RAS Wild-type Colorectal Carcinoma (SAPPHIRE)â€Study Design andÂRationale. <i>Clinical Colorectal Cancer</i> , 2017, 16, 154-157.e1.	2.3	4

#	ARTICLE	IF	CITATIONS
163	Remission induction by Raising the dose of Remicade in RA (RRRR) study: Rationale and study protocol for a randomized controlled trial comparing for sustained clinical remission after discontinuation of infliximab in patients with rheumatoid arthritis. Contemporary Clinical Trials Communications, 2017, 8, 49-54.	1.1	4
164	A phase II trial to evaluate the efficacy of panitumumab combined with fluorouracil-based chemotherapy for metastatic colorectal cancer: the PF trial. Cancer Chemotherapy and Pharmacology, 2018, 81, 829-838.	2.3	4
165	Impact of Residual Stenosis on the Angiographic Edge Restenosis of a Second-Generation Drug-Eluting Stent. International Heart Journal, 2019, 60, 1050-1060.	1.0	4
166	Lymph node retrieval after colorectal cancer surgery: a comparative study of the efficacy between the conventional manual method and a new fat dissolution method. Surgery Today, 2020, 50, 726-733.	1.5	4
167	Rapid decrease of serum surfactant protein- α levels predicts the reactivity of rituximab therapy in systemic sclerosis-associated interstitial lung disease. Journal of Dermatology, 2020, 47, 796-800.	1.2	4
168	Efficacy and Safety of Polaprezinc (Zinc Compound) on Zinc Deficiency: A Systematic Review and Dose-Response Meta-Analysis of Randomized Clinical Trials Using Individual Patient Data. Nutrients, 2020, 12, 1128.	4.1	4
169	FOLFIRI plus bevacizumab (bev) as second-line therapy in patients (pts) with metastatic colorectal cancer (mCRC) who have failed first-line bev plus oxaliplatin-based therapy: The randomized phase III EAGLE study.. Journal of Clinical Oncology, 2013, 31, 3516-3516.	1.6	4
170	Association between polymorphisms in EGFR and tumor response during cetuximab and oxaliplatin-based combination therapy in metastatic colorectal cancer: Analysis of data from two clinical trials. Oncology Letters, 2019, 18, 4555-4562.	1.8	4
171	Association between hospital treatment volume and survival of women with gynecologic malignancy in Japan: a JSOG tumor registry-based data extraction study. Journal of Gynecologic Oncology, 2021, 33, .	2.2	4
172	Bi-weekly Capecitabine-Oxaliplatin (XELOX) plus Bevacizumab as First-line Treatment of Metastatic Colorectal Cancer -The PHOENIX Trial. Anticancer Research, 2016, 36, 3437-43.	1.1	4
173	Factors associated with compliance and non-compliance by physicians in a large-scale randomized clinical trial. Trials, 2006, 7, 26.	1.6	3
174	Impact of Left Ventricular Hypertrophy on the Time-Course of Renal Function in Hypertensive Patients - A Subanalysis of the CASE-J Trial -. Circulation Journal, 2010, 74, 2132-2138.	1.6	3
175	Plasma matrix metalloproteinase-3 correlates with the clinical severity in men with multiple system atrophy. Neurology and Clinical Neuroscience, 2013, 1, 69-77.	0.4	3
176	Lymph node retrieval after dissolution of surrounding adipose tissue for pathological examination of colorectal cancer. Oncology Letters, 2017, 15, 2495-2500.	1.8	3
177	A phase II study of bevacizumab and irinotecan plus alternate-day S-1 as a second-line therapy in patients with metastatic colorectal cancer: the AIRS study. Cancer Chemotherapy and Pharmacology, 2018, 81, 1035-1041.	2.3	3
178	A multicenter single-arm Phase II clinical trial of second-line FOLFIRI plus panitumumab after first-line treatment with FOLFOX plus panitumumab for initial RAS wild-type colorectal cancer with evaluation of circulating tumor DNA: A protocol study. Oncology Letters, 2018, 17, 1980-1985.	1.8	3
179	Biweekly S-1 plus oxaliplatin (SOX) reintroduction in previously treated metastatic colorectal cancer patients (ORION 2 study): a phase II study to evaluate the efficacy and safety. International Journal of Clinical Oncology, 2019, 24, 836-841.	2.2	3
180	SAMIT: Preliminary safety data from a 2x2 factorial randomized phase III trial to investigate weekly paclitaxel (PTX) followed by oral fluoropyrimidines (FPs) versus FPs alone as adjuvant chemotherapy in patients (pts) with gastric cancer.. Journal of Clinical Oncology, 2011, 29, 4017-4017.	1.6	3

#	ARTICLE	IF	CITATIONS
181	XELOX with bevacizumab in elderly patients age 75 or older with metastatic colorectal cancer: Results of a planned interim analysis for multicenter phase II ASCA study.. Journal of Clinical Oncology, 2013, 31, 502-502.	1.6	3
182	Bayesian adaptive randomization design incorporating propensity scoreâ€¢matched historical controls. Pharmaceutical Statistics, 2022, , .	1.3	3
183	Geographic difference in advanced gastric cancer prevalence and chemotherapy treatment results: could it really be an independent prognostic factor?. Gastric Cancer, 2012, 15, 232-234.	5.3	2
184	A prospective analysis of clinical efficacy and safety in chronic myeloid leukemiaâ€¢chronic phase patients with imatinib resistance or intolerance as evaluated using <scp>E</scp>uropean <scp>L</scp>eukemia<scp>N</scp>et 2013 criteria. European Journal of Haematology, 2015, 95, 558-565.	2.2	2
185	Risk factors associated with recurrence by peritoneal dissemination or paraaortic lymph node metastasis after curative surgery in patients with colorectal cancer. Annals of Cancer Research and Therapy, 2016, 24, 58-59.	0.3	2
186	Optimal Interval for 18F-FDG-PET After Chemoradiotherapy for Rectal Cancer. Clinical Colorectal Cancer, 2018, 17, e163-e170.	2.3	2
187	Efficacy of CapeOX plus Cetuximab Treatment as a First-Line Therapy for Patients with Extended RAS/BRAF/PIK3CA Wild-Type Advanced or Metastatic Colorectal Cancer. Journal of Cancer, 2018, 9, 4092-4098.	2.5	2
188	Reference value for expiratory time constant calculated from the maximal expiratory flow-volume curve. BMC Pulmonary Medicine, 2019, 19, 208.	2.0	2
189	Multivariable prediction model to estimate the probability of restenosis at proximal edge after 2nd-generation drug-eluting-stent implantation: development and internal validation using a quantitative coronary angiography from the post-marketing surveillance studies of everolimus-eluting stent in Japan. Cardiovascular Intervention and Therapeutics. 2021, 36, 190-197.	2.3	2
190	Focused Assessment with Sonography for Trauma (FAST) training for first-year resident physicians at a university hospital in Japan: A longitudinal, observational study. SAGE Open Medicine, 2021, 9, 205031212110443.	1.8	2
191	Serum TARC Levels in Patients with Systemic Sclerosis: Clinical Association with Interstitial Lung Disease. Journal of Clinical Medicine, 2021, 10, 660.	2.4	2
192	Phase II study of trastuzumab with irinotecan in HER2-positive metastatic or advanced gastric cancer patients previously treated with trastuzumab and failed: HGCSG 1201/OGSG1205.. Journal of Clinical Oncology, 2017, 35, 151-151.	1.6	2
193	Treatment outcome and recovery from peripheral sensory neuropathy during a 3-year follow-up in patients receiving modified FOLFOX6 as adjuvant treatment for stage II/III colon cancer (JFMC41-1001-C2:) Tj ETQq16l 0.784314 rgB		
194	Truth-telling to Patients with Advanced Cancer: A Comparison between Cancer center, University Hospitals, General City Hospitals, and Private General Practitioners in Japan. Annals of Cancer Research and Therapy, 2006, 14, 1-11.	0.3	2
195	Comparison of clinic pathological characters and survival between right- and left-side colon cancer. Annals of Cancer Research and Therapy, 2016, 24, 62-63.	0.3	2
196	Tolerability study of adjuvant modified FOLFOX6 treatment in curatively resected stage II/III colon cancer (JFMC41-1001-C2: JOIN trial).. Journal of Clinical Oncology, 2014, 32, 530-530.	1.6	2
197	Factors influencing the number of retrieved lymph nodes after colorectal resection: a retrospective study from a single institute. International Journal of Clinical and Experimental Pathology, 2018, 11, 1694-1700.	0.5	2
198	Improvement in Surgical Outcomes Using 3D Printed Models for Lateral Pelvic Lymph Node Dissection in Rectal Cancer. Diseases of the Colon and Rectum, 2021, Publish Ahead of Print, .	1.3	2

#	ARTICLE	IF	CITATIONS
199	Increased Red Blood Cell Distribution Width in the First Year after Diagnosis Predicts Worsening of Systemic Sclerosis-Associated Interstitial Lung Disease at 5 Years: A Pilot Study. <i>Diagnostics</i> , 2021, 11, 2274.	2.6	2
200	Phase II Study of Continued Trastuzumab Plus Irinotecan in Patients with HER2-positive Gastric Cancer Previously Treated with Trastuzumab (HGCSG 1201). <i>Oncologist</i> , 2022, 27, 340-e374.	3.7	2
201	Laparoscopic Colectomy: A Risk Factor for Postoperative Peritoneal Metastasis. <i>Clinical Colorectal Cancer</i> , 2022, 21, e205-e212.	2.3	2
202	Multicentre Phase II Study of XELOX with Bevacizumab in Late-stage Elderly Patients with Unresectable Advanced/Recurrent Colorectal Cancer: An ASCA Study. <i>Japanese Journal of Clinical Oncology</i> , 2011, 41, 134-138.	1.3	1
203	Optimal Dose Period for Indisetrone Tablets for Preventing Chemotherapy-Induced Nausea and Vomiting with Modified FOLFOX6: A Randomized Pilot Study. <i>Chemotherapy</i> , 2012, 58, 439-444.	1.6	1
204	Preservation of Deep Inferior Epigastric Artery at Kidney Transplantation Prevents Atrophy of Lower Rectus Abdominis Muscle. <i>Transplantation</i> , 2012, 93, 1013-1016.	1.0	1
205	Long-term effects of antihypertensive therapy on cardiovascular events and new-onset diabetes mellitus in high-risk hypertensive patients in Japan. <i>Journal of Hypertension</i> , 2018, 36, 1921-1928.	0.5	1
206	Phase II study of an oxaliplatin-based regimen for relapsed colon cancer patients treated with oxaliplatin-based adjuvant chemotherapy (INSPIRE study). <i>Cancer Chemotherapy and Pharmacology</i> , 2021, 87, 665-672.	2.3	1
207	Serum Calponin 3 Levels in Patients with Systemic Sclerosis: Possible Association with Skin Sclerosis and Arthralgia. <i>Journal of Clinical Medicine</i> , 2021, 10, 280.	2.4	1
208	SAMIT: A phase III randomized clinical trial of adjuvant paclitaxel followed by oral fluorinated pyrimidines for locally advanced gastric cancer.. <i>Journal of Clinical Oncology</i> , 2013, 31, LBA4002-LBA4002.	1.6	1
209	Clinical features of colorectal mucinous and poorly differentiated adenocarcinomas; study concept of a propensity score analysis in a pooled data of 5530 patients. <i>Annals of Cancer Research and Therapy</i> , 2016, 24, 52-53.	0.3	1
210	The relation between postoperative surgical complications and colorectal cancer survival. <i>Annals of Cancer Research and Therapy</i> , 2016, 24, 54-55.	0.3	1
211	Lymph node retrieval using fat dissolution technique after surgical resection of colorectal cancer: a protocol paper. <i>Annals of Cancer Research and Therapy</i> , 2017, 25, 90-91.	0.3	1
212	Effect of antispasmodic drugs on endoscopic ultrasound/endoscopic ultrasound-guided fine-needle aspiration: A multicenter randomized controlled trial. <i>Endoscopic Ultrasound</i> , 2014, 3, 167.	1.5	1
213	Clinical trial comparing UFT-PSK combination adjuvant therapy and surgery-alone for Stage II rectal cancer. <i>Annals of Cancer Research and Therapy</i> , 2017, 25, 15-16.	0.3	1
214	COVID-19 pandemic highlighted the importance of telemedicine in the collagen disease of systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 160-160.	0.8	1
215	THU0231â€¦The efficacy of tacrolimus in patients with interstitial lung diseases complicated with polymyositis or dermatomyositis. <i>Annals of the Rheumatic Diseases</i> , 2013, 71, 233.2-233.	0.9	0
216	The Potential Diagnostic Role of the Number of Ultrasonographic Characteristics for Patients with Thyroid Nodules Evaluated as Bethesda IÃ¢â‚¬â€œV. <i>Frontiers in Oncology</i> , 2014, 4, 261.	2.8	0

#	ARTICLE	IF	CITATIONS
217	AB0293â€¦High Serum Levels of Anti-Cyclic Citrullinated Peptide Antibody and Matrix Metalloproteinase-3 at the Time of Diagnosis of Rheumatoid Arthritis Are Possible Predictors of Future Initiation of Biological Agents. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 991.1-991.	0.9	0
218	APOLLON study: a phase I/II study for the safety and efficacy of panitumumab in combination with TAS-102 for patients with RAS wild-type metastatic colorectal cancer refractory to standard chemotherapy. <i>Annals of Oncology</i> , 2016, 27, vi201.	1.2	0
219	Evaluation of Surrogate Endpoints Using a Meta-Analysis Approach with Individual Patient Data: Summary of a Gastric Cancer Meta-Analysis Project. , 2017, , 179-192.		0
220	HGCSG 1201: Phase II study of trastuzumab with irinotecan in HER2-positive metastatic or advanced gastric cancer patients previously treated with trastuzumab. <i>Annals of Oncology</i> , 2017, 28, iii37.	1.2	0
221	P4566Predictors for angiographic edge restenosis of 2nd generation drug-eluting stents: Integrated angiographic analyses from the post-marketing surveillance studies of everolimus-eluting stents in Japan. <i>European Heart Journal</i> , 2018, 39, .	2.2	0
222	Multicenter phase Ib/II study of biweekly TAS-102 with bevacizumab combination for patients with metastatic colorectal cancer refractory to standard therapies (BiTS study) - Trial in progress. <i>Annals of Oncology</i> , 2018, 29, v83.	1.2	0
223	A phase I/II study of panitumumab combined with TAS-102 in patients (pts) with RAS wild-type (wt) metastatic colorectal cancer (mCRC) refractory to standard chemotherapy: APOLLON study. <i>Annals of Oncology</i> , 2018, 29, v71.	1.2	0
224	Response to: â€˜Comment on â€˜Sustained discontinuation of infliximab with a raising-dose strategy after obtaining remission in patients with rheumatoid arthritis: the RRRR study, a randomised controlled trialâ€™ by Tanaka <i>et al</i>â€™ by Berkhout <i>et al</i>. <i>Annals of the Rheumatic Diseases</i> , 2021, 80, e173-e173.	0.9	0
225	Impact of follow-up on generalized pairwise comparisons for estimating the irinotecan benefit in advanced/metastatic gastric cancer. <i>Contemporary Clinical Trials</i> , 2021, 105, 106400.	1.8	0
226	ANTIREFLUX CERVICAL ESOPHAGOGASTRICANASTOMOSIS IN THE SURGERY OF ESOPHAGEAL CANCER. <i>Annals of Cancer Research and Therapy</i> , 2010, 18, 59-64.	0.3	0
227	PROSPECTIVE STUDY ON THE EFFECT OF TOREMIFENE IN PATIENTS WITH ADJUVANT ANASTROZOLE FAILURE IN POSTMENOPAUSAL BREAST CANCER. <i>Annals of Cancer Research and Therapy</i> , 2011, 19, 9-14.	0.3	0
228	A novel prognostic classification for overall survival in Japanese patients with untreated metastatic renal cell carcinoma.. <i>Journal of Clinical Oncology</i> , 2011, 29, e15051-e15051.	1.6	0
229	Dasatinib Induces a Higher Molecular Response in Japanese Patients with Chronic Myeloid Leukemia After Imatinib Failure Than in Western Populations: Kanto CML Study. <i>Blood</i> , 2011, 118, 4448-4448.	1.4	0
230	Adjuvant paclitaxel followed by oral fluoropyrimidines for gastric cancer: Safety data of the factorial phase III SAMIT trial.. <i>Journal of Clinical Oncology</i> , 2012, 30, 72-72.	1.6	0
231	Retrospective cohort study on the risk factors of admission due to serious adverse events during S-1 (tegafur, gimeracil, oteracil potassium) containing chemotherapy for gastric cancer.. <i>Journal of Clinical Oncology</i> , 2012, 30, 141-141.	1.6	0
232	FOLFIRI plus 5 mg/kg of bevacizumab versus 10 mg/kg as second-line therapy in patients with metastatic colorectal cancer who have failed first-line bevacizumab plus oxaliplatin-based therapy: A randomized phase III study (EAGLE Study).. <i>Journal of Clinical Oncology</i> , 2012, 30, TPS3642-TPS3642.	1.6	0
233	Multicenter phase II study ofÂFOLFOX or biweekly XELOX and cetuximab as first-line treatment in patients with wild-type KRAS/BRAF metastatic colorectal cancer (mCRC) (FLEET study).. <i>Journal of Clinical Oncology</i> , 2013, 31, 463-463.	1.6	0
234	The Prospective Analysis Of Clinical Efficacy and Adverse Events In Chronic Myeloid Leukemia In Chronic Phase (CML-CP) Patients With Imatinib Resistance Or Intolerance, Evaluated By European Leukemia Net (ELN) 2013 Criteria. <i>Blood</i> , 2013, 122, 4027-4027.	1.4	0

#	ARTICLE	IF	CITATIONS
235	Randomized controlled trial on the skin toxicity of panitumumab in third-line treatment of KRAS wild-type metastatic colorectal cancer: HGCSG1001 (Japanese Skin Toxicity Evaluation Protocol with) Tj ETQq1 1 0z84314 rgBT /Overl	1.6	0
236	Matched pair analysis to examine the effects of a planned preoperative exercise program in early gastric cancer patients with metabolic syndrome to reduce operative risk: The Adjuvant Exercise for General Elective Surgery (AEGES) study group.. Journal of Clinical Oncology, 2014, 32, 166-166.	1.6	0
237	Phase II study of 5-FU dose-adjusted mFOLFOX7 plus bevacizumab in patients with metastatic colorectal cancer (AJUST study).. Journal of Clinical Oncology, 2014, 32, 592-592.	1.6	0
238	Randomized controlled trial on the skin toxicity of panitumumab in third-line treatment of KRAS wild-type metastatic colorectal cancer: HGSG1001 (Japanese Skin Toxicity Evaluation Protocol with) Tj ETQq0 0 0 rgBT /Overlck 10 Tf 5 32, 3587-3587.	1.6	0
239	Prognosis of Japanese patients with metastatic renal cell carcinoma who underwent subsequent systemic therapy due to prior failure of VEGF-TKI therapy.. Journal of Clinical Oncology, 2015, 33, e15615-e15615.	1.6	0
240	Velocity of Early BCR-ABL Transcript Elimination As an Optimized Predictor of Deep Molecular Response in Chronic Myeloid Leukemia Patients in Chronic Phase on Treatment with Dasatinib. Blood, 2015, 126, 4052-4052.	1.4	0
241	Hazard rate of tumor recurrence over time: a pooled-analysis of three clinical trials with fluoropyrimidine-based adjuvant chemotherapy for colorectal cancer achieved by the Japanese Foundation for Multidisciplinary Treatment of Cancer (JFMC). Annals of Cancer Research and Therapy, 2016, 24, 60-61.	0.3	0
242	Prognostic Nomogram for Colorectal Cancer after Radical Resection in Japanese population. Annals of Cancer Research and Therapy, 2016, 24, 56-57.	0.3	0
243	Phase II study of irinotecan and bevacizumab plus alternate day S-1 as second-line treatment in patients with metastatic colorectal cancer.. Journal of Clinical Oncology, 2016, 34, 749-749.	1.6	0
244	First-Line Dasatinib Treatment of CML-CP Leads to Earlier Achievement of MMR and MR4.5 with High Safety. Blood, 2016, 128, 3102-3102.	1.4	0
245	Combined analysis of three randomized phase III trials comparing S-1 monotherapy and S-1 combination therapy for first-line treatment of advanced gastric cancer.. Journal of Clinical Oncology, 2017, 35, 49-49.	1.6	0
246	Systematic review and individual patient data based meta-analysis of palonosetron trials for chemotherapy induced nausea and vomiting.. Journal of Clinical Oncology, 2017, 35, e21688-e21688.	1.6	0
247	Trial in progress: A phase II study of second-line FOLFIRI plus panitumumab after first-line FOLFOX plus panitumumab for RAS wild-type colorectal cancer with evaluation of circulating tumor DNA.. Journal of Clinical Oncology, 2018, 36, TPS886-TPS886.	1.6	0
248	A Study of Persistence of Anti-HBs Antibodies after Vaccination with Recombinant Hepatitis B Vaccine in Healthcare Workers. Japanese Journal of Environmental Infections, 2018, 33, 117-122.	0.1	0
249	Risk factors for paraaortic lymph-node recurrence in colorectal cancer. Annals of Cancer Research and Therapy, 2019, 27, 52-56.	0.3	0
250	COVID-19 pandemic highlighted the importance of telemedicine in the collagen disease of systemic sclerosis. Clinical and Experimental Rheumatology, 2021, 39 Suppl 131, 160.	0.8	0
251	Analysis of the radial and ulnar arteries using ultrasound and a vascular visualization device in children. Paediatric Anaesthesia, 2022, , .	1.1	0