

Koji Oba

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/224072/koji-oba-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

232
papers

4,327
citations

32
h-index

58
g-index

254
ext. papers

5,201
ext. citations

3.7
avg, IF

5.07
L-index

#	Paper	IF	Citations
232	Predictors of Rituximab Effect on Modified Rodnan Skin Score in Systemic Sclerosis: a machine learning analysis of the DESIRES trial.. <i>Rheumatology</i> , 2022 ,	3.9	1
231	Preoperative sarcopenia is a poor prognostic factor in lower rectal cancer patients undergoing neoadjuvant chemoradiotherapy: a retrospective study. <i>International Journal of Clinical Oncology</i> , 2021 , 27, 141	4.2	0
230	Intraperitoneal Chemotherapy as Adjuvant or Perioperative Chemotherapy for Patients with Type 4 Scirrhus Gastric Cancer: PHOENIX-GC2 Trial. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	1
229	Association between working overtime and psychological stress reactions in elementary and junior high school teachers in Japan: a large-scale cross-sectional study. <i>Industrial Health</i> , 2021 ,	2.5	1
228	Randomized phase 2 study of perampanel for sporadic amyotrophic lateral sclerosis. <i>Journal of Neurology</i> , 2021 , 1	5.5	5
227	Impact of follow-up on generalized pairwise comparisons for estimating the irinotecan benefit in advanced/metastatic gastric cancer. <i>Contemporary Clinical Trials</i> , 2021 , 105, 106400	2.3	
226	Safety and efficacy of rituximab in systemic sclerosis (DESIRES): a double-blind, investigator-initiated, randomised, placebo-controlled trial. <i>Lancet Rheumatology, The</i> , 2021 , 3, e489-e497	14.2	20
225	Response to: Comment on Sustained discontinuation of infliximab with a raising-dose strategy after obtaining remission in patients with rheumatoid arthritis: the RRRR study, a randomised controlled trial by Tanaka and Berkhout. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, e173	2.4	
224	Multivariable prediction model to estimate the probability of restenosis at proximal edge after 2nd-generation drug-eluting-stent implantation: development and internal validation using a quantitative coronary angiography from the post-marketing surveillance studies of	2.5	0
223	Risk factors for outlet obstruction after laparoscopic surgery and diverting ileostomy for rectal cancer. <i>Surgery Today</i> , 2021 , 51, 366-373	3	3
222	Efficacy of bezafibrate for preventing myopathic attacks in patients with very long-chain acyl-CoA dehydrogenase deficiency. <i>Brain and Development</i> , 2021 , 43, 214-219	2.2	0
221	Focused Assessment with Sonography for Trauma (FAST) training for first-year resident physicians at a university hospital in Japan: A longitudinal, observational study. <i>SAGE Open Medicine</i> , 2021 , 9, 20503121211044367	2.4	
220	Phase II study of an oxaliplatin-based regimen for relapsed colon cancer patients treated with oxaliplatin-based adjuvant chemotherapy (INSPIRE study). <i>Cancer Chemotherapy and Pharmacology</i> , 2021 , 87, 665-672	3.5	1
219	Clinical TNM staging for esophageal, gastric, and colorectal cancers in the era of neoadjuvant therapy: A systematic review of the literature. <i>Annals of Gastroenterological Surgery</i> , 2021 , 5, 404-418	4.3	1
218	Sites of blood collection and topical antiseptics associated with contaminated cultures: prospective observational study. <i>Scientific Reports</i> , 2021 , 11, 6211	4.9	1
217	COVID-19 pandemic highlighted the importance of telemedicine in the collagen disease of systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39 Suppl 131, 160	2.2	
216	COVID-19 pandemic highlighted the importance of telemedicine in the collagen disease of systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39, 160-160	2.2	1

215	Rapid decrease of serum surfactant protein-D levels predicts the reactivity of rituximab therapy in systemic sclerosis-associated interstitial lung disease. <i>Journal of Dermatology</i> , 2020 , 47, 796-800	1.6	2
214	A retrospective review of 10-year trends in general anesthesia for cesarean delivery at a university hospital: the impact of a newly launched team on obstetric anesthesia practice. <i>BMC Health Services Research</i> , 2020 , 20, 421	2.9	3
213	Oxidized albumin in blood reflects the severity of multiple vascular complications in diabetes mellitus. <i>Metabolism Open</i> , 2020 , 6, 100032	2.8	10
212	Rationale and design of the EMPYREAN study. <i>ESC Heart Failure</i> , 2020 , 7, 3134-3141	3.7	4
211	Associations between visceral obesity and renal impairment in health checkup participants: a retrospective cohort study. <i>Clinical and Experimental Nephrology</i> , 2020 , 24, 935-945	2.5	3
210	Relationship Between the Waiting Times for Surgery and Survival in Patients with Gastric Cancer. <i>World Journal of Surgery</i> , 2020 , 44, 1209-1215	3.3	3
209	Efficacy and Safety of Polaprezinc (Zinc Compound) on Zinc Deficiency: A Systematic Review and Dose-Response Meta-Analysis of Randomized Clinical Trials Using Individual Patient Data. <i>Nutrients</i> , 2020 , 12,	6.7	2
208	Lymph node retrieval after colorectal cancer surgery: a comparative study of the efficacy between the conventional manual method and a new fat dissolution method. <i>Surgery Today</i> , 2020 , 50, 726-733	3	2
207	Sustained discontinuation of infliximab with a raising-dose strategy after obtaining remission in patients with rheumatoid arthritis: the RRRR study, a randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 94-102	2.4	7
206	Comprehensive Comparative Analysis of Prognostic Value of Systemic Inflammatory Biomarkers for Patients with Stage II/III Colon Cancer. <i>Annals of Surgical Oncology</i> , 2020 , 27, 844-852	3.1	16
205	Rate of Reoperation Decreased Significantly After Year 2002 in Patients With Crohn's Disease. <i>Clinical Gastroenterology and Hepatology</i> , 2020 , 18, 898-907.e5	6.9	11
204	Radiomics Approach Outperforms Diameter Criteria for Predicting Pathological Lateral Lymph Node Metastasis After Neoadjuvant (Chemo)Radiotherapy in Advanced Low Rectal Cancer. <i>Annals of Surgical Oncology</i> , 2020 , 27, 4273-4283	3.1	10
203	SAPPHERE: a randomised phase II study of planned discontinuation or continuous treatment of oxaliplatin after six cycles of modified FOLFOX6 plus panitumumab in patients with colorectal cancer. <i>European Journal of Cancer</i> , 2019 , 119, 158-167	7.5	9
202	Impact of Residual Stenosis on the Angiographic Edge Restenosis of a Second-Generation Drug-Eluting Stent. <i>International Heart Journal</i> , 2019 , 60, 1050-1060	1.8	3
201	JOIN trial: treatment outcome and recovery status of peripheral sensory neuropathy during a 3-year follow-up in patients receiving modified FOLFOX6 as adjuvant treatment for stage II/III colon cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2019 , 84, 1269-1277	3.5	11
200	One-Day Versus Three-Day Dexamethasone in Combination with Palonosetron for the Prevention of Chemotherapy-Induced Nausea and Vomiting: A Systematic Review and Individual Patient Data-Based Meta-Analysis. <i>Oncologist</i> , 2019 , 24, 1593-1600	5.7	8
199	Immunogenomic profiles associated with response to neoadjuvant chemoradiotherapy in patients with rectal cancer. <i>British Journal of Surgery</i> , 2019 , 106, 1381-1392	5.3	27
198	The role of ultrasound imaging in adult patients with testicular torsion: a systematic review and meta-analysis. <i>Journal of Medical Ultrasonics (2001)</i> , 2019 , 46, 325-334	1.4	12

197	Biweekly S-1 plus oxaliplatin (SOX) reintroduction in previously treated metastatic colorectal cancer patients (ORION 2 study): a phase II study to evaluate the efficacy and safety. <i>International Journal of Clinical Oncology</i> , 2019 , 24, 836-841	4.2	3
196	Association of blood mercury levels during pregnancy with infant birth size by blood selenium levels in the Japan Environment and Children's Study: A prospective birth cohort. <i>Environment International</i> , 2019 , 125, 418-429	12.9	14
195	Open-label clinical trial of bezafibrate treatment in patients with fatty acid oxidation disorders in Japan; 2nd report QOL survey. <i>Molecular Genetics and Metabolism Reports</i> , 2019 , 20, 100496	1.8	4
194	Detection of atypical data in multicenter clinical trials using unsupervised statistical monitoring. <i>Clinical Trials</i> , 2019 , 16, 512-522	2.2	5
193	Prognostic value of metastatic lymph node regression grade after neoadjuvant chemoradiotherapy in patients with locally advanced rectal cancer. <i>Surgery</i> , 2019 , 166, 1061-1067	3.6	3
192	Utility of a Three-Dimensional Printed Pelvic Model for Lateral Pelvic Lymph Node Dissection Education: A Randomized Controlled Trial. <i>Journal of the American College of Surgeons</i> , 2019 , 229, 552-559	4.4	6
191	Risk-Stratified Incidence of Renal Replacement Therapy Initiation: A Longitudinal Analysis Using Medical Claims and Health Checkup Data. <i>Tohoku Journal of Experimental Medicine</i> , 2019 , 248, 125-135	2.4	4
190	Association between polymorphisms in EGFR and tumor response during cetuximab and oxaliplatin-based combination therapy in metastatic colorectal cancer: Analysis of data from two clinical trials. <i>Oncology Letters</i> , 2019 , 18, 4555-4562	2.6	0
189	Risk factors for paraaortic lymph-node recurrence in colorectal cancer. <i>Annals of Cancer Research and Therapy</i> , 2019 , 27, 52-56	0.2	
188	Reference value for expiratory time constant calculated from the maximal expiratory flow-volume curve. <i>BMC Pulmonary Medicine</i> , 2019 , 19, 208	3.5	1
187	Development and Validation of a Prediction Model for Organ-Specific Recurrences After Curative Resection of Colon Cancer. <i>Diseases of the Colon and Rectum</i> , 2019 , 62, 1043-1054	3.1	1
186	A multicenter single-arm Phase II clinical trial of second-line FOLFIRI plus panitumumab after first-line treatment with FOLFOX plus panitumumab for initial wild-type colorectal cancer with evaluation of circulating tumor DNA: A protocol study. <i>Oncology Letters</i> , 2019 , 17, 1980-1985	2.6	2
185	Development and Validation of a Prediction Model for Postoperative Peritoneal Metastasis After Curative Resection of Colon Cancer. <i>Annals of Surgical Oncology</i> , 2018 , 25, 1366-1373	3.1	8
184	A phase II trial to evaluate the efficacy of panitumumab combined with fluorouracil-based chemotherapy for metastatic colorectal cancer: the PF trial. <i>Cancer Chemotherapy and Pharmacology</i> , 2018 , 81, 829-838	3.5	3
183	Clinical effectiveness of four neuraminidase inhibitors (oseltamivir, zanamivir, laninamivir, and peramivir) for children with influenza A and B in the 2014-2015 to 2016-2017 influenza seasons in Japan. <i>Journal of Infection and Chemotherapy</i> , 2018 , 24, 449-457	2.2	21
182	A phase II study of bevacizumab and irinotecan plus alternate-day S-1 as a second-line therapy in patients with metastatic colorectal cancer: the AIRS study. <i>Cancer Chemotherapy and Pharmacology</i> , 2018 , 81, 1035-1041	3.5	2
181	Pattern of programmed cell death-ligand 1 expression and CD8-positive T-cell infiltration before and after chemoradiotherapy in rectal cancer. <i>European Journal of Cancer</i> , 2018 , 91, 11-20	7.5	36
180	Vasoactive-inotropic score as a predictor of morbidity and mortality in adults after cardiac surgery with cardiopulmonary bypass. <i>Journal of Anesthesia</i> , 2018 , 32, 167-173	2.2	45

179	Meta-analysis of randomized clinical trials in the era of individual patient data sharing. <i>International Journal of Clinical Oncology</i> , 2018 , 23, 403-409	4.2	9
178	Risk Factors for Peritoneal Recurrence in Stage II to III Colon Cancer. <i>Diseases of the Colon and Rectum</i> , 2018 , 61, 803-808	3.1	22
177	Open-label clinical trial of bezafibrate treatment in patients with fatty acid oxidation disorders in Japan. <i>Molecular Genetics and Metabolism Reports</i> , 2018 , 15, 55-63	1.8	14
176	Clinical Signatures of Mucinous and Poorly Differentiated Subtypes of Colorectal Adenocarcinomas by a Propensity Score Analysis of an Independent Patient Database from Three Phase III Trials. <i>Diseases of the Colon and Rectum</i> , 2018 , 61, 461-471	3.1	9
175	Randomized phase III trial comparing surgery alone to UFT + PSK for stage II rectal cancer (JFMC38 trial). <i>Cancer Chemotherapy and Pharmacology</i> , 2018 , 81, 65-71	3.5	4
174	Rapid reduction in BCR-ABL1 transcript predicts deep molecular response in dasatinib-treated chronic-phase chronic myeloid leukaemia patients. <i>European Journal of Haematology</i> , 2018 , 100, 27-35	3.8	8
173	Treatment using oxaliplatin and S-1 adjuvant chemotherapy for pathological stage III gastric cancer: a multicenter phase II study (TOSA trial) protocol. <i>BMC Cancer</i> , 2018 , 18, 186	4.8	5
172	Incidence rate and characteristics of symptomatic vitamin D deficiency in children: a nationwide survey in Japan. <i>Endocrine Journal</i> , 2018 , 65, 593-599	2.9	7
171	Long-term effects of antihypertensive therapy on cardiovascular events and new-onset diabetes mellitus in high-risk hypertensive patients in Japan. <i>Journal of Hypertension</i> , 2018 , 36, 1921-1928	1.9	1
170	Lymph node retrieval after dissolution of surrounding adipose tissue for pathological examination of colorectal cancer. <i>Oncology Letters</i> , 2018 , 15, 2495-2500	2.6	2
169	Factors influencing the number of retrieved lymph nodes after colorectal resection: a retrospective study from a single institute. <i>International Journal of Clinical and Experimental Pathology</i> , 2018 , 11, 1694-1700	1.7	1
168	A randomized phase III study of hepatic arterial infusion chemotherapy with 5-fluorouracil and subsequent systemic chemotherapy versus systemic chemotherapy alone for colorectal cancer patients with curatively resected liver metastases (Japanese Foundation for Multidisciplinary Treatment of Cancer Study 6201). <i>Journal of Clinical Oncology</i> , 2018 , 36, 2711-2719	1.2	5
167	Trial in progress: A phase II study of second-line FOLFIRI plus panitumumab after first-line FOLFOX plus panitumumab for RAS wild-type colorectal cancer with evaluation of circulating tumor DNA.. <i>Journal of Clinical Oncology</i> , 2018 , 36, TPS886-TPS886	2.2	
166	A Study of Persistence of Anti-HBs Antibodies after Vaccination with Recombinant Hepatitis B Vaccine in Healthcare Workers. <i>Japanese Journal of Environmental Infections</i> , 2018 , 33, 117-122	0.1	
165	Does a laparoscopic approach attenuate the body weight loss and lean body mass loss observed in open distal gastrectomy for gastric cancer? a single-institution exploratory analysis of the JCOG 0912 phase III trial. <i>Gastric Cancer</i> , 2018 , 21, 345-352	7.6	8
164	Optimal Interval for F-FDG-PET After Chemoradiotherapy for Rectal Cancer. <i>Clinical Colorectal Cancer</i> , 2018 , 17, e163-e170	3.8	0
163	Clinical Efficacy and Safety of First-Line Dasatinib Therapy and the Relevance of Velocity of BCR-ABL1 Transcript Decline for Achievement of Molecular Responses in Newly Diagnosed Chronic-Phase Chronic Myeloid Leukemia: Report from the Juntendo Yamanashi Cooperative Study Group. <i>Oncology</i> , 2018 , 94, 85-91	3.6	5
162	Efficacy of CapeOX plus Cetuximab Treatment as a First-Line Therapy for Patients with Extended RAS/BRAF/PIK3CA Wild-Type Advanced or Metastatic Colorectal Cancer. <i>Journal of Cancer</i> , 2018 , 9, 4092-4098	4.5	1

161	Pretreatment evaluation of fluorescence resonance energy transfer-based drug sensitivity test for patients with chronic myelogenous leukemia treated with dasatinib. <i>Cancer Science</i> , 2018 , 109, 2256-2265	6.9	4
160	A prospective, multicenter survey on the validity of shorter periendoscopic cessation of antithrombotic agents in Japan. <i>Journal of Gastroenterology</i> , 2017 , 52, 50-60	6.9	14
159	Two-by-two cross-over study to evaluate agreement between versions of a quantitative coronary analysis system (QAngio XA). <i>International Journal of Cardiovascular Imaging</i> , 2017 , 33, 779-787	2.5	3
158	Safety of laparoscopic surgery for colorectal cancer in patients over 80 years old: a propensity score matching study. <i>Surgery Today</i> , 2017 , 47, 951-958	3	18
157	Does postoperative morbidity worsen the oncological outcome after radical surgery for gastrointestinal cancers? A systematic review of the literature. <i>Annals of Gastroenterological Surgery</i> , 2017 , 1, 11-23	4.3	49
156	Multicenter phase II study of infusional 5-fluorouracil (5-FU), leucovorin, and oxaliplatin, plus biweekly cetuximab as first-line treatment in patients with metastatic colorectal cancer (CELINE trial). <i>Oncology Letters</i> , 2017 , 13, 747-753	2.6	9
155	mFOLFOX6 Plus Panitumumab Versus 5-FU/LV Plus Panitumumab After Six Cycles of Frontline mFOLFOX6 Plus Panitumumab: A Randomized Phase II Study of Patients With Unresectable or Advanced/Recurrent, RAS Wild-type Colorectal Carcinoma (SAPPHIRE)-Study Design and Rationale. <i>Clinical Colorectal Cancer</i> , 2017 , 16, 154-157.e1	3.8	3
154	Clinical impact of tumor location on the colon cancer survival and recurrence: analyses of pooled data from three large phase III randomized clinical trials. <i>Cancer Medicine</i> , 2017 , 6, 2523-2530	4.8	18
153	Evaluation of Surrogate Endpoints Using a Meta-Analysis Approach with Individual Patient Data: Summary of a Gastric Cancer Meta-Analysis Project 2017 , 179-192		
152	Relationship between pre-anesthetic and intra-anesthetic airway resistance in patients undergoing general anesthesia: A prospective observational study. <i>PLoS ONE</i> , 2017 , 12, e0172421	3.7	5
151	Therapeutic efficacy of azithromycin, clarithromycin, minocycline and tosufloxacin against macrolide-resistant and macrolide-sensitive <i>Mycoplasma pneumoniae</i> pneumonia in pediatric patients. <i>PLoS ONE</i> , 2017 , 12, e0173635	3.7	35
150	Development and validation of a prognostic nomogram for colorectal cancer after radical resection based on individual patient data from three large-scale phase III trials. <i>Oncotarget</i> , 2017 , 8, 99150-99160	3.3	4
149	Hazard rate of tumor recurrence over time in patients with colon cancer: implications for postoperative surveillance from three Japanese Foundation for Multidisciplinary Treatment of Cancer (JFMC) clinical trials. <i>Journal of Cancer</i> , 2017 , 8, 4057-4064	4.5	10
148	Remission induction by Raising the dose of Remicade in RA (RRRR) study: Rationale and study protocol for a randomized controlled trial comparing for sustained clinical remission after discontinuation of infliximab in patients with rheumatoid arthritis. <i>Contemporary Clinical Trials Communications</i> , 2017 , 8, 49-54	1.8	4
147	Prognostic Significance of Peritoneal Metastasis in Stage IV Colorectal Cancer Patients With R0 Resection: A Multicenter, Retrospective Study. <i>Diseases of the Colon and Rectum</i> , 2017 , 60, 1041-1049	3.1	12
146	Impact of postoperative complications on the colorectal cancer survival and recurrence: analyses of pooled individual patients data from three large phase III randomized trials. <i>Cancer Medicine</i> , 2017 , 6, 1573-1580	4.8	42
145	The COMET Open-label Phase II Study of Neoadjuvant FOLFOX or XELOX Treatment Combined with Molecular Targeting Monoclonal Antibodies in Patients with Resectable Liver Metastasis of Colorectal Cancer. <i>Annals of Surgical Oncology</i> , 2017 , 24, 546-553	3.1	6
144	Randomized phase II study of cetuximab versus irinotecan and cetuximab in patients with chemo-refractory KRAS codon G13D metastatic colorectal cancer (G13D-study). <i>Cancer Chemotherapy and Pharmacology</i> , 2017 , 79, 29-36	3.5	8

143	Phase II study of trastuzumab with irinotecan in HER2-positive metastatic or advanced gastric cancer patients previously treated with trastuzumab and failed: HGCSG 1201/OGSG1205.. <i>Journal of Clinical Oncology</i> , 2017 , 35, 151-151	2.2	2
142	Treatment outcome and recovery from peripheral sensory neuropathy during a 3-year follow-up in patients receiving modified FOLFOX6 as adjuvant treatment for stage II/III colon cancer (JFMC41-1001-C2: JOIN Trial).. <i>Journal of Clinical Oncology</i> , 2017 , 35, 703-703	2.2	1
141	Randomized controlled Phase III study comparing hepatic arterial infusion with systemic chemotherapy after curative resection for liver metastasis of colorectal carcinoma: JFMC 29-0003. <i>Journal of Cancer Research and Therapeutics</i> , 2017 , 13, 84-90	1.2	10
140	Lymph node retrieval using fat dissolution technique after surgical resection of colorectal cancer: a protocol paper. <i>Annals of Cancer Research and Therapy</i> , 2017 , 25, 90-91	0.2	1
139	Clinical trial comparing UFT-PSK combination adjuvant therapy and surgery-alone for Stage II rectal cancer. <i>Annals of Cancer Research and Therapy</i> , 2017 , 25, 15-16	0.2	1
138	Combined analysis of three randomized phase III trials comparing S-1 monotherapy and S-1 combination therapy for first-line treatment of advanced gastric cancer.. <i>Journal of Clinical Oncology</i> , 2017 , 35, 49-49	2.2	
137	Systematic review and individual patient data based meta-analysis of palonosetron trials for chemotherapy induced nausea and vomiting.. <i>Journal of Clinical Oncology</i> , 2017 , 35, e21688-e21688	2.2	
136	Comparison of Different Symptom Assessment Scales for Multiple System Atrophy. <i>Cerebellum</i> , 2016 , 15, 190-200	4.3	10
135	Serum myostatin levels are independently associated with skeletal muscle wasting in patients with heart failure. <i>International Journal of Cardiology</i> , 2016 , 220, 483-7	3.2	25
134	Higher Hemoglobin A1c After Discharge Is an Independent Predictor of Adverse Outcomes in Patients With Acute Coronary Syndrome - Findings From the PACIFIC Registry. <i>Circulation Journal</i> , 2016 , 80, 1607-14	2.9	3
133	Pharmacokinetic dose adjustment of 5-FU in modified FOLFOX7 plus bevacizumab for metastatic colorectal cancer in Japanese patients: a-JUST phase II clinical trial. <i>Cancer Chemotherapy and Pharmacology</i> , 2016 , 78, 1253-1261	3.5	9
132	Statistical evaluation of surrogate endpoints with examples from cancer clinical trials. <i>Biometrical Journal</i> , 2016 , 58, 104-32	1.5	54
131	Statistical challenges for central monitoring in clinical trials: a review. <i>International Journal of Clinical Oncology</i> , 2016 , 21, 28-37	4.2	8
130	Multicenter randomized phase II clinical trial of oxaliplatin reintroduction as a third- or later-line therapy for metastatic colorectal cancer-biweekly versus standard triweekly XELOX (The ORION Study). <i>International Journal of Clinical Oncology</i> , 2016 , 21, 566-72	4.2	10
129	CD133 Expression at the Metastatic Site Predicts PatientsPOutcome in Colorectal Cancer with Synchronous Liver Metastasis. <i>Annals of Surgical Oncology</i> , 2016 , 23, 1916-23	3.1	10
128	Does remnant gastric cancer really differ from primary gastric cancer? A systematic review of the literature by the Task Force of Japanese Gastric Cancer Association. <i>Gastric Cancer</i> , 2016 , 19, 339-349	7.6	26
127	Clinical features of colorectal mucinous and poorly differentiated adenocarcinomas; study concept of a propensity score analysis in a pooled data of 5530 patients. <i>Annals of Cancer Research and Therapy</i> , 2016 , 24, 52-53	0.2	1
126	The relation between postoperative surgical complications and colorectal cancer survival. <i>Annals of Cancer Research and Therapy</i> , 2016 , 24, 54-55	0.2	1

125	Comparison of clinic pathological characters and survival between right- and left-side colon cancer. <i>Annals of Cancer Research and Therapy</i> , 2016 , 24, 62-63	0.2	2
124	Hazard rate of tumor recurrence over time: a pooled-analysis of three clinical trials with fluoropyrimidine-based adjuvant chemotherapy for colorectal cancer achieved by the Japanese Foundation for Multidisciplinary Treatment of Cancer (JFMC). <i>Annals of Cancer Research and Therapy</i> , 2016 , 24, 68-81	0.2	
123	Prognostic Nomogram for Colorectal Cancer after Radical Resection in Japanese population. <i>Annals of Cancer Research and Therapy</i> , 2016 , 24, 56-57	0.2	
122	Phase II study of irinotecan and bevacizumab plus alternate day S-1 as second-line treatment in patients with metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 749-749	2.2	
121	First-Line Dasatinib Treatment of CML-CP Leads to Earlier Achievement of MMR and MR4.5 with High Safety. <i>Blood</i> , 2016 , 128, 3102-3102	2.2	
120	Regional Differences in Prevalence of Macrolide Resistance among Pediatric Mycoplasma pneumoniae Infections in Hokkaido, Japan. <i>Japanese Journal of Infectious Diseases</i> , 2016 , 69, 186-90	2.7	18
119	Risk factors associated with recurrence by peritoneal dissemination or paraaortic lymph node metastasis after curative surgery in patients with colorectal cancer. <i>Annals of Cancer Research and Therapy</i> , 2016 , 24, 58-59	0.2	2
118	A Randomized Controlled Trial Comparing the Effects of Sitagliptin and Glimepiride on Endothelial Function and Metabolic Parameters: Sapporo Athero-Incretin Study 1 (SAIS1). <i>PLoS ONE</i> , 2016 , 11, e0164255	2.7	23
117	A Phase II Study of XELOX and Cetuximab as First-Line Therapy in Patients With KRAS Wild Type Metastatic Colorectal Cancer (FLEET2 Study). <i>Clinical Colorectal Cancer</i> , 2016 , 15, 329-336	3.8	14
116	Comparison of Targeted vs Random Biopsies for Surveillance of Ulcerative Colitis-Associated Colorectal Cancer. <i>Gastroenterology</i> , 2016 , 151, 1122-1130	13.3	117
115	Multicenter randomized controlled study to assess the effect of prophylactic clipping on post-polypectomy delayed bleeding. <i>Digestive Endoscopy</i> , 2016 , 28, 570-6	3.7	44
114	Bi-weekly Capecitabine-Oxaliplatin (XELOX) plus Bevacizumab as First-line Treatment of Metastatic Colorectal Cancer -The PHOENIX Trial. <i>Anticancer Research</i> , 2016 , 36, 3437-43	2.3	4
113	Prediction of the preoperative chemoradiotherapy response for rectal cancer by peripheral blood lymphocyte subsets. <i>World Journal of Surgical Oncology</i> , 2015 , 13, 30	3.4	13
112	A prospective analysis of clinical efficacy and safety in chronic myeloid leukemia-chronic phase patients with imatinib resistance or intolerance as evaluated using European LeukemiaNet 2013 criteria. <i>European Journal of Haematology</i> , 2015 , 95, 558-65	3.8	2
111	Immunohistochemical validation and expression profiling of NKG2D ligands in a wide spectrum of human epithelial neoplasms. <i>Journal of Histochemistry and Cytochemistry</i> , 2015 , 63, 217-27	3.4	6
110	Transcatheter Arterial Embolization with Ethanol Injection in Symptomatic Patients with Enlarged Polycystic Kidneys. <i>Radiology</i> , 2015 , 277, 277-85	20.5	9
109	FOLFIRI plus bevacizumab as second-line therapy in patients with metastatic colorectal cancer after first-line bevacizumab plus oxaliplatin-based therapy: the randomized phase III EAGLE study. <i>Annals of Oncology</i> , 2015 , 26, 1427-33	10.3	35
108	Initial safety report on the tolerability of modified FOLFOX6 as adjuvant therapy in patients with curatively resected stage II or III colon cancer (JFMC41-1001-C2: JOIN trial). <i>Cancer Chemotherapy and Pharmacology</i> , 2015 , 76, 75-84	3.5	23

107	Effect of a stylet on a histological specimen in EUS-guided fine-needle tissue acquisition by using 22-gauge needles: a multicenter, prospective, randomized, controlled trial. <i>Gastrointestinal Endoscopy</i> , 2015 , 82, 837-844.e1	5.2	31
106	Randomized controlled trial on the skin toxicity of panitumumab in Japanese patients with metastatic colorectal cancer: HGCSG1001 study; J-STEPP. <i>Future Oncology</i> , 2015 , 11, 617-27	3.6	45
105	The efficacy of tacrolimus in patients with interstitial lung diseases complicated with polymyositis or dermatomyositis. <i>Rheumatology</i> , 2015 , 54, 39-44	3.9	78
104	Efficacy of corticosteroids for cancer-related fatigue: A pilot randomized placebo-controlled trial of advanced cancer patients. <i>Palliative and Supportive Care</i> , 2015 , 13, 1301-8	2.5	9
103	Prognosis of Japanese patients with previously untreated metastatic renal cell carcinoma in the era of molecular-targeted therapy. <i>Cancer Science</i> , 2015 , 106, 618-26	6.9	17
102	Open-label, randomized, comparative, phase III study on effects of reducing steroid use in combination with Palonosetron. <i>Cancer Science</i> , 2015 , 106, 891-5	6.9	32
101	Multicenter Phase II study of FOLFOX or biweekly XELOX and Erbitux (cetuximab) as first-line therapy in patients with wild-type KRAS/BRAF metastatic colorectal cancer: The FLEET study. <i>BMC Cancer</i> , 2015 , 15, 695	4.8	20
100	Capecitabine and oxaliplatin combined with bevacizumab are feasible for treating selected Japanese patients at least 75 years of age with metastatic colorectal cancer. <i>BMC Cancer</i> , 2015 , 15, 786	4.8	12
99	AB0293 High Serum Levels of Anti-Cyclic Citrullinated Peptide Antibody and Matrix Metalloproteinase-3 at the Time of Diagnosis of Rheumatoid Arthritis Are Possible Predictors of Future Initiation of Biological Agents. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 991.1-991	2.4	
98	Adaptive Servo-Ventilation Has More Favorable Acute Effects on Hemodynamics Than Continuous Positive Airway Pressure in Patients With Heart Failure. <i>International Heart Journal</i> , 2015 , 56, 527-32	1.8	10
97	Serum brain-derived neurotrophic factor level predicts adverse clinical outcomes in patients with heart failure. <i>Journal of Cardiac Failure</i> , 2015 , 21, 300-6	3.3	21
96	Phase II study on hepatic arterial infusion chemotherapy using percutaneous catheter placement techniques for liver metastases from colorectal cancer (JFMC28 study). <i>Asia-Pacific Journal of Clinical Oncology</i> , 2015 , 11, 41-8	1.9	12
95	A Comparison of the Effects of the GLP-1 Analogue Liraglutide and Insulin Glargine on Endothelial Function and Metabolic Parameters: A Randomized, Controlled Trial Sapporo Athero-Incretin Study 2 (SAIS2). <i>PLoS ONE</i> , 2015 , 10, e0135854	3.7	29
94	Prognosis of Japanese patients with metastatic renal cell carcinoma who underwent subsequent systemic therapy due to prior failure of VEGF-TKI therapy.. <i>Journal of Clinical Oncology</i> , 2015 , 33, e15615-e15615	2.2	15615
93	Velocity of Early BCR-ABL Transcript Elimination As an Optimized Predictor of Deep Molecular Response in Chronic Myeloid Leukemia Patients in Chronic Phase on Treatment with Dasatinib. <i>Blood</i> , 2015 , 126, 4052-4052	2.2	
92	Differences in UGT1A1, UGT1A7, and UGT1A9 polymorphisms between Uzbek and Japanese populations. <i>Molecular Diagnosis and Therapy</i> , 2014 , 18, 333-42	4.5	18
91	Matched pair analysis to examine the effects of a planned preoperative exercise program in early gastric cancer patients with metabolic syndrome to reduce operative risk: the Adjuvant Exercise for General Elective Surgery (AEGES) study group. <i>Annals of Surgical Oncology</i> , 2014 , 21, 2044-50	3.1	46
90	Long-term outcome in Japanese patients with lupus nephritis. <i>Lupus</i> , 2014 , 23, 1124-32	2.6	29

89	Sequential paclitaxel followed by tegafur and uracil (UFT) or S-1 versus UFT or S-1 monotherapy as adjuvant chemotherapy for T4a/b gastric cancer (SAMIT): a phase 3 factorial randomised controlled trial. <i>Lancet Oncology, The</i> , 2014 , 15, 886-93	21.7	84
88	Diamine oxidase as a marker of intestinal mucosal injury and the effect of soluble dietary fiber on gastrointestinal tract toxicity after intravenous 5-fluorouracil treatment in rats. <i>Medical Molecular Morphology</i> , 2014 , 47, 100-7	2.3	57
87	High and low negative pressure suction techniques in EUS-guided fine-needle tissue acquisition by using 25-gauge needles: a multicenter, prospective, randomized, controlled trial. <i>Gastrointestinal Endoscopy</i> , 2014 , 80, 1030-7.e1	5.2	64
86	Clinical significance of serum tumor markers for gastric cancer: a systematic review of literature by the Task Force of the Japanese Gastric Cancer Association. <i>Gastric Cancer</i> , 2014 , 17, 26-33	7.6	260
85	A novel system for predicting the toxicity of irinotecan based on statistical pattern recognition with UGT1A genotypes. <i>International Journal of Oncology</i> , 2014 , 45, 1381-90	4.4	13
84	Cardiac magnetic resonance performs better in the detection of functionally significant coronary artery stenosis compared to single-photon emission computed tomography and dobutamine stress echocardiography. <i>Circulation Journal</i> , 2014 , 78, 2468-76	2.9	19
83	Efficacy of molecular response at 1 or 3 months after the initiation of dasatinib treatment can predict an improved response to dasatinib in imatinib-resistant or imatinib-intolerant Japanese patients with chronic myelogenous leukemia during the chronic phase. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2014 , 54, 197-204	1.9	8
82	The potential diagnostic role of the number of ultrasonographic characteristics for patients with thyroid nodules evaluated as Bethesda I-v. <i>Frontiers in Oncology</i> , 2014 , 4, 261	5.3	
81	Prognostic significance of epithelial-mesenchymal transition-related markers in extrahepatic cholangiocarcinoma: comprehensive immunohistochemical study using a tissue microarray. <i>British Journal of Cancer</i> , 2014 , 111, 1363-72	8.7	45
80	The influence of childhood abuse, adult stressful life events and temperaments on depressive symptoms in the nonclinical general adult population. <i>Journal of Affective Disorders</i> , 2014 , 158, 101-7	6.6	56
79	Comparison of the clinical effectiveness of zanamivir and laninamivir octanoate for children with influenza A(H3N2) and B in the 2011-2012 season. <i>Influenza and Other Respiratory Viruses</i> , 2014 , 8, 151-8	5.6	19
78	Relative increase in lymphocytes from as early as 1 month predicts improved response to dasatinib in chronic-phase chronic myelogenous leukemia. <i>International Journal of Hematology</i> , 2014 , 99, 41-52	2.3	21
77	Randomized controlled trial on the skin toxicity of panitumumab in third-line treatment of KRAS wild-type metastatic colorectal cancer: HGSG1001 (Japanese Skin Toxicity Evaluation Protocol with Panitumumab: J-STEPP) Additional analysis of antitumor efficacy.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 3587-3587	2.2	
76	Effect of antispasmodic drugs on endoscopic ultrasound/endoscopic ultrasound-guided fine-needle aspiration: A multicenter randomized controlled trial. <i>Endoscopic Ultrasound</i> , 2014 , 3, 167-73	3.6	1
75	Randomized controlled trial on the skin toxicity of panitumumab in third-line treatment of KRAS wild-type metastatic colorectal cancer: HGCSG1001 (Japanese Skin Toxicity Evaluation Protocol with Panitumumab; J-STEPP).. <i>Journal of Clinical Oncology</i> , 2014 , 32, 541-541	2.2	
74	Matched pair analysis to examine the effects of a planned preoperative exercise program in early gastric cancer patients with metabolic syndrome to reduce operative risk: The Adjuvant Exercise for General Elective Surgery (AEGES) study group.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 166-166	2.2	
73	Phase II study of 5-FU dose-adjusted mFOLFOX7 plus bevacizumab in patients with metastatic colorectal cancer (AJUST study).. <i>Journal of Clinical Oncology</i> , 2014 , 32, 592-592	2.2	
72	Tolerability study of adjuvant modified FOLFOX6 treatment in curatively resected stage II/III colon cancer (JFMC41-1001-C2: JOIN trial).. <i>Journal of Clinical Oncology</i> , 2014 , 32, 530-530	2.2	2

71	Prospective phase II trial of second-line FOLFIRI in patients with advanced colorectal cancer including analysis of UGT1A1 polymorphisms: FLIGHT 2 study. <i>Anticancer Research</i> , 2014 , 34, 195-201	2.3	6
70	Multicenter phase II study of second-line cetuximab plus folinic acid/5-fluorouracil/irinotecan (FOLFIRI) in KRAS wild-type metastatic colorectal cancer: the FLIER study. <i>Anticancer Research</i> , 2014 , 34, 1967-73	2.3	13
69	Low dose capecitabine plus weekly paclitaxel in patients with metastatic breast cancer: a multicenter phase II study KBCSG-0609. <i>Cancer Chemotherapy and Pharmacology</i> , 2013 , 71, 741-7	3.5	5
68	Optimal predictive value of preoperative serum carcinoembryonic antigen for surgical outcomes in stage I non-small cell lung cancer: differences according to histology and smoking status. <i>Journal of Surgical Oncology</i> , 2013 , 107, 619-24	2.8	11
67	Reduced-intensity vs myeloablative conditioning allogeneic hematopoietic SCT for patients aged over 45 years with ALL in remission: a study from the Adult ALL Working Group of the Japan Society for Hematopoietic Cell Transplantation (JSHCT). <i>Bone Marrow Transplantation</i> , 2013 , 48, 1389-94	4.4	23
66	Progression-free survival as a surrogate for overall survival in advanced/recurrent gastric cancer trials: a meta-analysis. <i>Journal of the National Cancer Institute</i> , 2013 , 105, 1667-70	9.7	58
65	Role of chemotherapy for advanced/recurrent gastric cancer: an individual-patient-data meta-analysis. <i>European Journal of Cancer</i> , 2013 , 49, 1565-77	7.5	119
64	Is Memorial Sloan-Kettering Cancer Center risk classification appropriate for Japanese patients with metastatic renal cell carcinoma in the cytokine era?. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 1276-82	2.8	19
63	Plasma matrix metalloproteinase-3 correlates with the clinical severity in men with multiple system atrophy. <i>Neurology and Clinical Neuroscience</i> , 2013 , 1, 69-77	0.3	0
62	Temperament and character profiles of Japanese university students with depressive episodes and ideas of suicide or self-harm: a PHQ-9 screening study. <i>Comprehensive Psychiatry</i> , 2013 , 54, 1215-21	7.3	15
61	Disease-free survival as a surrogate for overall survival in adjuvant trials of gastric cancer: a meta-analysis. <i>Journal of the National Cancer Institute</i> , 2013 , 105, 1600-7	9.7	102
60	Decreased serum brain-derived neurotrophic factor levels are correlated with exercise intolerance in patients with heart failure. <i>International Journal of Cardiology</i> , 2013 , 168, e142-4	3.2	30
59	Phase II trial of paclitaxel and cisplatin as neoadjuvant chemotherapy for locally advanced gastric cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2013 , 71, 1309-14	3.5	37
58	Longitudinal comparison of quality of life after real-time tumor-tracking intensity-modulated radiation therapy and radical prostatectomy in patients with localized prostate cancer. <i>Journal of Radiation Research</i> , 2013 , 54, 1095-101	2.4	13
57	UGT1A1*6, 1A7*3, and 1A9*22 genotypes predict severe neutropenia in FOLFIRI-treated metastatic colorectal cancer in two prospective studies in Japan. <i>Cancer Science</i> , 2013 , 104, 1662-9	6.9	24
56	Effects of KIR ligand incompatibility on clinical outcomes of umbilical cord blood transplantation without ATG for acute leukemia in complete remission. <i>Blood Cancer Journal</i> , 2013 , 3, e164	7	31
55	Acute hemodynamic effects of adaptive servo-ventilation in patients with heart failure. <i>Circulation Journal</i> , 2013 , 77, 1214-20	2.9	33
54	THU0231 The efficacy of tacrolimus in patients with interstitial lung diseases complicated with polymyositis or dermatomyositis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 71, 233.2-233	2.4	

53	FOLFIRI plus bevacizumab (bev) as second-line therapy in patients (pts) with metastatic colorectal cancer (mCRC) who have failed first-line bev plus oxaliplatin-based therapy: The randomized phase III EAGLE study.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 3516-3516	2.2	4
52	SAMIT: A phase III randomized clinical trial of adjuvant paclitaxel followed by oral fluorinated pyrimidines for locally advanced gastric cancer.. <i>Journal of Clinical Oncology</i> , 2013 , 31, LBA4002-LBA4002	2.2	1
51	XELOX with bevacizumab in elderly patients age 75 or older with metastatic colorectal cancer: Results of a planned interim analysis for multicenter phase II ASCA study.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 502-502	2.2	2
50	Multicenter phase II study of FOLFOX or biweekly XELOX and cetuximab as first-line treatment in patients with wild-type KRAS/BRAF metastatic colorectal cancer (mCRC) (FLEET study).. <i>Journal of Clinical Oncology</i> , 2013 , 31, 463-463	2.2	
49	The Prospective Analysis Of Clinical Efficacy and Adverse Events In Chronic Myeloid Leukemia In Chronic Phase (CML-CP) Patients With Imatinib Resistance Or Intolerance, Evaluated By European Leukemia Net (ELN) 2013 Criteria. <i>Blood</i> , 2013 , 122, 4027-4027	2.2	
48	A new prognostic classification for overall survival in Asian patients with previously untreated metastatic renal cell carcinoma. <i>Cancer Science</i> , 2012 , 103, 1695-700	6.9	29
47	Sleep-disordered breathing is an independent risk factor of aborted sudden cardiac arrest in patients with coronary artery spasm. <i>Circulation Journal</i> , 2012 , 76, 2204-10	2.9	5
46	FOLFIRI plus bevacizumab 5 mg/kg versus 10 mg/kg as second-line therapy in patients with metastatic colorectal cancer who have failed first-line bevacizumab plus oxaliplatin-based therapy: a randomized phase III study (EAGLE Study). <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 134-8	2.8	9
45	Optimal dose period for indisetron tablets for preventing chemotherapy-induced nausea and vomiting with modified FOLFOX6: a randomized pilot study. <i>Chemotherapy</i> , 2012 , 58, 439-44	3.2	
44	Preservation of deep inferior epigastric artery at kidney transplantation prevents atrophy of lower rectus abdominis muscle. <i>Transplantation</i> , 2012 , 93, 1013-6	1.8	1
43	Is there diversity among UGT1A1 polymorphism in Japan?. <i>World Journal of Gastrointestinal Oncology</i> , 2012 , 4, 170-5	3.4	9
42	FOLFIRI plus 5 mg/kg of bevacizumab versus 10 mg/kg as second-line therapy in patients with metastatic colorectal cancer who have failed first-line bevacizumab plus oxaliplatin-based therapy: A randomized phase III study (EAGLE Study).. <i>Journal of Clinical Oncology</i> , 2012 , 30, TPS3642-TPS3642	2.2	
41	Relative Increase of Lymphocytes As Early As 1 Month After Initiation of Dasatinib Is a Reliable Predictor for Achieving Complete Molecular Response At 12 Months in Chronic Phase CML Patients Treated with Dasatinib. <i>Blood</i> , 2012 , 120, 691-691	2.2	1
40	Influence of coronary risk factors on coronary events in Japanese high-risk hypertensive patients. - Primary and secondary prevention of ischemic heart disease in a subanalysis of the Candesartan Antihypertensive Survival Evaluation in Japan (CASE-J) trial-. <i>Circulation Journal</i> , 2011 , 75, 2411-6	2.9	9
39	Assessing clinical benefit response in the treatment of gastric malignant ascites with non-measurable lesions: a multicenter phase II trial of paclitaxel for malignant ascites secondary to advanced/recurrent gastric cancer. <i>Gastric Cancer</i> , 2011 , 14, 81-90	7.6	29
38	Phase II study of weekly paclitaxel, cisplatin, and 5-fluorouracil for advanced gastric cancer. <i>Gastric Cancer</i> , 2011 , 14, 332-8	7.6	4
37	Long-term effects of candesartan and amlodipine on cardiovascular morbidity and mortality in Japanese high-risk hypertensive patients: the Candesartan Antihypertensive Survival Evaluation in Japan Extension Study (CASE-J Ex). <i>Hypertension Research</i> , 2011 , 34, 1295-301	4.7	12
36	How to use marginal structural models in randomized trials to estimate the natural direct and indirect effects of therapies mediated by causal intermediates. <i>Clinical Trials</i> , 2011 , 8, 277-87	2.2	5

35	Multicentre phase II study of XELOX with bevacizumab in late-stage elderly patients with unresectable advanced/recurrent colorectal cancer: an ASCA study. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 134-8	2.8	1
34	Prospective phase II study of FOLFIRI for mCRC in Japan, including the analysis of UGT1A1 28/6 polymorphisms. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 477-82	2.8	43
33	SAMIT: Preliminary safety data from a 2x2 factorial randomized phase III trial to investigate weekly paclitaxel (PTX) followed by oral fluoropyrimidines (FPs) versus FPs alone as adjuvant chemotherapy in patients (pts) with gastric cancer.. <i>Journal of Clinical Oncology</i> , 2011 , 29, 4017-4017	2.2	3
32	PROSPECTIVE STUDY ON THE EFFECT OF TOREMIFENE IN PATIENTS WITH ADJUVANT ANASTROZOLE FAILURE IN POSTMENOPAUSAL BREAST CANCER. <i>Annals of Cancer Research and Therapy</i> , 2011 , 19, 9-14	0.2	
31	Dasatinib Induces a Higher Molecular Response in Japanese Patients with Chronic Myeloid Leukemia After Imatinib Failure Than in Western Populations: Kanto CML Study. <i>Blood</i> , 2011 , 118, 4448-4448	2.2	48
30	Benefit of adjuvant chemotherapy for resectable gastric cancer: a meta-analysis. <i>JAMA - Journal of the American Medical Association</i> , 2010 , 303, 1729-37	27.4	586
29	Role of diabetes and obesity in outcomes of the candesartan antihypertensive survival evaluation in Japan (CASE-J) trial. <i>Hypertension Research</i> , 2010 , 33, 600-6	4.7	28
28	Is pulse pressure a predictor of new-onset diabetes in high-risk hypertensive patients?: a subanalysis of the Candesartan Antihypertensive Survival Evaluation in Japan (CASE-J) trial. <i>Diabetes Care</i> , 2010 , 33, 1122-7	14.6	43
27	Impact of left ventricular hypertrophy on the time-course of renal function in hypertensive patients: a subanalysis of the CASE-J trial <i>Circulation Journal</i> , 2010 , 74, 2132-8	2.9	3
26	Patterns of drug use and HIV-related risk behaviors among incarcerated people in a prison in Iran. <i>Journal of Urban Health</i> , 2010 , 87, 603-16	5.8	29
25	ANTIREFLUX CERVICAL ESOPHAGOGASTRICANASTOMOSIS IN THE SURGERY OF ESOPHAGEAL CANCER. <i>Annals of Cancer Research and Therapy</i> , 2010 , 18, 59-64	0.2	
24	Relationship between the achieved blood pressure and the incidence of cardiovascular events in Japanese hypertensive patients with complications: a sub-analysis of the CASE-J trial. <i>Hypertension Research</i> , 2009 , 32, 248-54	4.7	25
23	Efficacy of adjuvant chemotherapy using tegafur-based regimen for curatively resected gastric cancer: update of a meta-analysis. <i>International Journal of Clinical Oncology</i> , 2009 , 14, 85-9	4.2	8
22	A phase I study of palliative chemoradiation therapy with paclitaxel and cisplatin for local symptoms due to an unresectable primary advanced or locally recurrent gastric adenocarcinoma. <i>Cancer Chemotherapy and Pharmacology</i> , 2009 , 64, 1071-7	3.5	8
21	Long-term effects of candesartan and amlodipine on cardiovascular mortality and morbidity in Japanese high-risk hypertensive patients: rationale, design, and characteristics of candesartan antihypertensive survival evaluation in Japan extension (CASE-J Ex). <i>Contemporary Clinical Trials</i> , 2009 , 30, 97-101	2.3	9
20	Effects of cardiac complications on cardiovascular events in Japanese high-risk hypertensive patients: subanalysis of the CASE-J trial. <i>Circulation Journal</i> , 2009 , 73, 1080-5	2.9	8
19	Clinical significance of left ventricular hypertrophy and changes in left ventricular mass in high-risk hypertensive patients: a subanalysis of the Candesartan Antihypertensive Survival Evaluation in Japan trial. <i>Journal of Hypertension</i> , 2009 , 27, 1705-12	1.9	22
18	Multicenter phase II study of FOLFOX for metastatic colorectal cancer (mCRC) in Japan; SWIFT-1 and 2 study. <i>Hepato-Gastroenterology</i> , 2009 , 56, 1346-53		6

17	A randomized phase II trial of preoperative exercise to reduce operative risk in gastric cancer patients with metabolic syndrome: adjuvant exercise for general elective surgery (AEGES) study group. <i>Japanese Journal of Clinical Oncology</i> , 2008 , 38, 71-3	2.8	9
16	The optimal target blood pressure for antihypertensive treatment in Japanese elderly patients with high-risk hypertension: a subanalysis of the Candesartan Antihypertensive Survival Evaluation in Japan (CASE-J) trial. <i>Hypertension Research</i> , 2008 , 31, 1595-601	4.7	34
15	Effects of candesartan compared with amlodipine in hypertensive patients with high cardiovascular risks: candesartan antihypertensive survival evaluation in Japan trial. <i>Hypertension</i> , 2008 , 51, 393-8	8.5	201
14	The domain structure of the EORTC QLQ-STO22 supported by Japanese validation data. <i>Psycho-Oncology</i> , 2008 , 17, 474-9	3.9	12
13	Meta-analyses of randomized trials assessing the interest chemotherapy in advanced gastric cancer. <i>Journal of Clinical Oncology</i> , 2008 , 26, 4563-4563	2.2	3
12	An individual patient data meta-analysis of adjuvant therapy with uracil-tegafur (UFT) in patients with curatively resected rectal cancer. <i>British Journal of Cancer</i> , 2007 , 96, 1170-7	8.7	52
11	Efficacy of adjuvant immunochemotherapy with polysaccharide K for patients with curative resections of gastric cancer. <i>Cancer Immunology, Immunotherapy</i> , 2007 , 56, 905-11	7.4	78
10	Pharmacokinetic study of weekly administration dose of paclitaxel in patients with advanced or recurrent gastric cancer in Japan. <i>Gastric Cancer</i> , 2007 , 10, 52-7	7.6	20
9	Multicenter phase II trial of combination chemotherapy with weekly paclitaxel and 5-fluorouracil for the treatment of advanced or recurrent gastric carcinoma. <i>Journal of Chemotherapy</i> , 2007 , 19, 444-50	2.3	10
8	Efficacy of adjuvant immunochemotherapy with polysaccharide K for patients with curatively resected colorectal cancer: a meta-analysis of centrally randomized controlled clinical trials. <i>Cancer Immunology, Immunotherapy</i> , 2006 , 55, 404-11	7.4	66
7	Efficacy of adjuvant chemotherapy using oral fluorinated pyrimidines for curatively resected gastric cancer: a meta-analysis of centrally randomized controlled clinical trials in Japan. <i>Journal of Chemotherapy</i> , 2006 , 18, 311-7	2.3	23
6	Factors associated with compliance and non-compliance by physicians in a large-scale randomized clinical trial. <i>Trials</i> , 2006 , 7, 26	2.8	2
5	Efficacy of oral anticancer agents for colorectal cancer. <i>Diseases of the Colon and Rectum</i> , 2006 , 49, S82-91	3.1	11
4	A feasibility study of sequential paclitaxel and S-1 (PTX/S-1) chemotherapy as postoperative adjuvant chemotherapy for advanced gastric cancer. <i>Gastric Cancer</i> , 2006 , 9, 114-9	7.6	12
3	Adjuvant chemotherapy with tegafur/uracil (UFT) for gastric cancer. A meta-analysis of centrally randomized clinical trials. <i>Journal of Clinical Oncology</i> , 2006 , 24, 4033-4033	2.2	4
2	Truth-telling to Patients with Advanced Cancer: A Comparison between Cancer center, University Hospitals, General City Hospitals, and Private General Practitioners in Japan. <i>Annals of Cancer Research and Therapy</i> , 2006 , 14, 1-11	0.2	2
1	Detection of early gastric cancer by a real-time autofluorescence imaging system. <i>Cancer Letters</i> , 2001 , 165, 155-9	9.9	60