

Ishan Agarwal

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2267235/publications.pdf>

Version: 2024-02-01

53
papers

770
citations

623734

14
h-index

677142

22
g-index

53
all docs

53
docs citations

53
times ranked

429
citing authors

#	ARTICLE	IF	CITATIONS
1	Insights into Himalayan biogeography from geckos: A molecular phylogeny of <i>Cyrtodactylus</i> (Squamata: Gekkonidae). <i>Molecular Phylogenetics and Evolution</i> , 2014, 80, 145-155.	2.7	73
2	A phylogeny of the only ground-dwelling radiation of <i>Cyrtodactylus</i> (Squamata, Gekkonidae): diversification of <i>Geckoella</i> across peninsular India and Sri Lanka. <i>Molecular Phylogenetics and Evolution</i> , 2015, 82, 193-199.	2.7	60
3	A phylogeny of open-habitat lizards (Squamata: Lacertidae: <i>Ophisops</i>) supports the antiquity of Indian grassy biomes. <i>Journal of Biogeography</i> , 2017, 44, 2021-2032.	3.0	36
4	The hills are alive with geckos! A radiation of a dozen species on sky islands across peninsular India (Squamata: Gekkonidae, <i>Hemiphyllodactylus</i>) with the description of three new species. <i>Organisms Diversity and Evolution</i> , 2019, 19, 341-361.	1.6	33
5	The role of wet-zone fragmentation in shaping biodiversity patterns in peninsular India: insights from the caecilian amphibian <i>Gegeneophis</i> . <i>Journal of Biogeography</i> , 2016, 43, 1091-1102.	3.0	30
6	Two new species of ground-dwelling <i>Cyrtodactylus</i> (<i>Geckoella</i>) from the Mysore Plateau, south India. <i>Zootaxa</i> , 2016, 4193, zootaxa.4193.2.2.	0.5	26
7	Phylogenetic partitioning of the third-largest vertebrate genus in the world, <i>Cyrtodactylus</i> Gray, 1827 (Reptilia; Squamata; Gekkonidae) and its relevance to taxonomy and conservation. <i>Vertebrate Zoology</i> , 0, 71, 101-154.	2.0	25
8	Cryptic species, taxonomic inflation, or a bit of both? New species phenomenon in Sri Lanka as suggested by a phylogeny of dwarf geckos (Reptilia, Squamata, Gekkonidae, <i>Cnemaspis</i>). <i>Systematics and Biodiversity</i> , 2017, 15, 427-439.	1.2	24
9	A review of the genus <i>Hemidactylus</i> Goldfuss, 1820 (Squamata: Gekkonidae) from Angola, with the description of two new species. <i>Zootaxa</i> , 2020, 4746, zootaxa.4746.1.1.	0.5	24
10	A new cryptic rock-dwelling <i>Hemidactylus</i> (Squamata: Gekkonidae) from south India. <i>Zootaxa</i> , 2011, 2765, 21.	0.5	24
11	A new species of the <i>Cyrtodactylus</i> (<i>Geckoella</i>) <i>collegalensis</i> (Beddome, 1870) complex (Squamata: Tj ETQq1 1 0.784314 rgBT /Over	0.5	22
12	An expanded ND2 phylogeny of the brookii and prashadi groups with the description of three new Indian <i>Hemidactylus</i> Oken (Squamata: Gekkonidae). <i>Zootaxa</i> , 2019, 4619, zootaxa.4619.3.2.	0.5	21
13	Granite boulders act as deep-time climate refugia: A Miocene divergent clade of rupicolous <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) from the Mysore Plateau, India, with descriptions of three new species. <i>Journal of Zoological Systematics and Evolutionary Research</i> , 2020, 58, 1234-1261.	1.4	21
14	The first teresomatian caecilian (Amphibia: Gymnophiona) from the Eastern Ghats of India—a new species of <i>Gegeneophis</i> ; Peters, 1880. <i>Zootaxa</i> , 2013, 3693, 534.	0.5	20
15	Cryptic species and Miocene diversification of Palaearctic naked-toed geckos (Squamata: Gekkonidae) in the Indian dry zone. <i>Zoologica Scripta</i> , 2014, 43, 455-471.	1.7	18
16	Two new <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) from the Shevaroy massif, Tamil Nadu, India, with a preliminary ND2 phylogeny of Indian <i>Cnemaspis</i> . <i>Zootaxa</i> , 2019, 4609, zootaxa.4609.1.3.	0.5	18
17	A new earless species of <i>Poyntonophrynus</i> (Anura, Bufonidae) from the Serra da Neve Inselberg, Namibe Province, Angola. <i>ZooKeys</i> , 2018, 780, 109-136.	1.1	17
18	How the African house gecko (<i>Hemidactylus mabouia</i>) conquered the world. <i>Royal Society Open Science</i> , 2021, 8, 210749.	2.4	17

#	ARTICLE	IF	CITATIONS
19	Six new <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from northeast India. <i>Zootaxa</i> , 2018, 4524, 501-535.	0.5	16
20	Phylogenetic partitioning of the third-largest vertebrate genus in the world, <i>Cyrtodactylus</i> Gray, 1827 (Reptilia; Squamata; Gekkonidae) and its relevance to taxonomy and conservation. <i>Vertebrate Zoology</i> , 0, 71, 101-154.	2.0	16
21	A new species of large-bodied, tuberculate <i>Hemidactylus</i> Oken (Squamata: Gekkonidae) from the Eastern Ghats, India. <i>Zootaxa</i> , 2017, 4347, 331.	0.5	13
22	The evolutionary history of an accidental model organism, the leopard gecko <i>Eublepharis macularius</i> (Squamata: Eublepharidae). <i>Molecular Phylogenetics and Evolution</i> , 2022, 168, 107414.	2.7	13
23	Two new species of bent toed geckos, <i>Cyrtodactylus</i> Gray, 1827 (Squamata: Gekkonidae) from Northeast India with comments on name-bearing types from the region. <i>Zootaxa</i> , 2018, 4420, 334-356.	0.5	12
24	A new rock dwelling <i>Hemidactylus</i> (Squamata: Gekkonidae) from Chhattisgarh, India. <i>Zootaxa</i> , 2015, 4021, 334-50.	0.5	11
25	Two more new species of <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) from the northern Western Ghats, Maharashtra, India. <i>Zootaxa</i> , 2019, 4656, zootaxa.4656.1.2.	0.5	11
26	A new species of South Asian <i>Cnemaspis</i> (Squamata: Gekkonidae) from the Eastern Ghats, India. <i>Zootaxa</i> , 2020, 4802, zootaxa.4802.3.3.	0.5	10
27	The phylogenetic position of the enigmatic Assam day gecko <i>Cnemaspis</i> cf. <i>assamensis</i> (Squamata: Gekkonidae) from India-Sri Lanka. <i>Amphibia - Reptilia</i> , 2021, 42, 355-367.	0.5	10
28	Revalidation of <i>Natrix</i> <i>clerkii</i> Wall, 1925, an overlooked species in the genus <i>Amphiesma</i> Duméril, Bibron & Duméril, 1854 (Squamata: Natricidae). <i>Zootaxa</i> , 2015, 3919, 375.	0.5	9
29	A new medium-bodied rupicolous <i>Hemidactylus</i> Goldfuss, 1820 (Squamata: Gekkonidae) from the Sirumalai massif, Tamil Nadu, India. <i>Zootaxa</i> , 2020, 4852, zootaxa.4852.1.4.	0.5	9
30	A new large-bodied, rupicolous <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) allied to <i>Cnemaspis heteropholis</i> Bauer, 2002 from the Central Western Ghats of Karnataka, India. <i>Zootaxa</i> , 2020, 4801, zootaxa.4801.1.2.	0.5	9
31	Another new <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from Guwahati, Assam, India. <i>Zootaxa</i> , 2020, 4732, zootaxa.4732.3.2.	0.5	9
32	The curious case of <i>Hemidactylus gujaratensis</i> (Squamata: Gekkonidae) from Gujarat, India. <i>Zootaxa</i> , 2018, 4388, 137-142.	0.5	8
33	Description of a new diminutive, rupicolous species of day-gecko (Squamata: Gekkonidae: <i>Cnemaspis</i>) from southern Sri Lanka. <i>Zootaxa</i> , 2019, 4565, 223.	0.5	8
34	A new cryptic <i>Cnemaspis</i> Strauch (Squamata: Gekkonidae) from an isolated granite hill on the Mysore Plateau, Karnataka, India. <i>Zootaxa</i> , 2020, 4845, zootaxa.4845.4.3.	0.5	8
35	Geckos in the granite: two new geckos (Squamata: Gekkonidae) from rocky, scrub habitats in Rishi Valley, Andhra Pradesh, India. <i>Zootaxa</i> , 2020, 4838, zootaxa.4838.4.1.	0.5	8
36	Two more new <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) from Tamil Nadu, India. <i>Zootaxa</i> , 2020, 4729, zootaxa.4729.2.6.	0.5	8

#	ARTICLE	IF	CITATIONS
37	Two new species of the <i>Ophisops microlepis</i> (Squamata: Lacertidae) complex from northwestern India with a key to Indian <i>Ophisops</i> . <i>Journal of Natural History</i> , 2018, 52, 819-847.	0.5	7
38	Two new species of yellow-tailed <i>Hemidactylus</i> Goldfuss, 1820 (Squamata: Gekkonidae) from rocky outcrops on the Telangana Plateau, India. <i>Zootaxa</i> , 2020, 4895, zootaxa.4895.4.2.	0.5	7
39	On the Status of <i>Cyrtodactylus malcolmsmithi</i> (Constable, 1949). <i>Breviora</i> , 2018, 557, 1-11.	0.5	6
40	A cryptic new species of rupicolous <i>Hemidactylus</i> Goldfuss, 1820 (Squamata: Gekkonidae) allied to <i>H. aaronbaueri</i> Giri, 2008 from the northern Western Ghats of Maharashtra, India. <i>Zootaxa</i> , 2021, 5020, 434-456.	0.5	6
41	Three new cryptic species of South Asian <i>Cnemaspis</i> Strauch, 1887 (Squamata, Gekkonidae) from Karnataka, India. <i>Vertebrate Zoology</i> , 0, 72, 115-142.	2.0	6
42	A new bent-toed gecko (Squamata: Gekkonidae: <i>Cyrtodactylus</i>) from the Western Himalayas, Himachal Pradesh, India. <i>Zootaxa</i> , 2018, 4446, 442.	0.5	5
43	A novel member of the <i>Hemidactylus brookii</i> complex (Squamata: Gekkonidae) from the Western Ghats of Maharashtra, India. <i>Zootaxa</i> , 2019, 4646, zootaxa.4646.2.2.	0.5	5
44	A new medium-sized rupicolous <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) of the <i>C. bangara</i> clade from granite boulder habitats in Krishnagiri, Tamil Nadu, India. <i>Zootaxa</i> , 2021, 4969, 351366.	0.5	5
45	Three more novel species of South Asian <i>Cnemaspis</i> Strauch, 1887 (Squamata, Gekkonidae) from Kalakad Mundanthurai Tiger Reserve, Tamil Nadu, India. <i>Vertebrate Zoology</i> , 0, 72, 385-422.	2.0	5
46	A novel, diminutive <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) from a sacred grove in Odisha, eastern India. <i>Zootaxa</i> , 2020, 4852, zootaxa.4852.4.6.	0.5	4
47	The identity and probable origin of the <i>Hemidactylus</i> geckos of the Maldives. <i>Herpetological Journal</i> , 2019, 29, 230-236.	0.6	4
48	A new small-bodied, polymorphic <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) from the Central Western Ghats of Karnataka, India. <i>Zootaxa</i> , 2021, 4950, 501-527.	0.5	3
49	A novel small-bodied rupicolous <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) from the Central Western Ghats of Karnataka, India. <i>Zootaxa</i> , 2021, 4969, 331-350.	0.5	3
50	A long-lost relic from the Eastern Ghats: Morphology, distribution and habitat of <i>Sepsophis punctatus</i> Beddome, 1870 (Squamata: Scincidae). <i>Zootaxa</i> , 2013, 3670, 55-62.	0.5	3
51	Lizards of the Thar Desert " Resource partitioning and community composition. <i>Journal of Arid Environments</i> , 2015, 118, 58-64.	2.4	2
52	Integrative Taxonomy in the Indian Subcontinent: Current Progress and Prospects. <i>Journal of the Indian Institute of Science</i> , 2021, 101, 125-149.	1.9	2
53	A new species of the genus <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) from Goa, India. <i>Zootaxa</i> , 2021, 5027, 254-268.	0.5	0