

Saemundur Gudmundsson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2269339/publications.pdf>

Version: 2024-02-01

28
papers

714
citations

623188

14
h-index

525886

27
g-index

28
all docs

28
docs citations

28
times ranked

676
citing authors

#	ARTICLE	IF	CITATIONS
1	Placental pulsatility index: a new, more sensitive parameter for predicting adverse outcome in pregnancies suspected of fetal growth restriction. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2017, 96, 216-222.	1.3	16
2	Alterations in ductus venosus velocity indices in relation to umbilical venous pulsations and perinatal outcome. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2016, 95, 645-651.	1.3	2
3	Increased pulsatility in the fetal ductus venosus is not related to altered cardiac strain in high-risk pregnancies. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2016, 29, 1328-1333.	0.7	5
4	Investigations on atrial function in fetuses with signs of impaired placental function. <i>Prenatal Diagnosis</i> , 2015, 35, 605-611.	1.1	9
5	Uterine artery Doppler in high-risk pregnancies at 23-24 gestational weeks is of value in predicting adverse outcome of pregnancy and selecting cases for more intense surveillance. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2014, 93, 1276-1281.	1.3	22
6	Fetal pulmonary and cerebral artery Doppler velocimetry in normal and high risk pregnancy. <i>Ginekologia Polska</i> , 2014, 85, 26-30.	0.3	3
7	Multivascular Doppler surveillance in high risk pregnancies. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2012, 25, 970-974.	0.7	5
8	Venous Doppler in the Evaluation of Fetal Hydrops. <i>Obstetrics and Gynecology International</i> , 2010, 2010, 1-7.	0.5	9
9	Signs of fetal brain sparing are not related to umbilical cord blood gases at birth. <i>Early Human Development</i> , 2009, 85, 467-470.	0.8	13
10	Placental Morphologic and Functional Imaging in High-Risk Pregnancies. <i>Seminars in Perinatology</i> , 2009, 33, 270-280.	1.1	25
11	Abnormal uterine artery Doppler in pregnancies suspected of a SGA fetus is related to increased risk of recurrence during next pregnancy. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2009, 88, 814-817.	1.3	2
12	Flow index evaluation of 3-d volume flow images: An in vivo and in vitro study. <i>Ultrasound in Medicine and Biology</i> , 2006, 32, 665-671.	0.7	31
13	Correlation of birth injury with maternal height and birthweight. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2005, 112, 764-767.	1.1	61
14	Preeclampsia - abnormal uterine artery Doppler is related to recurrence of symptoms during the next pregnancy. <i>Journal of Perinatal Medicine</i> , 2004, 32, 400-3.	0.6	10
15	New score indicating placental vascular resistance. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2003, 82, 807-812.	1.3	44
16	New score indicating placental vascular resistance. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2003, 82, 807-812.	1.3	3
17	Prospect for vaginal delivery of growth restricted fetuses with abnormal umbilical artery blood flow. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2003, 82, 828-833.	1.3	1
18	Are some perinatal deaths in immigrant groups linked to suboptimal perinatal care services?. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2002, 109, 677-682.	1.1	133

#	ARTICLE	IF	CITATIONS
19	Two types of umbilical venous pulsations and outcome of high-risk pregnancy. <i>Early Human Development</i> , 2001, 61, 111-117.	0.8	46
20	Fetal and placental power Doppler imaging in normal and high-risk pregnancy. <i>European Journal of Ultrasound: Official Journal of the European Federation of Societies for Ultrasound in Medicine and Biology</i> , 1999, 9, 223-230.	1.4	20
21	Uterine and umbilical artery velocimetry in pre-eclampsia. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1998, 77, 614-619.	1.3	27
22	Frequency of anal sphincter rupture at delivery in Sweden and Finland - result of difference in manual help to the baby's head. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1998, 77, 974-977.	1.3	42
23	Socioeconomic status and perinatal outcome according to residence area in the city of Malmö. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1997, 76, 318-323.	1.3	34
24	Can women with intrapartum rupture of anal sphincter still suffer after-effects two decades later?. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1997, 76, 601-603.	1.3	49
25	Fetal pulse oximetry a methodological study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1996, 75, 144-148.	1.3	15
26	Effects of hydralazine on placental and renal circulation in pre-eclampsia. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1995, 74, 415-418.	1.3	14
27	Doppler ultrasound recordings of placental blood velocity waveforms. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1991, 70, 389-390.	1.3	1
28	Blood Velocity Waveforms in the Fetal Aorta and Umbilical Artery as Predictors of Fetal Outcome: A Comparison. <i>American Journal of Perinatology</i> , 1991, 8, 1-6.	0.6	72