

Chris Wallace

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2277101/chris-wallace-publications-by-citations.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

134
papers

25,924
citations

48
h-index

156
g-index

156
ext. papers

29,908
ext. citations

13.2
avg, IF

7.45
L-index

#	Paper	IF	Citations
134	Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. <i>Nature</i> , 2007 , 447, 661-78	50.4	7801
133	Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes. <i>Science</i> , 2007 , 316, 1336-41	33.3	1823
132	Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. <i>Nature Genetics</i> , 2009 , 41, 25-34	36.3	1368
131	Robust associations of four new chromosome regions from genome-wide analyses of type 1 diabetes. <i>Nature Genetics</i> , 2007 , 39, 857-64	36.3	1159
130	Association scan of 14,500 nonsynonymous SNPs in four diseases identifies autoimmunity variants. <i>Nature Genetics</i> , 2007 , 39, 1329-37	36.3	1130
129	Common variants near MC4R are associated with fat mass, weight and risk of obesity. <i>Nature Genetics</i> , 2008 , 40, 768-75	36.3	1048
128	Genome-wide association study identifies eight loci associated with blood pressure. <i>Nature Genetics</i> , 2009 , 41, 666-76	36.3	970
127	Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. <i>PLoS Genetics</i> , 2014 , 10, e1004383	6	868
126	Genome-wide association analysis identifies 20 loci that influence adult height. <i>Nature Genetics</i> , 2008 , 40, 575-83	36.3	654
125	Genome-wide association study of CNVs in 16,000 cases of eight common diseases and 3,000 shared controls. <i>Nature</i> , 2010 , 464, 713-20	50.4	639
124	Lineage-Specific Genome Architecture Links Enhancers and Non-coding Disease Variants to Target Gene Promoters. <i>Cell</i> , 2016 , 167, 1369-1384.e19	56.2	556
123	Dense genotyping identifies and localizes multiple common and rare variant association signals in celiac disease. <i>Nature Genetics</i> , 2011 , 43, 1193-201	36.3	535
122	Meta-analysis of 28,141 individuals identifies common variants within five new loci that influence uric acid concentrations. <i>PLoS Genetics</i> , 2009 , 5, e1000504	6	495
121	Genome-wide association study identifies loci influencing concentrations of liver enzymes in plasma. <i>Nature Genetics</i> , 2011 , 43, 1131-8	36.3	415
120	Fine mapping of type 1 diabetes susceptibility loci and evidence for colocalization of causal variants with lymphoid gene enhancers. <i>Nature Genetics</i> , 2015 , 47, 381-6	36.3	414
119	Pervasive sharing of genetic effects in autoimmune disease. <i>PLoS Genetics</i> , 2011 , 7, e1002254	6	413
118	Genome-wide association scan meta-analysis identifies three Loci influencing adiposity and fat distribution. <i>PLoS Genetics</i> , 2009 , 5, e1000508	6	393

117	Genome-wide association study identifies genes for biomarkers of cardiovascular disease: serum urate and dyslipidemia. <i>American Journal of Human Genetics</i> , 2008 , 82, 139-49	11	361
116	Widespread seasonal gene expression reveals annual differences in human immunity and physiology. <i>Nature Communications</i> , 2015 , 6, 7000	17.4	268
115	SLC2A9 is a high-capacity urate transporter in humans. <i>PLoS Medicine</i> , 2008 , 5, e197	11.6	254
114	The imprinted DLK1-MEG3 gene region on chromosome 14q32.2 alters susceptibility to type 1 diabetes. <i>Nature Genetics</i> , 2010 , 42, 68-71	36.3	238
113	The International Human Epigenome Consortium: A Blueprint for Scientific Collaboration and Discovery. <i>Cell</i> , 2016 , 167, 1145-1149	56.2	232
112	A trans-acting locus regulates an anti-viral expression network and type 1 diabetes risk. <i>Nature</i> , 2010 , 467, 460-4	50.4	224
111	Inherited variation in vitamin D genes is associated with predisposition to autoimmune disease type 1 diabetes. <i>Diabetes</i> , 2011 , 60, 1624-31	0.9	222
110	Genetic loci influencing kidney function and chronic kidney disease. <i>Nature Genetics</i> , 2010 , 42, 373-5	36.3	205
109	Genome-wide association analysis of autoantibody positivity in type 1 diabetes cases. <i>PLoS Genetics</i> , 2011 , 7, e1002216	6	195
108	A type I interferon transcriptional signature precedes autoimmunity in children genetically at risk for type 1 diabetes. <i>Diabetes</i> , 2014 , 63, 2538-50	0.9	188
107	Blood pressure loci identified with a gene-centric array. <i>American Journal of Human Genetics</i> , 2011 , 89, 688-700	11	137
106	Capture Hi-C reveals novel candidate genes and complex long-range interactions with related autoimmune risk loci. <i>Nature Communications</i> , 2015 , 6, 10069	17.4	121
105	Seven newly identified loci for autoimmune thyroid disease. <i>Human Molecular Genetics</i> , 2012 , 21, 5202-85.6		120
104	Dense genotyping of immune-related susceptibility loci reveals new insights into the genetics of psoriatic arthritis. <i>Nature Communications</i> , 2015 , 6, 6046	17.4	103
103	Meta-analysis of Dense Genecentric Association Studies Reveals Common and Uncommon Variants Associated with Height. <i>American Journal of Human Genetics</i> , 2011 , 88, 6-18	11	103
102	Statistical colocalization of genetic risk variants for related autoimmune diseases in the context of common controls. <i>Nature Genetics</i> , 2015 , 47, 839-46	36.3	97
101	Integration of disease association and eQTL data using a Bayesian colocalisation approach highlights six candidate causal genes in immune-mediated diseases. <i>Human Molecular Genetics</i> , 2015 , 24, 3305-13	5.6	94
100	Long-range DNA looping and gene expression analyses identify DEXI as an autoimmune disease candidate gene. <i>Human Molecular Genetics</i> , 2012 , 21, 322-33	5.6	91

99	IL-21 production by CD4+ effector T cells and frequency of circulating follicular helper T cells are increased in type 1 diabetes patients. <i>Diabetologia</i> , 2015 , 58, 781-90	10.3	86
98	Haplotypes of the WNK1 gene associate with blood pressure variation in a severely hypertensive population from the British Genetics of Hypertension study. <i>Human Molecular Genetics</i> , 2005 , 14, 1805-14	5.6	82
97	Meta-analysis of Dense Genecentric Association Studies Reveals Common and Uncommon Variants Associated with Height. <i>American Journal of Human Genetics</i> , 2012 , 90, 1116-1117	11	78
96	Regulatory T Cell Responses in Participants with Type 1 Diabetes after a Single Dose of Interleukin-2: A Non-Randomised, Open Label, Adaptive Dose-Finding Trial. <i>PLoS Medicine</i> , 2016 , 13, e1002139	11.6	76
95	Comparative analysis of genome-wide association studies signals for lipids, diabetes, and coronary heart disease: Cardiovascular Biomarker Genetics Collaboration. <i>European Heart Journal</i> , 2012 , 33, 393-407	9.5	75
94	Genome-wide association study of eosinophilic granulomatosis with polyangiitis reveals genomic loci stratified by ANCA status. <i>Nature Communications</i> , 2019 , 10, 5120	17.4	71
93	Statistical colocalization of monocyte gene expression and genetic risk variants for type 1 diabetes. <i>Human Molecular Genetics</i> , 2012 , 21, 2815-24	5.6	71
92	Reduced expression of IFIH1 is protective for type 1 diabetes. <i>PLoS ONE</i> , 2010 , 5, e12646	3.7	66
91	Eliciting priors and relaxing the single causal variant assumption in colocalisation analyses. <i>PLoS Genetics</i> , 2020 , 16, e1008720	6	54
90	Trends in drug overdose deaths in England and Wales 1993-98: methadone does not kill more people than heroin. <i>Addiction</i> , 2003 , 98, 419-25	4.6	54
89	Statistical testing of shared genetic control for potentially related traits. <i>Genetic Epidemiology</i> , 2013 , 37, 802-13	2.6	53
88	Effects of long-term averaging of quantitative blood pressure traits on the detection of genetic associations. <i>American Journal of Human Genetics</i> , 2014 , 95, 49-65	11	52
87	Common genetic variation near the phospholamban gene is associated with cardiac repolarisation: meta-analysis of three genome-wide association studies. <i>PLoS ONE</i> , 2009 , 4, e6138	3.7	50
86	Extreme clonality in lymphoblastoid cell lines with implications for allele specific expression analyses. <i>PLoS ONE</i> , 2008 , 3, e2966	3.7	48
85	Two-dimensional genome-scan identifies novel epistatic loci for essential hypertension. <i>Human Molecular Genetics</i> , 2006 , 15, 1365-74	5.6	47
84	Cells with Treg-specific FOXP3 demethylation but low CD25 are prevalent in autoimmunity. <i>Journal of Autoimmunity</i> , 2017 , 84, 75-86	15.5	46
83	A method for gene-based pathway analysis using genomewide association study summary statistics reveals nine new type 1 diabetes associations. <i>Genetic Epidemiology</i> , 2014 , 38, 661-70	2.6	43
82	Dissection of a Complex Disease Susceptibility Region Using a Bayesian Stochastic Search Approach to Fine Mapping. <i>PLoS Genetics</i> , 2015 , 11, e1005272	6	42

81	Chromosome contacts in activated T cells identify autoimmune disease candidate genes. <i>Genome Biology</i> , 2017 , 18, 165	18.3	41
80	Multi-parametric flow cytometric and genetic investigation of the peripheral B cell compartment in human type 1 diabetes. <i>Clinical and Experimental Immunology</i> , 2014 , 177, 571-85	6.2	41
79	An allele of IKZF1 (Ikaros) conferring susceptibility to childhood acute lymphoblastic leukemia protects against type 1 diabetes. <i>Diabetes</i> , 2011 , 60, 1041-4	0.9	41
78	Improved power offered by a score test for linkage disequilibrium mapping of quantitative-trait loci by selective genotyping. <i>American Journal of Human Genetics</i> , 2006 , 78, 498-504	11	38
77	A pleiotropy-informed Bayesian false discovery rate adapted to a shared control design finds new disease associations from GWAS summary statistics. <i>PLoS Genetics</i> , 2015 , 11, e1004926	6	37
76	Polymorphisms in the WNK1 gene are associated with blood pressure variation and urinary potassium excretion. <i>PLoS ONE</i> , 2009 , 4, e5003	3.7	36
75	Genetic association analyses of atopic illness and proinflammatory cytokine genes with type 1 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2011 , 27, 838-43	7.5	35
74	Natural Variation in Interleukin-2 Sensitivity Influences Regulatory T-Cell Frequency and Function in Individuals With Long-standing Type 1 Diabetes. <i>Diabetes</i> , 2015 , 64, 3891-902	0.9	34
73	Promoter interactome of human embryonic stem cell-derived cardiomyocytes connects GWAS regions to cardiac gene networks. <i>Nature Communications</i> , 2018 , 9, 2526	17.4	34
72	Postthymic expansion in human CD4 naive T cells defined by expression of functional high-affinity IL-2 receptors. <i>Journal of Immunology</i> , 2013 , 190, 2554-66	5.3	34
71	Chromosome 2p shows significant linkage to antihypertensive response in the British Genetics of Hypertension Study. <i>Hypertension</i> , 2006 , 47, 603-8	8.5	31
70	Glutathione S-transferase variants and hypertension. <i>Journal of Hypertension</i> , 2008 , 26, 1343-52	1.9	30
69	Neonatal and adult recent thymic emigrants produce IL-8 and express complement receptors CR1 and CR2. <i>JCI Insight</i> , 2017 , 2,	9.9	26
68	Epigenetic analysis of regulatory T cells using multiplex bisulfite sequencing. <i>European Journal of Immunology</i> , 2015 , 45, 3200-3	6.1	21
67	Plasma concentrations of soluble IL-2 receptor (CD25) are increased in type 1 diabetes and associated with reduced C-peptide levels in young patients. <i>Diabetologia</i> , 2014 , 57, 366-72	10.3	21
66	Increased support for linkage of a novel locus on chromosome 5q13 for essential hypertension in the British Genetics of Hypertension Study. <i>Hypertension</i> , 2006 , 48, 105-11	8.5	21
65	Linkage analysis using co-phenotypes in the BRIGHT study reveals novel potential susceptibility loci for hypertension. <i>American Journal of Human Genetics</i> , 2006 , 79, 323-31	11	20
64	The chromosome 6q22.33 region is associated with age at diagnosis of type 1 diabetes and disease risk in those diagnosed under 5 years of age. <i>Diabetologia</i> , 2018 , 61, 147-157	10.3	19

63	Stochastic search and joint fine-mapping increases accuracy and identifies previously unreported associations in immune-mediated diseases. <i>Nature Communications</i> , 2019 , 10, 3216	17.4	19
62	A method for identifying genetic heterogeneity within phenotypically defined disease subgroups. <i>Nature Genetics</i> , 2017 , 49, 310-316	36.3	17
61	Improving the coverage of credible sets in Bayesian genetic fine-mapping. <i>PLoS Computational Biology</i> , 2020 , 16, e1007829	5	16
60	Multivariate Genome-wide Association Analysis of a Cytokine Network Reveals Variants with Widespread Immune, Haematological, and Cardiometabolic Pleiotropy. <i>American Journal of Human Genetics</i> , 2019 , 105, 1076-1090	11	16
59	A genome-wide assessment of the role of untagged copy number variants in type 1 diabetes. <i>PLoS Genetics</i> , 2014 , 10, e1004367	6	16
58	Resolving mechanisms of immune-mediated disease in primary CD4 T cells. <i>EMBO Molecular Medicine</i> , 2020 , 12, e12112	12	14
57	Seropositivity in blood donors and pregnant women during the first year of SARS-CoV-2 transmission in Stockholm, Sweden. <i>Journal of Internal Medicine</i> , 2021 , 290, 666-676	10.8	14
56	Genetic association analysis of inositol polyphosphate phosphatase-like 1 (INPPL1, SHIP2) variants with essential hypertension. <i>Journal of Medical Genetics</i> , 2007 , 44, 603-5	5.8	13
55	Haplotypes of the beta-2 adrenergic receptor associate with high diastolic blood pressure in the Caerphilly prospective study. <i>Journal of Hypertension</i> , 2006 , 24, 471-7	1.9	13
54	Fine-mapping genetic associations. <i>Human Molecular Genetics</i> , 2020 , 29, R81-R88	5.6	13
53	A hybrid qPCR/SNP array approach allows cost efficient assessment of KIR gene copy numbers in large samples. <i>BMC Genomics</i> , 2014 , 15, 274	4.5	12
52	VSEAMS: a pipeline for variant set enrichment analysis using summary GWAS data identifies IKZF3, BATF and ESRRA as key transcription factors in type 1 diabetes. <i>Bioinformatics</i> , 2014 , 30, 3342-8	7.2	12
51	Functional effects of variation in transcription factor binding highlight long-range gene regulation by epromoters. <i>Nucleic Acids Research</i> , 2020 , 48, 2866-2879	20.1	11
50	Fine mapping chromatin contacts in capture Hi-C data. <i>BMC Genomics</i> , 2019 , 20, 77	4.5	10
49	Mendelian randomization. <i>Nature Reviews Methods Primers</i> , 2022 , 2,		10
48	simGWAS: a fast method for simulation of large scale case-control GWAS summary statistics. <i>Bioinformatics</i> , 2019 , 35, 1901-1906	7.2	9
47	Linkage analysis of susceptibility to leprosy type using an IBD regression method. <i>Genes and Immunity</i> , 2004 , 5, 221-5	4.4	8
46	Type 1 diabetes genome-wide association analysis with imputation identifies five new risk regions		8

45	Identification of susceptibility loci for Takayasu arteritis through a large multi-ancestral genome-wide association study. <i>American Journal of Human Genetics</i> , 2021 , 108, 84-99	11	8
44	Information capture using SNPs from HapMap and whole-genome chips differs in a sample of inflammatory and cardiovascular gene-centric regions from genome-wide estimates. <i>Genome Research</i> , 2007 , 17, 1596-602	9.7	7
43	Estimating the relative recurrence risk ratio using a global cross-ratio model. <i>Genetic Epidemiology</i> , 2003 , 25, 293-302	2.6	7
42	Seropositivity in blood donors and pregnant women during the first year of SARS-CoV-2 transmission in Stockholm, Sweden		5
41	Patient-reported wellbeing and clinical disease measures over time captured by multivariate trajectories of disease activity in individuals with juvenile idiopathic arthritis in the UK: a multicentre prospective longitudinal study. <i>Lancet Rheumatology, The</i> , 2021 , 3, e111-e121	14.2	5
40	A more accurate method for colocalisation analysis allowing for multiple causal variants. <i>PLoS Genetics</i> , 2021 , 17, e1009440	6	5
39	Extra-binomial variation approach for analysis of pooled DNA sequencing data. <i>Bioinformatics</i> , 2012 , 28, 2898-904	7.2	4
38	Accurate error control in high dimensional association testing using conditional false discovery rates		4
37	Estimating the relative recurrence risk ratio for leprosy in Karonga District, Malawi. <i>Leprosy Review</i> , 2003 , 74, 133-40	0.6	4
36	Appropriate use of information on family history of disease in recruitment for linkage analysis studies. <i>Annals of Human Genetics</i> , 2006 , 70, 360-71	2.2	3
35	Statistical Colocalization of Genetic Risk Variants for Related Autoimmune Diseases in the Context of Common Controls		3
34	Probabilistic approaches for classifying highly variable anti-SARS-CoV-2 antibody responses		3
33	Genetically distinct clinical subsets, and associations with asthma and eosinophil abundance, within Eosinophilic Granulomatosis with Polyangiitis		3
32	A Pleiotropy-Informed Bayesian False Discovery Rate adapted to a Shared Control Design Finds New Disease Associations From GWAS Summary Statistics		3
31	Genetic feature engineering enables characterisation of shared risk factors in immune-mediated diseases. <i>Genome Medicine</i> , 2020 , 12, 106	14.4	3
30	Accurate error control in high-dimensional association testing using conditional false discovery rates. <i>Biometrical Journal</i> , 2021 , 63, 1096-1130	1.5	3
29	Eliciting priors and relaxing the single causal variant assumption in colocalisation analyses		2
28	A rare IL2RA haplotype identifies SNP rs61839660 as causal for autoimmunity		2

27	Detecting chromosomal interactions in Capture Hi-C data with CHiCAGO and companion tools. <i>Nature Protocols</i> , 2021 , 16, 4144-4176	18.8	2
26	Multi-tissue transcriptome-wide association studies. <i>Genetic Epidemiology</i> , 2021 , 45, 324-337	2.6	2
25	Fine mapping chromatin contacts in capture Hi-C data		1
24	A method for identifying genetic heterogeneity within phenotypically-defined disease subgroups		1
23	Cells with Treg-specific FOXP3 demethylation but low CD25 are prevalent in autoimmunity		1
22	Informed dimension reduction of clinically-related genome-wide association summary data characterises cross-trait axes of genetic risk		1
21	EPISPOT: an epigenome-driven approach for detecting and interpreting hotspots in molecular QTL studies		1
20	simGWAS: a fast method for simulation of large scale case-control GWAS summary statistics		1
19	Improving the coverage of credible sets in Bayesian genetic fine-mapping		1
18	Recent thymic emigrants produce antimicrobial IL-8, express complement receptors and are precursors of a tissue-homing Th8 lineage of memory cells		1
17	Comparison of sparse biclustering algorithms for gene expression datasets. <i>Briefings in Bioinformatics</i> , 2021 , 22,	13.4	1
16	EPISPOT: An epigenome-driven approach for detecting and interpreting hotspots in molecular QTL studies. <i>American Journal of Human Genetics</i> , 2021 , 108, 983-1000	11	1
15	Detection of quantitative trait loci from RNA-seq data with or without genotypes using BaseQTL.. <i>Nature Computational Science</i> , 2021 , 1, 421-432		1
14	A more accurate method for colocalisation analysis allowing for multiple causal variants		1
13	Probabilistic classification of anti-SARS-CoV-2 antibody responses improves seroprevalence estimates.. <i>Clinical and Translational Immunology</i> , 2022 , 11, e1379	6.8	1
12	INNODIA Master Protocol for the evaluation of investigational medicinal products in children, adolescents and adults with newly diagnosed type 1 diabetes.. <i>Trials</i> , 2022 , 23, 414	2.8	1
11	The flashfm approach for fine-mapping multiple quantitative traits. <i>Nature Communications</i> , 2021 , 12, 6147	17.4	0
10	Leveraging auxiliary data from arbitrary distributions to boost GWAS discovery with Flexible cFDR. <i>PLoS Genetics</i> , 2021 , 17, e1009853	6	0

- 9 O1-11-04: Temporal Clustering Reveals Heterogeneity Of Cognitive Decline In Dementia **2016**, 12, P203-P203
- 8 Improving the coverage of credible sets in Bayesian genetic fine-mapping **2020**, 16, e1007829
- 7 Improving the coverage of credible sets in Bayesian genetic fine-mapping **2020**, 16, e1007829
- 6 Improving the coverage of credible sets in Bayesian genetic fine-mapping **2020**, 16, e1007829
- 5 Improving the coverage of credible sets in Bayesian genetic fine-mapping **2020**, 16, e1007829
- 4 Eliciting priors and relaxing the single causal variant assumption in colocalisation analyses **2020**, 16, e1008720
- 3 Eliciting priors and relaxing the single causal variant assumption in colocalisation analyses **2020**, 16, e1008720
- 2 Eliciting priors and relaxing the single causal variant assumption in colocalisation analyses **2020**, 16, e1008720
- 1 Eliciting priors and relaxing the single causal variant assumption in colocalisation analyses **2020**, 16, e1008720