

Hiroshi Itagaki

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2277554/publications.pdf>

Version: 2024-02-01

34
papers

1,131
citations

687363

13
h-index

414414

32
g-index

34
all docs

34
docs citations

34
times ranked

928
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Co-Culture of THP-1 Cells and Normal Human Epidermal Keratinocytes (NHEK) for Modified Human Cell Line Activation Test (h-CLAT). <i>Applied Sciences (Switzerland)</i> , 2022, 12, 6207. | 2.5 | 1 |
| 2 | An acid-hydrolyzed wheat protein activates the inflammatory and NF- κ B pathways leading to long TSLP transcription in human keratinocytes. <i>Journal of Toxicological Sciences</i> , 2020, 45, 327-337. | 1.5 | 1 |
| 3 | Predicting the results of a 24-hr human patch test for surfactants: utility of margin-setting in a reconstructed human epidermis model. <i>Journal of Toxicological Sciences</i> , 2019, 44, 393-403. | 1.5 | 3 |
| 4 | Some non-sensitizers upregulate CD54 expression by activation of the NLRP3 inflammasome in THP-1 cells. <i>Journal of Toxicological Sciences</i> , 2019, 44, 213-224. | 1.5 | 6 |
| 5 | Eliminating the contribution of lipopolysaccharide to protein allergenicity in the human cell-line activation test (h-CLAT). <i>Journal of Toxicological Sciences</i> , 2019, 44, 283-297. | 1.5 | 3 |
| 6 | Lipopolysaccharide interferes with the use of the human Cell Line Activation Test to determine the allergic potential of proteins. <i>Journal of Pharmacological and Toxicological Methods</i> , 2018, 92, 34-42. | 0.7 | 4 |
| 7 | Predictive performance and inter-laboratory reproducibility in assessing eye irritation potential of water- and oil-soluble mixtures using the Short Time Exposure test method. <i>Toxicology in Vitro</i> , 2018, 48, 78-85. | 2.4 | 4 |
| 8 | Expansion of the applicability domain for highly volatile substances on the Short Time Exposure test method and the predictive performance in assessing eye irritation potential. <i>Journal of Toxicological Sciences</i> , 2018, 43, 407-422. | 1.5 | 6 |
| 9 | Improvement of human cell line activation test (h-CLAT) using short-time exposure methods for prevention of false-negative results. <i>Journal of Toxicological Sciences</i> , 2018, 43, 229-240. | 1.5 | 10 |
| 10 | Assessment of the skin sensitizing potential of chemicals, contained in foods and/or cosmetic ingredients, using a modified local lymph node assay with an elicitation phase (LLNA:DAE) method. <i>Journal of Toxicological Sciences</i> , 2018, 43, 513-520. | 1.5 | 7 |
| 11 | Acidic conditions induce the suppression of CD86 and CD54 expression in THP-1 cells. <i>Journal of Toxicological Sciences</i> , 2018, 43, 299-309. | 1.5 | 8 |
| 12 | Preventing false-negatives in the in vitro skin sensitization testing of acid anhydrides using interleukin-8 release assays. <i>Toxicology in Vitro</i> , 2017, 42, 69-75. | 2.4 | 5 |
| 13 | Long form of thymic stromal lymphopoietin of keratinocytes is induced by protein allergens. <i>Journal of Immunotoxicology</i> , 2017, 14, 178-187. | 1.7 | 10 |
| 14 | Quantitative analysis of the relationship between the LLNA:DAE method results and the LLNA EC3 values highlights the connection between the elicitation and induction phases during skin sensitization. <i>Fundamental Toxicological Sciences</i> , 2016, 3, 27-31. | 0.6 | 1 |
| 15 | Tributyltin induces mitochondrial fission through Mfn1 degradation in human induced pluripotent stem cells. <i>Toxicology in Vitro</i> , 2016, 34, 257-263. | 2.4 | 24 |
| 16 | Unsaturated fatty acids show clear elicitation responses in a modified local lymph node assay with an elicitation phase, and test positive in the direct peptide reactivity assay. <i>Journal of Toxicological Sciences</i> , 2015, 40, 843-853. | 1.5 | 6 |
| 17 | Further development of LLNA:DAE method as stand-alone skin-sensitization testing method and applied for evaluation of relative skin-sensitizing potency between chemicals. <i>Journal of Toxicological Sciences</i> , 2015, 40, 137-150. | 1.5 | 8 |
| 18 | Development of LLNA:DAE: a new local lymph node assay that includes the elicitation phase, discriminates borderline-positive chemicals, and is useful for cross-sensitization testing. <i>Journal of Toxicological Sciences</i> , 2014, 39, 147-161. | 1.5 | 8 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Changes of cell-surface thiols and intracellular signaling in human monocytic cell line THP-1 treated with diphenylcyclopropenone. <i>Journal of Toxicological Sciences</i> , 2010, 35, 871-879. | 1.5 | 13 |
| 20 | A Comparative Evaluation of <i>In Vitro</i> Skin Sensitisation Tests: The Human Cell-line Activation Test (h-CLAT) versus the Local Lymph Node Assay (LLNA). <i>ATLA Alternatives To Laboratory Animals</i> , 2010, 38, 275-284. | 1.0 | 107 |
| 21 | Oxidation of Cell Surface Thiol Groups by Contact Sensitizers Triggers the Maturation of Dendritic Cells. <i>Journal of Investigative Dermatology</i> , 2010, 130, 175-183. | 0.7 | 31 |
| 22 | The relationship between CD86/CD54 expression and THP-1 cell viability in an in vitro skin sensitization test – human cell line activation test (h-CLAT). <i>Cell Biology and Toxicology</i> , 2009, 25, 109-126. | 5.3 | 92 |
| 23 | Evaluation of changes of cell-surface thiols as a new biomarker for in vitro sensitization test. <i>Toxicology in Vitro</i> , 2009, 23, 687-696. | 2.4 | 39 |
| 24 | Development of an in vitro photosensitization assay using human monocyte-derived cells. <i>Toxicology in Vitro</i> , 2009, 23, 911-918. | 2.4 | 23 |
| 25 | Modification of cell-surface thiols elicits activation of human monocytic cell line THP-1: Possible involvement in effect of haptens 2,4-dinitrochlorobenzene and nickel sulfate. <i>Journal of Toxicological Sciences</i> , 2009, 34, 139-150. | 1.5 | 24 |
| 26 | Development of an in vitro skin sensitization test using human cell lines: The human Cell Line Activation Test (h-CLAT). <i>Toxicology in Vitro</i> , 2006, 20, 767-773. | 2.4 | 266 |
| 27 | Development of an in vitro skin sensitization test using human cell lines; human Cell Line Activation Test (h-CLAT) II. An inter-laboratory study of the h-CLAT. <i>Toxicology in Vitro</i> , 2006, 20, 774-784. | 2.4 | 197 |
| 28 | QUANTITATIVE MEASUREMENT OF SPLICED XBP1 mRNA AS AN INDICATOR OF ENDOPLASMIC RETICULUM STRESS. <i>Journal of Toxicological Sciences</i> , 2006, 31, 149-156. | 1.5 | 101 |
| 29 | SIRC-CVS CYTOTOXICITY TEST: AN ALTERNATIVE FOR PREDICTING RODENT ACUTE SYSTEMIC TOXICITY. <i>Journal of Toxicological Sciences</i> , 2006, 31, 371-379. | 1.5 | 10 |
| 30 | Evaluation of CD86 expression and MHC class II molecule internalization in THP-1 human monocyte cells as predictive endpoints for contact sensitizers. <i>Toxicology in Vitro</i> , 2002, 16, 711-716. | 2.4 | 86 |
| 31 | Utility of MTT assay in three-dimensional cultured human skin model as an alternative for draize skin irritation test: approach using diffusion law of irritant in skin and toxicokinetics-toxicodynamics correlation. <i>Pharmaceutical Research</i> , 2002, 19, 669-675. | 3.5 | 24 |
| 32 | Selection of the First Inspection Time Based on Maximization of Amount of Information. <i>Journal of the Society of Naval Architects of Japan</i> , 1994, 1994, 597-602. | 0.2 | 0 |
| 33 | Reliability Assessment by Simulation of Fatigue Crack Growth. <i>Journal of the Society of Naval Architects of Japan</i> , 1989, 1989, 253-264. | 0.2 | 3 |
| 34 | Fatigue Crack Propagation under Controlled Stress Intensity Factor- (I). <i>Journal of the Society of Naval Architects of Japan</i> , 1973, 1973, 221-234. | 0.2 | 0 |