

Ryo Toya

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2281631/publications.pdf>

Version: 2024-02-01

56
papers

967
citations

623734

14
h-index

477307

29
g-index

63
all docs

63
docs citations

63
times ranked

1537
citing authors

#	ARTICLE	IF	CITATIONS
1	Grading Astrocytic Tumors by Using Apparent Diffusion Coefficient Parameters: Superiority of a One-versus Two-Parameter Pilot Method. <i>Radiology</i> , 2009, 251, 838-845.	7.3	170
2	Conformal radiation therapy for portal vein tumor thrombosis of hepatocellular carcinoma. <i>Radiotherapy and Oncology</i> , 2007, 84, 266-271.	0.6	104
3	IL-6 controls resistance to radiation by suppressing oxidative stress via the Nrf2-antioxidant pathway in oral squamous cell carcinoma. <i>British Journal of Cancer</i> , 2016, 115, 1234-1244.	6.4	87
4	Detection of Hemorrhagic Hypointense Foci in the Brain on Susceptibility-Weighted Imaging. <i>Academic Radiology</i> , 2007, 14, 1011-1019.	2.5	78
5	Tumor motion changes in stereotactic body radiotherapy for liver tumors: an evaluation based on four-dimensional cone-beam computed tomography and fiducial markers. <i>Radiation Oncology</i> , 2017, 12, 61.	2.7	47
6	Radiation-induced Parotid Gland Changes in Oral Cancer Patients: Correlation Between Parotid Volume and Saliva Production. <i>Japanese Journal of Clinical Oncology</i> , 2010, 40, 42-46.	1.3	45
7	Spleen Dose-Volume Parameters as a Predictor of Treatment-related Lymphopenia During Definitive Chemoradiotherapy for Esophageal Cancer. <i>In Vivo</i> , 2018, 32, 1519-1525.	1.3	29
8	Tumor budding as a novel predictor of occult metastasis in cT2N0 tongue squamous cell carcinoma. <i>Human Pathology</i> , 2018, 76, 1-8.	2.0	27
9	Circulating miRNA-1290 as a potential biomarker for response to chemoradiotherapy and prognosis of patients with advanced oral squamous cell carcinoma: A single-center retrospective study. <i>Tumor Biology</i> , 2019, 41, 101042831982685.	1.8	26
10	Dosimetric predictors of treatment-related lymphopenia induced by palliative radiotherapy: predictive ability of dose-volume parameters based on body surface contour. <i>Radiology and Oncology</i> , 2017, 51, 228-234.	1.7	20
11	Plan quality and delivery time comparisons between volumetric modulated arc therapy and intensity modulated radiation therapy for scalp angiosarcoma: A planning study. <i>Journal of Medical Radiation Sciences</i> , 2018, 65, 39-47.	1.5	19
12	Extracellular vesicles derived from radioresistant oral squamous cell carcinoma cells contribute to the acquisition of radioresistance via the miR-503a-3p-BAK axis. <i>Journal of Extracellular Vesicles</i> , 2021, 10, e12169.	12.2	18
13	The antioxidative stress regulator Nrf2 potentiates radioresistance of oral squamous cell carcinoma accompanied with metabolic modulation. <i>Laboratory Investigation</i> , 2022, 102, 896-907.	3.7	18
14	Hypofractionated palliative volumetric modulated arc radiotherapy with the Radiation Oncology Study Group 8502 "QUAD shot" regimen for incurable head and neck cancer. <i>Radiation Oncology</i> , 2020, 15, 123.	2.7	17
15	Impact of hybrid FDG-PET/CT on gross tumor volume definition of cervical esophageal cancer: reducing interobserver variation. <i>Journal of Radiation Research</i> , 2019, 60, 348-352.	1.6	15
16	Diagnostic Value of FDG-PET/CT for the Identification of Extranodal Extension in Patients With Head and Neck Squamous Cell Carcinoma. <i>Anticancer Research</i> , 2020, 40, 2073-2077.	1.1	15
17	Onodera's prognostic nutritional index correlates with tumor immune environment and survival in patients with oral squamous cell carcinoma undergoing chemoradiotherapy. <i>Translational Oncology</i> , 2020, 13, 100850.	3.7	14
18	FDG-PET/CT-based Gross Tumor Volume Contouring for Radiation Therapy Planning: An Experimental Phantom Study. <i>Journal of Radiation Research</i> , 2012, 53, 338-341.	1.6	13

#	ARTICLE	IF	CITATIONS
19	Predictors of Pain Palliation After Radiation Therapy for Painful Tumors: A Prospective Observational Study. <i>International Journal of Radiation Oncology Biology Physics</i> , 2018, 101, 1061-1068.	0.8	12
20	Pain Response Rates After Conventional Radiation Therapy for Bone Metastases in Prospective Nonrandomized Studies: A Systematic Review. <i>Practical Radiation Oncology</i> , 2019, 9, 81-88.	2.1	11
21	FBXW7 expression affects the response to chemoradiotherapy and overall survival among patients with oral squamous cell carcinoma: A single-center retrospective study. <i>Tumor Biology</i> , 2017, 39, 101042831773177.	1.8	10
22	Image quality of four-dimensional cone-beam computed tomography obtained at various gantry rotation speeds for liver stereotactic body radiation therapy with fiducial markers. <i>Physica Medica</i> , 2018, 45, 19-24.	0.7	10
23	Impact of 99mTc-GSA SPECT Image-Guided Inverse Planning on Dose-Function Histogram Parameters for Stereotactic Body Radiation Therapy Planning for Patients With Hepatocellular Carcinoma: A Dosimetric Comparison Study. <i>Dose-Response</i> , 2019, 17, 155932581983214.	1.6	10
24	Radiation-induced Liver Injury after 3D-conformal Radiotherapy for Hepatocellular Carcinoma: Quantitative Assessment Using Gd-EOB-DTPA-enhanced MRI. <i>Acta Medica Okayama</i> , 2017, 71, 25-29.	0.2	10
25	Radiation therapy for nasopharyngeal carcinoma: the predictive value of interim survival assessment. <i>Journal of Radiation Research</i> , 2016, 57, 541-547.	1.6	9
26	Respiratory Gating during Stereotactic Body Radiotherapy for Lung Cancer Reduces Tumor Position Variability. <i>PLoS ONE</i> , 2014, 9, e112824.	2.5	8
27	Image quality evaluation of in-treatment four-dimensional cone-beam computed tomography in volumetric-modulated arc therapy for stereotactic body radiation therapy. <i>Physica Medica</i> , 2019, 68, 10-16.	0.7	8
28	Single- Versus Multiple-Fraction Radiation Therapy for Painful Bone Metastases: A Systematic Review and Meta-analysis of Nonrandomized Studies. <i>Advances in Radiation Oncology</i> , 2019, 4, 706-715.	1.2	8
29	Enhanced Expression of IGFBP-3 Reduces Radiosensitivity and Is Associated with Poor Prognosis in Oral Squamous Cell Carcinoma. <i>Cancers</i> , 2020, 12, 494.	3.7	8
30	Dose-Function Histogram Evaluation Using 99mTc-GSA SPECT/CT Images for Stereotactic Body Radiation Therapy Planning for Hepatocellular Carcinoma Patients: A Dosimetric Parameter Comparison. <i>Anticancer Research</i> , 2018, 38, 1511-1516.	1.1	8
31	Radiation therapy for lymph node metastases from hepatocellular carcinoma. <i>Hepato-Gastroenterology</i> , 2009, 56, 476-80.	0.5	8
32	Comparison of rigid and deformable image registration for nasopharyngeal carcinoma radiotherapy planning with diagnostic position PET/CT. <i>Japanese Journal of Radiology</i> , 2020, 38, 256-264.	2.4	7
33	Predictors of the Predominance of NonIndex Pain After Palliative Radiation Therapy for Painful Tumors. <i>Advances in Radiation Oncology</i> , 2019, 4, 118-126.	1.2	6
34	Can MRI-derived depth of invasion predict nodal recurrence in oral tongue cancer?. <i>Oral Radiology</i> , 2021, 37, 641-646.	1.9	6
35	Concurrent chemoradiotherapy with S-1 in patients with stage III-IV oral squamous cell carcinoma: A retrospective analysis of nodal classification based on the neck node level. <i>Molecular and Clinical Oncology</i> , 2017, 7, 140-144.	1.0	5
36	Effect of metal-containing topical agents on surface doses received during external irradiation. <i>Journal of Radiation Research</i> , 2018, 59, 794-799.	1.6	5

#	ARTICLE	IF	CITATIONS
37	A neuropathic pain component as a predictor of improvement in pain interference after radiotherapy for painful tumors: A secondary analysis of a prospective observational study. <i>Clinical and Translational Radiation Oncology</i> , 2018, 12, 34-39.	1.7	5
38	High Spatial Resolution Digital Positron Emission Tomography Images With Dedicated Source-to-background Algorithm for Radiotherapy Planning. <i>Anticancer Research</i> , 2020, 40, 2567-2572.	1.1	5
39	Impact of four-dimensional cone-beam computed tomography on target localization for gastric mucosa-associated lymphoid tissue lymphoma radiotherapy: reducing planning target volume. <i>Radiation Oncology</i> , 2021, 16, 14.	2.7	5
40	Four-dimensional cone-beam computed tomography-guided radiotherapy for gastric lymphoma. <i>Japanese Journal of Radiology</i> , 2018, 36, 159-163.	2.4	4
41	Influence of pain duration on pain outcomes following palliative radiotherapy for painful tumors: the sooner the irradiation, the better?. <i>Strahlentherapie Und Onkologie</i> , 2021, 197, 916-925.	2.0	4
42	Palliative radiotherapy for painful lymph node metastases. <i>Radiation Oncology</i> , 2021, 16, 178.	2.7	4
43	Stereotactic Body Radiotherapy Based on ^{99m} Tc-GSA SPECT Image-guided Inverse Planning for Hepatocellular Carcinoma. <i>In Vivo</i> , 2020, 34, 3583-3588.	1.3	4
44	Influence of the treatment schedule on the physicians' decisions to refer bone metastases patients for palliative radiotherapy: a questionnaire survey of physicians in various specialties. <i>Nagoya Journal of Medical Science</i> , 2016, 78, 275-84.	0.3	4
45	A prospective comparison of adaptive and fixed boost plans in radiotherapy for glioblastoma. <i>Radiation Oncology</i> , 2022, 17, 40.	2.7	4
46	Plan Quality Comparisons Between 3D-CRT, IMRT, and VMAT Based on 4D-CT for Gastric MALT Lymphoma. <i>Anticancer Research</i> , 2021, 41, 3941-3947.	1.1	3
47	Radiotherapy for T3N0 glottic carcinoma without cord fixation: elective nodal irradiation or not?. <i>Oncotarget</i> , 2017, 8, 79761-79766.	1.8	3
48	Semi-automated prediction approach of target shifts using machine learning with anatomical features between planning and pretreatment CT images in prostate radiotherapy. <i>Journal of Radiation Research</i> , 2020, 61, 285-297.	1.6	3
49	Prevalence and risk factors of retro-styloid lymph node metastasis in oropharyngeal carcinoma. <i>Annals of Medicine</i> , 2022, 54, 436-441.	3.8	3
50	Concurrent Chemoradiotherapy With Docetaxel, Cisplatin, and 5-Fluorouracil for T3 N0 Glottic Carcinoma Without Vocal Cord Fixation. <i>Anticancer Research</i> , 2022, 42, 205-209.	1.1	3
51	Index and Nonindex Pain Endpoints in Radiation Therapy for Painful Tumors: A Secondary Analysis of a Prospective Observational Study. <i>Advances in Radiation Oncology</i> , 2020, 5, 1118-1125.	1.2	2
52	Prognostic value of parameters derived from white blood cell and differential counts in patients receiving palliative radiotherapy. <i>Molecular and Clinical Oncology</i> , 2016, 5, 241-246.	1.0	0
53	Improvement in pain interference after palliative radiotherapy for solid and hematologic painful tumors: a secondary analysis of a prospective observational study. <i>Japanese Journal of Clinical Oncology</i> , 2018, 48, 982-987.	1.3	0
54	QUAD shot: an effective cyclical hypofractionated palliative radiotherapy for salivary gland carcinoma. <i>BJR case Reports</i> , 2020, 6, 20190132.	0.2	0

#	ARTICLE	IF	CITATIONS
55	Abrupt Exacerbation of Atrial Functional Mitral Regurgitation During Emergence From General Anesthesia Following Transcatheter Aortic Valve Replacement. <i>A&A Practice</i> , 2020, 14, e01260.	0.4	0
56	Implementation of ^{99m} Tc-GSA SPECT Image-guided Inverse Planning into Palliative Radiotherapy for Diffuse Liver Metastases: A Novel Approach. <i>In Vivo</i> , 2022, 36, 1523-1526.	1.3	0