

# Orhan Onalan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2289335/publications.pdf>

Version: 2024-02-01

45  
papers

847  
citations

516215

16  
h-index

500791

28  
g-index

47  
all docs

47  
docs citations

47  
times ranked

1172  
citing authors

#	ARTICLE	IF	CITATIONS
1	Left Atrial Appendage Exclusion for Stroke Prevention in Patients With Nonrheumatic Atrial Fibrillation. <i>Stroke</i> , 2007, 38, 624-630.	1.0	119
2	Evaluation of Endothelial Dysfunction: Flow-Mediated Dilation. <i>Endothelium: Journal of Endothelial Cell Research</i> , 2008, 15, 157-163.	1.7	84
3	Meta-Analysis of Magnesium Therapy for the Acute Management of Rapid Atrial Fibrillation. <i>American Journal of Cardiology</i> , 2007, 99, 1726-1732.	0.7	82
4	Nonpharmacologic stroke prevention in atrial fibrillation. <i>Expert Review of Cardiovascular Therapy</i> , 2005, 3, 619-633.	0.6	48
5	Determinants of Pacemaker Dependency After Coronary and/or Mitral or Aortic Valve Surgery With Long-Term Follow-Up. <i>American Journal of Cardiology</i> , 2008, 101, 203-208.	0.7	44
6	Longitudinal Left Ventricular Function in Normotensive Prediabetics: A Tissue Doppler and Strain/Strain Rate Echocardiography Study. <i>Journal of the American Society of Echocardiography</i> , 2012, 25, 349-356.	1.2	39
7	Plasminogen activator inhibitor-1 4G4G genotype is associated with myocardial infarction but not with stable coronary artery disease. <i>Journal of Thrombosis and Thrombolysis</i> , 2008, 26, 211-217.	1.0	38
8	The Effect of Slow Coronary Flow on Right and Left Ventricular Performance. <i>Medical Principles and Practice</i> , 2014, 23, 34-39.	1.1	23
9	Complete Atrioventricular Block Associated With Concomitant Use of Metoprolol and Paroxetine. <i>Mayo Clinic Proceedings</i> , 2008, 83, 595-599.	1.4	22
10	Early Effects of Treatment with Nebivolol and Quinapril on Endothelial Function in Patients with Hypertension. <i>Endothelium: Journal of Endothelial Cell Research</i> , 2008, 15, 149-155.	1.7	21
11	The signal-averaged P-wave duration is longer in hypertensive patients with history of paroxysmal atrial fibrillation as compared to those without. <i>International Journal of Cardiology</i> , 2005, 103, 37-40.	0.8	19
12	Relation between fragmented QRS and collateral circulation in patients with chronic total occlusion without prior myocardial infarction. <i>Anatolian Journal of Cardiology</i> , 2011, 11, 300-4.	0.4	19
13	Heart Rate Variability and Heart Rate Turbulence in Hypothyroidism before and after Treatment. , 2011, 16, 344-350.		18
14	Frequency of fragmented QRS on ECG is increased in patients with rheumatoid arthritis without cardiovascular disease: a pilot study. <i>Modern Rheumatology</i> , 2012, 22, 238-242.	0.9	18
15	Relationship between red cell distribution width and echocardiographic parameters in patients with diastolic heart failure. <i>Kaohsiung Journal of Medical Sciences</i> , 2012, 28, 165-172.	0.8	18
16	Empiric slow pathway ablation in non-inducible supraventricular tachycardia. <i>International Journal of Cardiology</i> , 2015, 179, 417-420.	0.8	18
17	Aortic arch calcification is strongly associated with coronary artery calcification. <i>Vasa - European Journal of Vascular Medicine</i> , 2015, 44, 106-114.	0.6	17
18	Mildly Decreased Glomerular Filtration Rate Is Associated With Poor Coronary Collateral Circulation in Patients With Coronary Artery Disease. <i>Clinical Cardiology</i> , 2011, 34, 617-621.	0.7	16

#	ARTICLE	IF	CITATIONS
19	The protective effect of single dose tadalafil in contrast-induced nephropathy: An experimental study. <i>Anatolian Journal of Cardiology</i> , 2015, 15, 306-310.	0.5	16
20	Adrenal Adenoma Presenting with Torsade de Pointes. <i>Angiology</i> , 2002, 53, 471-474.	0.8	15
21	Frequency of fragmented QRS on ECG is increased in patients with rheumatoid arthritis without cardiovascular disease: a pilot study. <i>Modern Rheumatology</i> , 2012, 22, 238-242.	0.9	15
22	Radiotherapy Induces Development of Fragmented QRS in Patients with Breast Cancer. <i>Breast Care</i> , 2015, 10, 277-280.	0.8	14
23	Relationship between Aortic Arch Calcification, Detected by Chest X-Ray, and Renal Resistive Index in Patients with Hypertension. <i>Medical Principles and Practice</i> , 2019, 28, 133-140.	1.1	13
24	Association of Fragmented QRS with Subclinical Left Ventricular Dysfunction in Patients with Obstructive Sleep Apnea. <i>Medical Principles and Practice</i> , 2015, 24, 376-381.	1.1	12
25	Prophylaxis and management of postoperative atrial fibrillation. <i>Current Cardiology Reports</i> , 2005, 7, 382-390.	1.3	10
26	Onychomycosis is associated with subclinical atherosclerosis in patients with diabetes. <i>Vasa - European Journal of Vascular Medicine</i> , 2015, 44, 59-64.	0.6	10
27	Effects of prediabetes on coronary collateral circulation in patients with coronary artery disease. <i>Coronary Artery Disease</i> , 2011, 22, 233-237.	0.3	9
28	Antioxidant status and levels of antioxidant vitamins in coronary artery ectasia. <i>Coronary Artery Disease</i> , 2011, 22, 306-310.	0.3	9
29	Fragmented QRS Complexes are a Marker of Myocardial Fibrosis in Hypertensive Heart Disease. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2016, 44, 554-560.	0.6	9
30	Impact of cytokine gene polymorphism on cardiovascular risk in renal transplant recipients. <i>Transplant International</i> , 2005, 18, 681-689.	0.8	8
31	Pregbalin induced recurrent syncopal attacks with prolong <math>\langle scp \rangle QT \langle /scp \rangle</math> interval. <i>Annals of Noninvasive Electrocardiology</i> , 2018, 23, e12489.	0.5	7
32	Relationship between L-arginine/asymmetric dimethylarginine, homocysteine, folic acid, vitamin B levels, and coronary artery ectasia. <i>Coronary Artery Disease</i> , 2010, 21, 445-449.	0.3	4
33	Relationship between ST-Segment Shifts in Lead aVR and Coronary Complexity in Patients with Acute Coronary Syndrome. <i>Acta Cardiologica Sinica</i> , 2019, 35, 11-19.	0.1	4
34	Coronary and vertebral subclavian steal demonstrated by subclavian angiography. <i>European Heart Journal</i> , 2009, 30, 2469-2469.	1.0	3
35	Angioplasty induced myocardial ischaemia prolongs the signal-averaged P-wave duration in single vessel coronary artery disease. <i>Europace</i> , 2002, 4, 289-293.	0.7	2
36	The relationship between renal functions and thrombolysis in myocardial infarction frame count in patients with slow coronary flow. <i>Kaohsiung Journal of Medical Sciences</i> , 2011, 27, 55-58.	0.8	2

#	ARTICLE	IF	CITATIONS
37	A strong and reliable indicator for early postoperative major cardiac events after elective orthopedic surgery: Aortic arch calcification. <i>Heart and Lung: Journal of Acute and Critical Care</i> , 2019, 48, 446-451.	0.8	2
38	Assesment of Right Ventricular Diastolic Function in Chronic Obstructive Lung Disease. <i>Turk Toraks Dergisi</i> , 2013, 13, 152-157.	0.2	1
39	Uma CombinaÃ§Ã£o Fatal: Indometacina e Dabigatrana. <i>Arquivos Brasileiros De Cardiologia</i> , 2020, 114, 13-15.	0.3	1
40	Nebivolol and quinapril reduce p-wave duration and dispersion in hypertensive patients. <i>Indian Pacing and Electrophysiology Journal</i> , 2009, 9, 158-66.	0.3	1
41	Development of Pacemaker Lead Thrombosis in a Patient with Atrial Fibrillation during Apixaban Treatment. <i>The Journal of Tehran Heart Center</i> , 2019, 14, 183-186.	0.3	1
42	Transthoracic Echocardiographic View of Aortic Dissection, Massive Pericardial Effusion and Severe Left Ventricular Hypertrophy in a Young Patient with Uncontrolled Hypertension. <i>Echocardiography</i> , 2010, 27, 94-94.	0.3	0
43	Reply to the Letter by Ertem et al. Entitled 'Is Coronary Circulation the Most Important Parameter for Right Ventricular Functions?'. <i>Medical Principles and Practice</i> , 2015, 24, 295-295.	1.1	0
44	Transthoracic Echocardiographic and Angiographic View of a Coronary-to-Left Ventricular Fistula. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2012, 40, 561-561.	0.6	0
45	Bilateral Coronary-to-Pulmonary Artery Fistulas: A Rare Cause of Chest Pain and Pulmonary Hypertension. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2012, 40, 285-285.	0.6	0