

Moammir H Aziz

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2289812/publications.pdf>

Version: 2024-02-01

14
papers

1,232
citations

759233

12
h-index

1058476

14
g-index

14
all docs

14
docs citations

14
times ranked

1607
citing authors

#	ARTICLE	IF	CITATIONS
1	Chemoprevention of skin cancer by grape constituent resveratrol: relevance to human disease?. <i>FASEB Journal</i> , 2005, 19, 1193-1195.	0.5	209
2	Resveratrol-caused apoptosis of human prostate carcinoma LNCaP cells is mediated via modulation of phosphatidylinositol 3-kinase/Akt pathway and Bcl-2 family proteins. <i>Molecular Cancer Therapeutics</i> , 2006, 5, 1335-1341.	4.1	189
3	Plumbagin, a Medicinal Plant-Derived Naphthoquinone, Is a Novel Inhibitor of the Growth and Invasion of Hormone-Refractory Prostate Cancer. <i>Cancer Research</i> , 2008, 68, 9024-9032.	0.9	172
4	Modulations of critical cell cycle regulatory events during chemoprevention of ultraviolet B-mediated responses by resveratrol in SKH-1 hairless mouse skin. <i>Oncogene</i> , 2004, 23, 5151-5160.	5.9	143
5	Protein Kinase C μ Interacts with Signal Transducers and Activators of Transcription 3 (Stat3), Phosphorylates Stat3Ser727, and Regulates Its Constitutive Activation in Prostate Cancer. <i>Cancer Research</i> , 2007, 67, 8828-8838.	0.9	124
6	Prevention of Ultraviolet-B Radiation Damage by Resveratrol in Mouse Skin Is Mediated via Modulation in Survivin \uparrow . <i>Photochemistry and Photobiology</i> , 2005, 81, 25.	2.5	118
7	Protein Kinase C μ , which Sensitizes Skin to Sun's UV Radiation-Induced Cutaneous Damage and Development of Squamous Cell Carcinomas, Associates with Stat3. <i>Cancer Research</i> , 2007, 67, 1385-1394.	0.9	72
8	The Upregulation of Integrin α 2 β 2 (CD11d/CD18) on Inflammatory Macrophages Promotes Macrophage Retention in Vascular Lesions and Development of Atherosclerosis. <i>Journal of Immunology</i> , 2017, 198, 4855-4867.	0.8	56
9	Protein kinase C δ interacts with Stat3 and regulates its activation that is essential for the development of skin cancer. <i>Molecular Carcinogenesis</i> , 2007, 46, 646-653.	2.7	54
10	Protein Kinase C γ Overexpressing Transgenic Mice Are Resistant to Chemically but not to UV Radiation-Induced Development of Squamous Cell Carcinomas: A Possible Link to Specific Cytokines and Cyclooxygenase-2. <i>Cancer Research</i> , 2006, 66, 713-722.	0.9	35
11	Ultraviolet-B Radiation Causes an Upregulation of Survivin in Human Keratinocytes and Mouse Skin \uparrow . <i>Photochemistry and Photobiology</i> , 2004, 80, 602.	2.5	25
12	Protective molecular mechanisms of resveratrol in UVR-induced Skin carcinogenesis. <i>Photodermatology Photoimmunology and Photomedicine</i> , 2018, 34, 35-41.	1.5	14
13	Protein Kinase C μ Inhibits UVR-Induced Expression of FADD, an Adaptor Protein, Linked to both Fas- and TNFR1-Mediated Apoptosis. <i>Journal of Investigative Dermatology</i> , 2009, 129, 2011-2021.	0.7	13
14	Prevention of Ultraviolet-B Radiation Damage by Resveratrol in Mouse Skin Is Mediated via Modulation in Survivin \uparrow . <i>Photochemistry and Photobiology</i> , 2005, 81, 25-31.	2.5	8