

Christian Medom Madsen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2303562/publications.pdf>

Version: 2024-02-01

13
papers

588
citations

933447

10
h-index

1125743

13
g-index

13
all docs

13
docs citations

13
times ranked

921
citing authors

#	ARTICLE	IF	CITATIONS
1	Worldwide Prevalence of Familial Hypercholesterolemia. <i>Journal of the American College of Cardiology</i> , 2020, 75, 2553-2566.	2.8	304
2	Molecular Characterization of Oxysterol Binding to the Epstein-Barr Virus-induced Gene 2 (GPR183). <i>Journal of Biological Chemistry</i> , 2012, 287, 35470-35483.	3.4	46
3	Orthogeriatric Service Reduces Mortality in Patients With Hip Fracture. <i>Geriatric Orthopaedic Surgery and Rehabilitation</i> , 2016, 7, 67-73.	1.4	42
4	Temporal trends in hip fracture incidence, mortality, and morbidity in Denmark from 1999 to 2012. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2018, 89, 170-176.	3.3	42
5	Genetic Variants Associated With Increased Plasma Levels of Triglycerides, via Effects on the Lipoprotein Lipase Pathway, Increase Risk of Acute Pancreatitis. <i>Clinical Gastroenterology and Hepatology</i> , 2021, 19, 1652-1660.e6.	4.4	30
6	Preoperative factors associated with red blood cell transfusion in hip fracture patients. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2014, 134, 375-382.	2.4	24
7	Low Levels of Hemoglobin at Admission Are Associated With Increased 30-Day Mortality in Patients With Hip Fracture. <i>Geriatric Orthopaedic Surgery and Rehabilitation</i> , 2016, 7, 115-120.	1.4	23
8	Is It Time for New Thinking About High-Density Lipoprotein?. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018, 38, 484-486.	2.4	21
9	Hyperkalemia is Associated with Increased 30-Day Mortality in Hip Fracture Patients. <i>Calcified Tissue International</i> , 2017, 101, 9-16.	3.1	18
10	Temporal trends in the use of antithrombotics at admission. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2016, 87, 368-373.	3.3	14
11	Pre-fracture medication use as a predictor of 30-day mortality in hip fracture patients: an analysis of 141,201 patients. <i>HIP International</i> , 2020, 30, 101-106.	1.7	10
12	Centenarian hip fracture patients: a nationwide population-based cohort study of 507 patients. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2019, 90, 342-347.	3.3	9
13	Thyroid-stimulating hormone (TSH) is associated with 30-day mortality in hip fracture patients. <i>European Journal of Trauma and Emergency Surgery</i> , 2021, 47, 1081-1087.	1.7	5