

# Christian Medom Madsen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2303562/publications.pdf>

Version: 2024-02-01

13  
papers

588  
citations

933447

10  
h-index

1125743

13  
g-index

13  
all docs

13  
docs citations

13  
times ranked

921  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Thyroid-stimulating hormone (TSH) is associated with 30-day mortality in hip fracture patients. <i>European Journal of Trauma and Emergency Surgery</i> , 2021, 47, 1081-1087.   | 1.7 | 5         |
| 2  | Genetic Variants Associated With Increased Plasma Levels of Triglycerides, via Effects on the Lipoprotein Lipase Pathway, Increase Risk of Acute Pancreatitis. <i>Clinical Gastroenterology and Hepatology</i> , 2021, 19, 1652-1660.e6. | 4.4 | 30        |
| 3  | Pre-fracture medication use as a predictor of 30-day mortality in hip fracture patients: an analysis of 141,201 patients. <i>HIP International</i> , 2020, 30, 101-106.  | 1.7 | 10        |
| 4  | Worldwide Prevalence of Familial Hypercholesterolemia. <i>Journal of the American College of Cardiology</i> , 2020, 75, 2553-2566.   | 2.8 | 304       |
| 5  | Centenarian hip fracture patients: a nationwide population-based cohort study of 507 patients. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2019, 90, 342-347.  | 3.3 | 9         |
| 6  | Is It Time for New Thinking About High-Density Lipoprotein?. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018, 38, 484-486.  | 2.4 | 21        |
| 7  | Temporal trends in hip fracture incidence, mortality, and morbidity in Denmark from 1999 to 2012. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2018, 89, 170-176.   | 3.3 | 42        |
| 8  | Hyperkalemia is Associated with Increased 30-Day Mortality in Hip Fracture Patients. <i>Calcified Tissue International</i> , 2017, 101, 9-16.  | 3.1 | 18        |
| 9  | Orthogeriatric Service Reduces Mortality in Patients With Hip Fracture. <i>Geriatric Orthopaedic Surgery and Rehabilitation</i> , 2016, 7, 67-73.  | 1.4 | 42        |
| 10 | Low Levels of Hemoglobin at Admission Are Associated With Increased 30-Day Mortality in Patients With Hip Fracture. <i>Geriatric Orthopaedic Surgery and Rehabilitation</i> , 2016, 7, 115-120.  | 1.4 | 23        |
| 11 | Temporal trends in the use of antithrombotics at admission. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2016, 87, 368-373.   | 3.3 | 14        |
| 12 | Preoperative factors associated with red blood cell transfusion in hip fracture patients. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2014, 134, 375-382.  | 2.4 | 24        |
| 13 | Molecular Characterization of Oxysterol Binding to the Epstein-Barr Virus-induced Gene 2 (GPR183). <i>Journal of Biological Chemistry</i> , 2012, 287, 35470-35483.  | 3.4 | 46        |