Gyung-Jae Oh

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2331758/publications.pdf

Version: 2024-02-01

687363 752698 27 466 13 20 citations h-index g-index papers 27 27 27 743 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Predictors of functional level and quality of life at 6Âmonths after a first-ever stroke: the KOSCO study. Journal of Neurology, 2016, 263, 1166-1177.	3.6	52
2	Effect of Cognitive Reserve on Risk of Cognitive Impairment and Recovery After Stroke. Stroke, 2020, 51, 99-107.	2.0	46
3	Representative levels of blood lead, mercury, and urinary cadmium in youth: Korean Environmental Health Survey in Children and Adolescents (KorEHS-C), 2012–2014. International Journal of Hygiene and Environmental Health, 2016, 219, 412-418.	4.3	40
4	Korean Stroke Cohort for functioning and rehabilitation (KOSCO): study rationale and protocol of a multi-centre prospective cohort study. BMC Neurology, 2015, 15, 42.	1.8	37
5	Association of current phthalate exposure with neurobehavioral development in a national sample. International Journal of Hygiene and Environmental Health, 2016, 219, 364-371.	4.3	34
6	Risk Factors and Functional Impact of Medical Complications in Stroke. Annals of Rehabilitation Medicine, 2017, 41, 753.	1.6	34
7	HbA1c is significantly associated with arterial stiffness but not with carotid atherosclerosis in a community-based population without type 2 diabetes: The Dong-gu study. Atherosclerosis, 2016, 247, 1-6.	0.8	26
8	Health Indicators Related to Disease, Death, and Reproduction. Journal of Preventive Medicine and Public Health, 2019, 52, 14-20.	1.9	23
9	INHIBITORY EFFECT OF IMMUNOGLOBULIN E PRODUCTION BY JIN-DEUK-CHAL (SIEGESBECKIA ORIENTALIS). Immunopharmacology and Immunotoxicology, 2001, 23, 555-563.	2.4	17
10	Public Awareness of Stroke and Its Predicting Factors in Korea: a National Public Telephone Survey, 2012 and 2014. Journal of Korean Medical Science, 2016, 31, 1703.	2.5	17
11	Factors influencing return to work after stroke: the Korean Stroke Cohort for Functioning and Rehabilitation (KOSCO) Study. BMJ Open, 2019, 9, e028673.	1.9	17
12	Normative and mean carotid intima-media thickness values according to metabolic syndrome in Koreans: The Namwon Study. Atherosclerosis, 2014, 234, 230-236.	0.8	16
13	Gender-Specific Factors Associated with Suicide Attempts among the Community-Dwelling General Population with Suicidal Ideation: the 2013 Korean Community Health Survey. Journal of Korean Medical Science, 2016, 31, 2010.	2.5	16
14	Status Of Dysphagia After Ischemic Stroke: A Korean Nationwide Study. Archives of Physical Medicine and Rehabilitation, 2021, 102, 2343-2352.e3.	0.9	12
15	Revisiting the Proportional Recovery Model in View of the Ceiling Effect of Fugl-Meyer Assessment. Stroke, 2021, 52, 3167-3175.	2.0	12
16	Impact of central facial palsy and dysarthria on quality of life in patients with stroke: TheÂKOSCO study. NeuroRehabilitation, 2016, 39, 253-259.	1.3	11
17	Differences in agricultural activities related to incidence of scrub typhus between Korea and Japan. Epidemiology and Health, 2017, 39, e2017051.	1.9	11
18	Differences in the awareness of stroke symptoms and emergency response by occupation in the Korean general population. PLoS ONE, 2019, 14, e0218608.	2.5	8

#	Article	IF	CITATIONS
19	The Prevalence of Musculoskeletal Symptoms and the Ergonomic Risk Factors among Oriental Melon-growing Farmers. Korean Journal of Occupational and Environmental Medicine, 2011, 23, 1.	0.4	8
20	Development and validation of a prediction model for home discharge in patients with moderate stroke: The Korean stroke cohort for functioning and rehabilitation study. Topics in Stroke Rehabilitation, 2020, 27, 453-461.	1.9	7
21	Association between Scrub Typhus Outbreaks and Meteorological Factors in Jeollabuk-do Province. Korean Journal of Environmental Health Sciences, 2016, 42, 41-52.	0.3	6
22	Factors Associated to Returning Home in the First Year after Stroke. Brain & Neurorehabilitation, 2020, 13, .	1.0	4
23	Post-stroke Hyperglycemia in Non-diabetic Ischemic Stroke is Related With Worse Functional Outcome: A Cohort Study. Annals of Rehabilitation Medicine, 2021, 45, 359-367.	1.6	4
24	Factors Associated with Changes in Functional Independence after Six Months of Ischemic Stroke. Brain & Neurorehabilitation, 2020, 13, .	1.0	3
25	Continuous Abstinence Rates from Smoking Over 12 Months according to the Frequency of Participation in a Hospital-based Smoking Cessation Program among Patients Discharged after Acute Myocardial Infarction. Korean Journal of Health Promotion, 2016, 16, 48.	0.2	2
26	Simultaneous Clustering and Classification of Function Recovery Patterns of Ischemic Stroke. Journal of Medical Imaging and Health Informatics, 2020, 10, 1401-1407.	0.3	2
27	Understanding of the Lower Extremity Motor Recovery After First-Ever Ischemic Stroke. Stroke, 0, , .	2.0	1