

Ludmila Matysova

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2336800/ludmila-matysova-publications-by-citations.pdf>
Version: 2024-04-11

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

27 papers	721 citations	11 h-index	26 g-index
27 ext. papers	774 ext. citations	3.9 avg, IF	3.53 L-index

#	Paper	IF	Citations
27	Advantages of application of UPLC in pharmaceutical analysis. <i>Talanta</i> , 2006 , 68, 908-18	6.2	339
26	Simultaneous HPLC determination of ketoprofen and its degradation products in the presence of preservatives in pharmaceuticals. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2004 , 36, 625-9	3.5	43
25	Determination of estradiol and its degradation products by liquid chromatography. <i>Journal of Chromatography A</i> , 2006 , 1119, 216-23	4.5	42
24	Comparison of performance of C18 monolithic rod columns and conventional C18 particle-packed columns in liquid chromatographic determination of Estrogel and Ketoprofen gel. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2004 , 813, 191-7	3.2	41
23	HPLC determination of estradiol, its degradation product, and preservatives in new topical formulation Estrogel HBF. <i>Analytical and Bioanalytical Chemistry</i> , 2004 , 379, 781-7	4.4	36
22	Development and validation of HPLC method for determination of indomethacin and its two degradation products in topical gel. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2005 , 37, 899-905	3.5	31
21	HPLC determination of chlorhexidine gluconate and p-chloroaniline in topical ointment. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2007 , 43, 1169-73	3.5	26
20	Determination of methylparaben, propylparaben, triamcinolone acetonide and its degradation product in a topical cream by RP-HPLC. <i>Analytical and Bioanalytical Chemistry</i> , 2003 , 376, 440-3	4.4	26
19	Development and validation of HPLC method for determination of clotrimazole and its two degradation products in spray formulation. <i>Talanta</i> , 2007 , 73, 483-9	6.2	22
18	Separation and determination of terbinafine and its four impurities of similar structure using simple RP-HPLC method. <i>Talanta</i> , 2006 , 68, 713-20	6.2	18
17	High-resolution monolithic columns--a new tool for effective and quick separation. <i>Analytical and Bioanalytical Chemistry</i> , 2013 , 405, 2255-63	4.4	14
16	Optimisation of an HPLC method for the simultaneous determination of pyrantel pamoate, praziquantel, fenbendazole, oxfendazole and butylhydroxyanisole using a phenyl stationary phase. <i>Analytical Methods</i> , 2012 , 4, 1592	3.2	11
15	Advantages of pentafluorophenylpropyl stationary phase over conventional C18 stationary phase--application to analysis of triamcinolone acetonide. <i>Talanta</i> , 2008 , 76, 597-601	6.2	11
14	Application of monolithic columns in pharmaceutical analysis. Determination of indomethacin and its degradation products. <i>Journal of Separation Science</i> , 2009 , 32, 2786-92	3.4	10
13	Simultaneous Determination of Propranolol Hydrochloride and Sodium Benzoate in Oral Liquid Preparations by HPLC. <i>Chromatographia</i> , 2013 , 76, 1553-1558	2.1	9
12	Application of HILIC stationary phase to determination of dimethindene maleate in topical gel. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2009 , 50, 23-6	3.5	8
11	HPLC determination of calcium pantothenate and two preservatives in topical cream. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2006 , 41, 671-5	3.5	6

10	Furosemide ethanol-free oral solutions for paediatric use: formulation, HPLC method and stability study. <i>European Journal of Hospital Pharmacy</i> , 2018 , 25, 144-149	1.6	6
9	Development of a gradient HPLC method for the simultaneous determination of sotalol and sorbate in oral liquid preparations using solid core stationary phase. <i>Journal of Analytical Methods in Chemistry</i> , 2015 , 2015, 806736	2	5
8	Terbinafine-loaded branched PLGA-based cationic nanoparticles with modifiable properties. <i>Pharmaceutical Development and Technology</i> , 2019 , 24, 1308-1316	3.4	3
7	Determination of urinary retinol and creatinine as an early sensitive marker of renal dysfunction. <i>Journal of Chromatography A</i> , 2019 , 1607, 460390	4.5	3
6	Extemporaneous sotalol hydrochloride oral solutions for use in paediatric cardiology: formulation and stability study. <i>European Journal of Hospital Pharmacy</i> , 2016 , 23, 33-37	1.6	3
5	Fast determination of omeprazole in extemporaneous suspensions used in paediatrics and stability studies. <i>Analytical Methods</i> , 2019 , 11, 517-523	3.2	2
4	Development of Novel Stability-Indicating Method for the Determination of Dimethindene Maleate and Its Impurities. <i>Chromatographia</i> , 2013 , 76, 1545-1551	2.1	2
3	Could urinary retinol be used as a new biomarker of kidney damage?. <i>TrAC - Trends in Analytical Chemistry</i> , 2017 , 95, 57-61	14.6	2
2	Chromatographic determination of active compounds in topical formulations. <i>Analytical Methods</i> , 2012 , 4, 1525	3.2	1
1	Omeprazole suspension 2 mg/ml - a new dosage form for pediatric use. <i>Praktická Farmazie</i> , 2017 , 13, 18-20	0.1	1