Olga Heidingsfeld

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2337859/publications.pdf

Version: 2024-02-01

1040056 940533 20 265 9 16 citations g-index h-index papers 20 20 20 357 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Structural determinants for subnanomolar inhibition of the secreted aspartic protease Sapp1p from Candida parapsilosis. Journal of Enzyme Inhibition and Medicinal Chemistry, 2021, 36, 914-921.	5.2	3
2	Cellular Localization of Carbonic Anhydrase Nce103p in Candida albicans and Candida parapsilosis. International Journal of Molecular Sciences, 2020, 21, 850.	4.1	1
3	PHO15 genes of Candida albicans and Candida parapsilosis encode HAD-type phosphatases dephosphorylating 2-phosphoglycolate. FEMS Yeast Research, 2019, 19, .	2.3	2
4	Functional Characterization of Secreted Aspartyl Proteases in Candida parapsilosis. MSphere, 2019, 4, .	2.9	29
5	Crystal structure of carbonic anhydrase CaNce103p from the pathogenic yeast Candida albicans. BMC Structural Biology, 2018, 18, 14.	2.3	13
6	Atomic resolution crystal structure of Sapp2p, a secreted aspartic protease from (i> Candida parapsilosis (i>. Acta Crystallographica Section D: Biological Crystallography, 2015, 71, 2494-2504.	2.5	9
7	Two SAPP2 gene homologs are present in Candida parapsilosis genome. Folia Microbiologica, 2015, 60, 373-374.	2.3	3
8	Intracellular aspartic proteinase Apr1p of Candida albicans is required for morphological transition under nitrogen-limited conditions but not for macrophage killing. Folia Microbiologica, 2014, 59, 485-493.	2.3	6
9	î"12-Fatty Acid Desaturase from Candida parapsilosis Is a Multifunctional Desaturase Producing a Range of Polyunsaturated and Hydroxylated Fatty Acids. PLoS ONE, 2014, 9, e93322.	2.5	27
10	Saccharomyces cerevisiae can secrete Sapp1p proteinase of Candida parapsilosis but cannot use it for efficient nitrogen acquisition. Journal of Microbiology, 2013, 51, 336-344.	2.8	1
11	The crystal structure of protease Sapp1p from <i>Candida parapsilosis</i> in complex with the HIV protease inhibitor ritonavir. Journal of Enzyme Inhibition and Medicinal Chemistry, 2012, 27, 160-165.	5 . 2	8
12	Nitrogen source and growth stage of Candida albicans influence expression level of vacuolar aspartic protease Apr1p and carboxypeptidase Cpy1p. Canadian Journal of Microbiology, 2012, 58, 678-681.	1.7	7
13	Raman Microspectroscopy of the Yeast Vacuoles. Spectroscopy, 2012, 27, 503-507.	0.8	8
14	Evidence for the presence of proteolytically active secreted aspartic proteinase 1 of <i>Candida parapsilosis</i> in the cell wall. Protein Science, 2011, 20, 2004-2012.	7.6	8
15	Two aspartic proteinases secreted by the pathogenic yeast Candida parapsilosis differ in expression pattern and catalytic properties. Biological Chemistry, 2009, 390, 259-68.	2.5	26
16	The crystal structure of the secreted aspartic protease 1 from Candida parapsilosis in complex with pepstatin A. Journal of Structural Biology, 2009, 167, 145-152.	2.8	19
17	Secreted proteins of Candida albicans. Frontiers in Bioscience - Landmark, 2008, Volume, 7227.	3.0	35
18	Cloning and characterization of Sapp2p, the second aspartic proteinase isoenzyme fromCandida parapsilosis. FEMS Yeast Research, 2006, 6, 1018-1026.	2.3	33

#	Article	lF	CITATIONS
19	Structure-based specificity mapping of secreted aspartic proteases of Candida parapsilosis, Candida albicans, and Candida tropicalis using peptidomimetic inhibitors and homology modeling. Biological Chemistry, 2006, 387, 1247-1254.	2.5	8
20	The precursor of secreted aspartic proteinase Sapp1p from Candida parapsilosis can be activated both autocatalytically and by a membrane-bound processing proteinase. Biological Chemistry, 2005, 386, 791-9.	2.5	19