

# Ananth Shankar

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2339690/publications.pdf>

Version: 2024-02-01

26  
papers

518  
citations

758635

12  
h-index

676716

22  
g-index

26  
all docs

26  
docs citations

26  
times ranked

731  
citing authors

#	ARTICLE	IF	CITATIONS
1	Resection alone in 58 children with limited stage, lymphocyte-predominant Hodgkin lymphoma—experience from the European network group on pediatric Hodgkin lymphoma. <i>Cancer</i> , 2007, 110, 179-185.	2.0	123
2	Treatment outcome after low intensity chemotherapy [CVP] in children and adolescents with early stage nodular lymphocyte predominant Hodgkin lymphoma—An Anglo-French collaborative report. <i>European Journal of Cancer</i> , 2012, 48, 1700-1706.	1.3	55
3	18F-Fluoroethylcholine (18F-Cho) PET/MRI Functional Parameters in Pediatric Astrocytic Brain Tumors. <i>Clinical Nuclear Medicine</i> , 2015, 40, e40-e45.	0.7	41
4	Role of FDG PET in the management of childhood lymphomas—case proven or is the jury still out?. <i>European Journal of Cancer</i> , 2008, 44, 663-673.	1.3	37
5	Quantitative diffusion weighted MRI: A functional biomarker of nodal disease in Hodgkin lymphoma?. <i>Cancer Biomarkers</i> , 2011, 7, 249-259.	0.8	36
6	Nodular lymphocyte predominant Hodgkin lymphoma in children and adolescents—a comprehensive review of biology, clinical course and treatment options. <i>British Journal of Haematology</i> , 2012, 159, 288-298.	1.2	36
7	Whole-body MRI for staging and interim response monitoring in paediatric and adolescent Hodgkin lymphoma: a comparison with multi-modality reference standard including 18F-FDG-PET-CT. <i>European Radiology</i> , 2019, 29, 202-212.	2.3	29
8	Imaging characteristics of H3 K27M histone-mutant diffuse midline glioma in teenagers and adults. <i>Quantitative Imaging in Medicine and Surgery</i> , 2021, 11, 43-56.	1.1	21
9	68Ga-DOTATATE and 123I-mIBG as imaging biomarkers of disease localisation in metastatic neuroblastoma: implications for molecular radiotherapy. <i>Nuclear Medicine Communications</i> , 2020, 41, 1169-1177.	0.5	21
10	The use of multiparametric 18F-fluoro-l-3,4-dihydroxy-phenylalanine PET/MRI in post-therapy assessment of patients with gliomas. <i>Nuclear Medicine Communications</i> , 2020, 41, 517-525.	0.5	19
11	Predicting outcome in childhood diffuse midline gliomas using magnetic resonance imaging based texture analysis. <i>Journal of Neuroradiology</i> , 2021, 48, 243-247.	0.6	16
12	Treatment outcome in children and adolescents with relapsed Hodgkin lymphoma - results of the UK HD3 relapse treatment strategy. <i>British Journal of Haematology</i> , 2014, 165, 534-544.	1.2	13
13	Clinical outcome in children and adolescents with Hodgkin lymphoma after treatment with chemotherapy alone—The results of the United Kingdom HD3 national cohort trial. <i>European Journal of Cancer</i> , 2012, 48, 108-113.	1.3	12
14	Evaluation of treatment response using integrated 18F-labeled choline positron emission tomography/magnetic resonance imaging in adolescents with intracranial non-germinomatous germ cell tumours. <i>Pediatric Blood and Cancer</i> , 2015, 62, 1661-1663.	0.8	11
15	Treatment outcome with a selective RET tyrosine kinase inhibitor selipercatinib in children with multiple endocrine neoplasia type 2 and advanced medullary thyroid carcinoma. <i>European Journal of Cancer</i> , 2021, 158, 38-46.	1.3	9
16	Management of children and adults with all stages of nodular lymphocyte predominant Hodgkin lymphoma—A consensus-based position paper from the Hodgkin lymphoma subgroup of the UK National Cancer Research Institute. <i>British Journal of Haematology</i> , 2022, 197, 679-690.	1.2	9
17	Preferred place of death in paediatric, teenage and young adult haemato-oncology patients: a retrospective review. <i>BMJ Supportive and Palliative Care</i> , 2020, , bmjspcare-2019-002005.	0.8	8
18	Practice patterns for the management of nodular lymphocyte-predominant Hodgkin lymphoma (NLPHL): an international survey by the Global NLPHL One Working Group (GLOW). <i>Leukemia and Lymphoma</i> , 2022, , 1-4.	0.6	8

#	ARTICLE	IF	CITATIONS
19	Hybrid PETâ€“MRI Imaging in Paediatric and TYA Brain Tumours: Clinical Applications and Challenges. Journal of Personalized Medicine, 2020, 10, 218.	1.1	7
20	Whole-body magnetic resonance imaging in paediatric Hodgkin lymphoma â€” evaluation of quantitative magnetic resonance metrics for nodal staging. Pediatric Radiology, 2019, 49, 1285-1298.	1.1	4
21	Variant histology nodular lymphocyte predominant Hodgkin lymphoma â€” a route to transformation?. British Journal of Haematology, 2018, 181, 403-406.	1.2	2
22	RADI-06. CORRELATION BETWEEN APT-CEST AND 18F-CHOLINE PET IN GLIOMA AT 3T. Neuro-Oncology, 2018, 20, i170-i171.	0.6	1
23	Multiparametric 18F-Choline PET/MR in childhood and teenage-young adult gliomas: assessment of suspected disease progression. Neuro-Oncology, 2019, 21, iv5-iv5.	0.6	0
24	1343â€”...Discussing organ and tissue donation by children with cancer; is there a training need?. , 2021, , .		0
25	DIPG-37. PREDICTING OUTCOME IN CHILDHOOD DIFFUSE MIDLINE GLIOMAS USING MAGNETIC RESONANCE IMAGING BASED TEXTURE ANALYSIS. Neuro-Oncology, 2020, 22, iii294-iii294.	0.6	0
26	RONC-01. A 10 year, single institution experience of re-irradiation for paediatric intracranial tumours. Neuro-Oncology, 2022, 24, i176-i176.	0.6	0