Matthew P Fox

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/2340399/matthew-p-fox-publications-by-year.pdf

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

260 9,543 51 90 h-index g-index citations papers 6.38 11,082 273 5.5 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
2 60	Virologic non-suppression and early loss to follow up among pregnant and non-pregnant adolescents aged 15-19 years initiating antiretroviral therapy in South Africa: a retrospective cohort study <i>Journal of the International AIDS Society</i> , 2022 , 25, e25870	5.4	1
259	Attrition from HIV care among youth initiating ART in youth-only clinics compared with general primary healthcare clinics in Khayelitsha, South Africa: a matched propensity score analysis <i>Journal of the International AIDS Society</i> , 2022 , 25, e25854	5.4	0
258	Validation of self-reported opioid agonist treatment among people who inject drugs using prescription dispensation records. <i>Epidemiology</i> , 2021 , 33,	3.1	2
257	Retention in care and viral suppression after same-day ART initiation: One-year outcomes of the SLATE I and II individually randomized clinical trials in South Africa. <i>Journal of the International AIDS Society</i> , 2021 , 24, e25825	5.4	O
256	Lash et al. Respond to "Better Bias Analysis" and "Toward Better Bias Analysis". <i>American Journal of Epidemiology</i> , 2021 , 190, 1622-1624	3.8	O
255	Fast-track treatment initiation counselling in South Africa: A cost-outcomes analysis. <i>PLoS ONE</i> , 2021 , 16, e0248551	3.7	0
254	Adverse psychosocial factors in pregnancy and preterm delivery. <i>Paediatric and Perinatal Epidemiology</i> , 2021 , 35, 519-529	2.7	1
253	Bias Analysis Gone Bad. American Journal of Epidemiology, 2021, 190, 1604-1612	3.8	7
252	A systematic review of quantitative bias analysis applied to epidemiological research. <i>International Journal of Epidemiology</i> , 2021 , 50, 1708-1730	7.8	2
251	A comorbid mental disorder paradox: Using causal diagrams to understand associations between posttraumatic stress disorder and suicide. <i>Psychological Trauma: Theory, Research, Practice, and Policy</i> , 2021 , 13, 725-729	7.8	2
250	The revolving door of HIV care: Revising the service delivery cascade to achieve the UNAIDS 95-95-95 goals. <i>PLoS Medicine</i> , 2021 , 18, e1003651	11.6	10
249	Multimonth dispensing of up to 6 months of antiretroviral therapy in Malawi and Zambia (INTERVAL): a cluster-randomised, non-blinded, non-inferiority trial. <i>The Lancet Global Health</i> , 2021 , 9, e628-e638	13.6	15
248	Tracing people living with HIV who are lost to follow-up at ART programs in Southern Africa: A sampling-based cohort study in six countries. <i>Clinical Infectious Diseases</i> , 2021 ,	11.6	2
247	Will Podcasting and Social Media Replace Journals and Traditional Science Communication? No, but. <i>American Journal of Epidemiology</i> , 2021 , 190, 1625-1631	3.8	1
246	An underappreciated misclassification mechanism: implications of nondifferential dependent misclassification of covariate and exposure. <i>Annals of Epidemiology</i> , 2021 , 58, 104-123	6.4	1
245	Elevated serum progesterone during in vitro fertilization treatment and the risk of ischemic placental disease. <i>Pregnancy Hypertension</i> , 2021 , 24, 7-12	2.6	1
244	Assessing knowledge, attitudes, and practices towards causal directed acyclic graphs: a qualitative research project. <i>European Journal of Epidemiology</i> , 2021 , 36, 659-667	12.1	1

243	Iron status and self-reported fatigue in blood donors. <i>Transfusion</i> , 2021 , 61, 124-133	2.9	2
242	Simulation as a Tool for Teaching and Learning Epidemiologic Methods. <i>American Journal of Epidemiology</i> , 2021 , 190, 900-907	3.8	1
241	Use of directed acyclic graphs (DAGs) to identify confounders in applied health research: review and recommendations. <i>International Journal of Epidemiology</i> , 2021 , 50, 620-632	7.8	45
240	Psychotropic medication use during pregnancy and gestational age at delivery. <i>Annals of Epidemiology</i> , 2021 , 53, 34-41.e2	6.4	Ο
239	Patient Perspectives of Quality of the Same-Day Antiretroviral Therapy Initiation Process in Gauteng Province, South Africa: Qualitative Dominant Mixed-Methods Analysis of the SLATE II Trial. <i>Patient</i> , 2021 , 14, 175-186	3.7	2
238	Concerns About the Special Article on Hydroxychloroquine and Azithromycin in High-Risk Outpatients With COVID-19. <i>American Journal of Epidemiology</i> , 2021 , 190, 491-495	3.8	5
237	Health provider perspectives on the implementation of the same-day-ART initiation policy in the Gauteng province of South Africa. <i>Health Research Policy and Systems</i> , 2021 , 19, 2	3.7	1
236	Understanding the Reasons for Deferring ART Among Patients Diagnosed Under the Same-Day-ART Policy in Johannesburg, South Africa. <i>AIDS and Behavior</i> , 2021 , 25, 2779-2792	4.3	2
235	Addressing Measurement Error in Random Forests Using Quantitative Bias Analysis. <i>American Journal of Epidemiology</i> , 2021 , 190, 1830-1840	3.8	10
234	Attrition in HIV care following HIV diagnosis: a comparison of the pre-UTT and UTT eras in South Africa. <i>Journal of the International AIDS Society</i> , 2021 , 24, e25652	5.4	5
233	Variation in HIV care and treatment outcomes by facility in South Africa, 2011-2015: A cohort study. <i>PLoS Medicine</i> , 2021 , 18, e1003479	11.6	3
232	Regression discontinuity analysis demonstrated varied effect of Treat-All on CD4 testing among Southern African countries. <i>Journal of Clinical Epidemiology</i> , 2021 , 140, 101-110	5.7	
231	Mortality following workplace injury: Quantitative bias analysis. <i>Annals of Epidemiology</i> , 2021 , 64, 155-7	16604	0
230	Multiple Bias Modeling. Statistics in the Health Sciences, 2021 , 415-440		
229	Misclassification. Statistics in the Health Sciences, 2021 , 141-195		
228	Introduction, Objectives, and an Alternative. Statistics in the Health Sciences, 2021, 1-24		O
227	Preparing for Probabilistic Bias Analysis. Statistics in the Health Sciences, 2021, 197-231		
226	Best Practices for Quantitative Bias Analysis. <i>Statistics in the Health Sciences</i> , 2021 , 441-452		

A Guide to Implementing Quantitative Bias Analysis. Statistics in the Health Sciences, **2021**, 25-55

224	Retention in care and viral suppression in differentiated service delivery models for HIV treatment delivery in sub-Saharan Africa: a rapid systematic review. <i>Journal of the International AIDS Society</i> , 2020 , 23, e25640	5.4	19
223	PS-SiZer map to investigate significant features of body-weight profile changes in HIV infected patients in the IeDEA Collaboration. <i>PLoS ONE</i> , 2020 , 15, e0220165	3.7	
222	HIV Treatment Outcomes Among Patients Initiated on Antiretroviral Therapy Pre and Post-Universal Test and Treat Guidelines in South Africa. <i>Therapeutics and Clinical Risk Management</i> , 2020 , 16, 169-180	2.9	8
221	Commentary: The value of E-values and why they are not enough. <i>International Journal of Epidemiology</i> , 2020 , 49, 1505-1506	7.8	3
220	Common misconceptions about validation studies. <i>International Journal of Epidemiology</i> , 2020 , 49, 1392	- 1 .3896	15
219	What the hashtag? Using twitter and podcasting to extend your scientific reach. <i>Paediatric and Perinatal Epidemiology</i> , 2020 , 34, 553-555	2.7	1
218	Quantitative bias analysis for study and grant planning. <i>Annals of Epidemiology</i> , 2020 , 43, 32-36	6.4	5
217	Flexibly Accounting for Exposure Misclassification With External Validation Data. <i>American Journal of Epidemiology</i> , 2020 , 189, 850-860	3.8	1
216	A Clinical Prediction Score Including Trial of Antibiotics and C-Reactive Protein to Improve the Diagnosis of Tuberculosis in Ambulatory People With HIV. <i>Open Forum Infectious Diseases</i> , 2020 , 7, of 25-	43	3
215	Characterizing the double-sided cascade of care for adolescents living with HIV transitioning to adulthood across Southern Africa. <i>Journal of the International AIDS Society</i> , 2020 , 23, e25447	5.4	4
214	Using a Self-Administered Electronic Adherence Questionnaire to Identify Poor Adherence Amongst Adolescents and Young Adults on First-Line Antiretroviral Therapy in Johannesburg, South Africa. <i>Patient Preference and Adherence</i> , 2020 , 14, 133-151	2.4	1
213	The Impact of Delayed Switch to Second-Line Antiretroviral Therapy on Mortality, Depending on Definition of Failure Time and CD4 Count at Failure. <i>American Journal of Epidemiology</i> , 2020 , 189, 811-8	1 3 9 ⁸	10
212	A clinical algorithm for same-day HIV treatment initiation in settings with high TB symptom prevalence in South Africa: The SLATE II individually randomized clinical trial. <i>PLoS Medicine</i> , 2020 , 17, e1003226	11.6	3
211	Recording of HIV viral loads and viral suppression in South African patients receiving antiretroviral treatment: a multicentre cohort study. <i>Antiviral Therapy</i> , 2020 , 25, 257-266	1.6	2
210	Using intervention mapping in motivational interviewing training to improve ART uptake in Gauteng, South Africa. <i>Journal of Health Psychology</i> , 2020 , 1359105320962241	3.1	
209	Will differentiated care for stable HIV patients reduce healthcare systems costs?. <i>Journal of the International AIDS Society</i> , 2020 , 23, e25541	5.4	3
208	Impact of Viral Load Monitoring on Retention and Viral Suppression: A Regression Discontinuity Analysis of South Africaß National Laboratory Cohort. <i>American Journal of Epidemiology</i> , 2020 , 189, 149	2 ²⁻⁸ 501	1

(2019-2020)

207	"Patients are not the same, so we cannot treat them the same" - A qualitative content analysis of provider, patient and implementer perspectives on differentiated service delivery models for HIV treatment in South Africa. <i>Journal of the International AIDS Society</i> , 2020 , 23, e25544	5.4	11
206	Delays in repeat HIV viral load testing for those with elevated viral loads: a national perspective from South Africa. <i>Journal of the International AIDS Society</i> , 2020 , 23, e25542	5.4	7
205	Trends in CD4 and viral load testing 2005 to 2018: multi-cohort study of people living with HIV in Southern Africa. <i>Journal of the International AIDS Society</i> , 2020 , 23, e25546	5.4	7
204	Comparison of pregnancy outcomes following preimplantation genetic testing for aneuploidy using a matched propensity score design. <i>Human Reproduction</i> , 2020 , 35, 2356-2364	5.7	6
203	Prevalence of TB symptoms, diagnosis and treatment among people living with HIV (PLHIV) not on ART presenting at outpatient clinics in South Africa and Kenya: baseline results from a clinical trial. <i>BMJ Open</i> , 2020 , 10, e035794	3	3
202	The Critical Importance of Asking Good Questions: The Role of Epidemiology Doctoral Training Programs. <i>American Journal of Epidemiology</i> , 2020 , 189, 261-264	3.8	8
201	Impact of the test and treat policy on delays in antiretroviral therapy initiation among adult HIV positive patients from six clinics in Johannesburg, South Africa: results from a prospective cohort study. <i>BMJ Open</i> , 2020 , 10, e030228	3	11
2 00	PS-SiZer map to investigate significant features of body-weight profile changes in HIV infected patients in the IeDEA Collaboration 2020 , 15, e0220165		
199	PS-SiZer map to investigate significant features of body-weight profile changes in HIV infected patients in the IeDEA Collaboration 2020 , 15, e0220165		
198	PS-SiZer map to investigate significant features of body-weight profile changes in HIV infected patients in the IeDEA Collaboration 2020 , 15, e0220165		
197	PS-SiZer map to investigate significant features of body-weight profile changes in HIV infected patients in the IeDEA Collaboration 2020 , 15, e0220165		
196	PS-SiZer map to investigate significant features of body-weight profile changes in HIV infected patients in the IeDEA Collaboration 2020 , 15, e0220165		
195	PS-SiZer map to investigate significant features of body-weight profile changes in HIV infected patients in the IeDEA Collaboration 2020 , 15, e0220165		
194	Simplified clinical algorithm for identifying patients eligible for same-day HIV treatment initiation (SLATE): Results from an individually randomized trial in South Africa and Kenya. <i>PLoS Medicine</i> , 2019 , 16, e1002912	11.6	16
193	Clinical predictor score to identify patients at risk of poor viral load suppression at six months on antiretroviral therapy: results from a prospective cohort study in Johannesburg, South Africa. <i>Clinical Epidemiology</i> , 2019 , 11, 359-373	5.9	5
192	Applying the E Value to Assess the Robustness of Epidemiologic Fields of Inquiry to Unmeasured Confounding. <i>American Journal of Epidemiology</i> , 2019 , 188, 1174-1180	3.8	15
191	Prevalence and predictors of postpartum depression by HIV status and timing of HIV diagnosis in Gauteng, South Africa. <i>PLoS ONE</i> , 2019 , 14, e0214849	3.7	11
190	Awareness of and potential for dependent error in the observational epidemiologic literature: A review. <i>Annals of Epidemiology</i> , 2019 , 36, 15-19.e2	6.4	5

189	Adherence clubs and decentralized medication delivery to support patient retention and sustained viral suppression in care: Results from a cluster-randomized evaluation of differentiated ART delivery models in South Africa. <i>PLoS Medicine</i> , 2019 , 16, e1002874	11.6	44
188	Risk of ischemic placental disease is increased following in vitro fertilization with oocyte donation: a retrospective cohort study. <i>Journal of Assisted Reproduction and Genetics</i> , 2019 , 36, 1917-1926	3.4	9
187	Alcohol Consumption in Later Life and Mortality in the United States: Results from 9 Waves of the Health and Retirement Study. <i>Alcoholism: Clinical and Experimental Research</i> , 2019 , 43, 1734-1746	3.7	18
186	Who is seeking antiretroviral treatment for HIV now? Characteristics of patients presenting in Kenya and South Africa in 2017-2018. <i>Journal of the International AIDS Society</i> , 2019 , 22, e25358	5.4	6
185	Differentiated HIV care in South Africa: the effect of fast-track treatment initiation counselling on ART initiation and viral suppression as partial results of an impact evaluation on the impact of a package of services to improve HIV treatment adherence. <i>Journal of the International AIDS Society</i> ,	5.4	9
184	2019 , 22, e25409 Global Health Research Mentoring Competencies for Individuals and Institutions in Low- and Middle-Income Countries. <i>American Journal of Tropical Medicine and Hygiene</i> , 2019 , 100, 15-19	3.2	17
183	Extending Visit Intervals for Clinically Stable Patients on Antiretroviral Therapy: Multicohort Analysis of HIV Programs in Southern Africa. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2019 , 81, 439-447	3.1	2
182	Growth curve modelling to determine distinct BMI trajectory groups in HIV-positive adults on antiretroviral therapy in South Africa. <i>Aids</i> , 2019 , 33, 2049-2059	3.5	4
181	Adolescent HIV treatment in South Africa® national HIV programme: a retrospective cohort study. <i>Lancet HIV,the</i> , 2019 , 6, e760-e768	7.8	27
180	A Meta-analysis Assessing Diarrhea and Pneumonia in HIV-Exposed Uninfected Compared With HIV-Unexposed Uninfected Infants and Children. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2019 , 82, 1-8	3.1	15
179	Third-Line Antiretroviral Therapy Program in the South African Public Sector: Cohort Description and Virological Outcomes. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2019 , 80, 73-78	3.1	23
178	Comparison of 3 Days Amoxicillin Versus 5 Days Co-Trimoxazole for Treatment of Fast-breathing Pneumonia by Community Health Workers in Children Aged 2-59 Months in Pakistan: A Cluster-randomized Trial. <i>Clinical Infectious Diseases</i> , 2019 , 69, 397-404	11.6	5
177	The right combination - treatment outcomes among HIV-positive patients initiating first-line fixed-dose antiretroviral therapy in a public sector HIV clinic in Johannesburg, South Africa. <i>Clinical Epidemiology</i> , 2018 , 10, 17-29	5.9	10
176	Retention and mortality on antiretroviral therapy in sub-Saharan Africa: collaborative analyses of HIV treatment programmes. <i>Journal of the International AIDS Society</i> , 2018 , 21, e25084	5.4	55
175	Social and behavioral factors associated with failing second-line ART - results from a cohort study at the Themba Lethu Clinic, Johannesburg, South Africa. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2018 , 30, 863-870	2.2	5
174	Regimen durability in HIV-infected children and adolescents initiating first-line antiretroviral therapy in a large public sector HIV cohort in South Africa. <i>Tropical Medicine and International Health</i> , 2018 , 23, 650-660	2.3	2
173	Seasonal variations in tuberculosis diagnosis among HIV-positive individuals in Southern Africa: analysis of cohort studies at antiretroviral treatment programmes. <i>BMJ Open</i> , 2018 , 8, e017405	3	4
172	Assessing the impact of the National Department of Health® National Adherence Guidelines for Chronic Diseases in South Africa using routinely collected data: a cluster-randomised evaluation. BM I Open 2018 8 e019680	3	12

171	Predictors of switch to and early outcomes on third-line antiretroviral therapy at a large public-sector clinic in Johannesburg, South Africa. <i>AIDS Research and Therapy</i> , 2018 , 15, 10	3	12
170	The Impact of Joint Misclassification of Exposures and Outcomes on the Results of Epidemiologic Research. <i>Current Epidemiology Reports</i> , 2018 , 5, 166-174	2.9	7
169	The WelTel Trial in context and the importance of null findings. Lancet Public Health, The, 2018, 3, e107	- <u>e</u> 21204β	3
168	Adverse Drug Reactions Among Patients Initiating Second-Line Antiretroviral Therapy in South Africa. <i>Drug Safety</i> , 2018 , 41, 1343-1353	5.1	12
167	Stratified Probabilistic Bias Analysis for Body Mass Index-related Exposure Misclassification in Postmenopausal Women. <i>Epidemiology</i> , 2018 , 29, 604-613	3.1	12
166	Outcomes of Patients Lost to Follow-up in African Antiretroviral Therapy Programs: Individual Patient Data Meta-analysis. <i>Clinical Infectious Diseases</i> , 2018 , 67, 1643-1652	11.6	47
165	Failure to initiate HIV treatment in patients with high CD4 counts: evidence from demographic surveillance in rural South Africa. <i>Tropical Medicine and International Health</i> , 2018 , 23, 206-220	2.3	14
164	Acceptability and feasibility of a financial incentive intervention to improve retention in HIV care among pregnant women in Johannesburg, South Africa. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2018 , 30, 453-460	2.2	12
163	Quantitative Bias Analysis for Collaborative Science. <i>Epidemiology</i> , 2018 , 29, 627-630	3.1	7
162	Low prevalence of depressive symptoms among stable patients on antiretroviral therapy in Johannesburg, South Africa. <i>PLoS ONE</i> , 2018 , 13, e0203797	3.7	4
161	Prevalence, incidence, predictors, treatment, and control of hypertension among HIV-positive adults on antiretroviral treatment in public sector treatment programs in South Africa. <i>PLoS ONE</i> , 2018 , 13, e0204020	3.7	22
160	Effectiveness of interventions for unstable patients on antiretroviral therapy in South Africa: results of a cluster-randomised evaluation. <i>Tropical Medicine and International Health</i> , 2018 , 23, 1314-13	3 2 5³	16
159	IVF success corrected for drop-out: use of inverse probability weighting. <i>Human Reproduction</i> , 2018 , 33, 2295-2301	5.7	14
158	"My future is brightIlwonR die with the cause of AIDS": ten-year patient ART outcomes and experiences in South Africa. <i>Journal of the International AIDS Society</i> , 2018 , 21, e25184	5.4	10
157	Routine data underestimates the incidence of first-line antiretroviral drug discontinuations due to adverse drug reactions: Observational study in two South African cohorts. <i>PLoS ONE</i> , 2018 , 13, e020353	3∂ ^{.7}	2
156	"I will leave the baby with my mother": Long-distance travel and follow-up care among HIV-positive pregnant and postpartum women in South Africa. <i>Journal of the International AIDS Society</i> , 2018 , 21 Suppl 4, e25121	5.4	18
155	Implementation of Option B and a fixed-dose combination antiretroviral regimen for prevention of mother-to-child transmission of HIV in South Africa: A model of uptake and adherence to care. <i>PLoS ONE</i> , 2018 , 13, e0201955	3.7	2
154	Estimating retention in HIV care accounting for patient transfers: A national laboratory cohort study in South Africa. <i>PLoS Medicine</i> , 2018 , 15, e1002589	11.6	45

153	Medication Side Effects and Retention in HIV Treatment: A Regression Discontinuity Study of Tenofovir Implementation in South Africa and Zambia. <i>American Journal of Epidemiology</i> , 2018 , 187, 199	9ð: ⁸ 200	1 ⁶
152	Understanding Predictors of Early Antenatal Care Initiation in Relationship to Timing of HIV Diagnosis in South Africa. <i>AIDS Patient Care and STDs</i> , 2018 , 32, 251-256	5.8	4
151	Persistent High Burden of Advanced HIV Disease Among Patients Seeking Care in South Africaß National HIV Program: Data From a Nationwide Laboratory Cohort. <i>Clinical Infectious Diseases</i> , 2018 , 66, S111-S117	11.6	73
150	Why do people living with HIV not initiate treatment? A systematic review of qualitative evidence from low- and middle-income countries. <i>Social Science and Medicine</i> , 2018 , 213, 72-84	5.1	56
149	Do HIV treatment eligibility expansions crowd out the sickest? Evidence from rural South Africa. <i>Tropical Medicine and International Health</i> , 2018 , 23, 968-979	2.3	6
148	A Clinical Validation of Self-Reported Periodontitis Among Participants in the Black Womenß Health Study. <i>Journal of Periodontology</i> , 2017 , 88, 582-592	4.6	26
147	Tenofovir stock shortages have limited impact on clinic- and patient-level HIV treatment outcomes in public sector clinics in South Africa. <i>Tropical Medicine and International Health</i> , 2017 , 22, 241-251	2.3	7
146	On the Need for Quantitative Bias Analysis in the Peer-Review Process. <i>American Journal of Epidemiology</i> , 2017 , 185, 865-868	3.8	21
145	Timing of pregnancy, postpartum risk of virologic failure and loss to follow-up among HIV-positive women. <i>Aids</i> , 2017 , 31, 1593-1602	3.5	18
144	Predicting the Need for Third-Line Antiretroviral Therapy by Identifying Patients at High Risk for Failing Second-Line Antiretroviral Therapy in South Africa. <i>AIDS Patient Care and STDs</i> , 2017 , 31, 205-21	2 ^{5.8}	24
143	Initiating antiretroviral therapy for HIV at a patient® first clinic visit: a cost-effectiveness analysis of the rapid initiation of treatment randomized controlled trial. <i>Aids</i> , 2017 , 31, 1611-1619	3.5	20
142	Simplified clinical algorithm for identifying patients eligible for immediate initiation of antiretroviral therapy for HIV (SLATE): protocol for a randomised evaluation. <i>BMJ Open</i> , 2017 , 7, e0163	4 0	12
141	Mobility and Clinic Switching Among Postpartum Women Considered Lost to HIV Care in South Africa. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2017 , 74, 383-389	3.1	55
140	Comparison of Kaposi Sarcoma Risk in Human Immunodeficiency Virus-Positive Adults Across 5 Continents: A Multiregional Multicohort Study. <i>Clinical Infectious Diseases</i> , 2017 , 65, 1316-1326	11.6	23
139	Citizenship status and engagement in HIV care: an observational cohort study to assess the association between reporting a national ID number and retention in public-sector HIV care in Johannesburg, South Africa. <i>BMJ Open</i> , 2017 , 7, e013908	3	3
138	Changes in estimated glomerular filtration rate over time in South African HIV-1-infected patients receiving tenofovir: a retrospective cohort study. <i>Journal of the International AIDS Society</i> , 2017 , 20, 213	3 7 7	26
137	Twelve-year mortality in adults initiating antiretroviral therapy in South Africa. <i>Journal of the International AIDS Society</i> , 2017 , 20, 21902	5.4	40
136	A new cascade of HIV care for the era of "treat all". <i>PLoS Medicine</i> , 2017 , 14, e1002268	11.6	48

Treatment eligibility and retention in clinical HIV care: A regression discontinuity study in South Africa. <i>PLoS Medicine</i> , 2017 , 14, e1002463	11.6	43
Varying intervals of antiretroviral medication dispensing to improve outcomes for HIV patients (The INTERVAL Study): study protocol for a randomized controlled trial. <i>Trials</i> , 2017 , 18, 476	2.8	11
Recurrent Yeast Infections and Vulvodynia: Can We Believe Associations Based on Self-Reported Data?. <i>Journal of Womenls Health</i> , 2017 , 26, 1069-1076	3	18
Has the phasing out of stavudine in accordance with changes in WHO guidelines led to a decrease in single-drug substitutions in first-line antiretroviral therapy for HIV in sub-Saharan Africa?. <i>Aids</i> , 2017 , 31, 147-157	3.5	12
Imputing HIV treatment start dates from routine laboratory data in South Africa: a validation study. <i>BMC Health Services Research</i> , 2017 , 17, 41	2.9	16
Health facility and skilled birth deliveries among poor women with Jamkesmas health insurance in Indonesia: a mixed-methods study. <i>BMC Health Services Research</i> , 2017 , 17, 105	2.9	18
Treatment outcomes of over 1000 patients on second-line, protease inhibitor-based antiretroviral therapy from four public-sector HIV treatment facilities across Johannesburg, South Africa. <i>Tropical Medicine and International Health</i> , 2017 , 22, 221-231	2.3	8
HIV viral load as an independent risk factor for tuberculosis in South Africa: collaborative analysis of cohort studies. <i>Journal of the International AIDS Society</i> , 2017 , 20, 21327	5.4	23
Cohort profile: the Right to Care Clinical HIV Cohort, South Africa. <i>BMJ Open</i> , 2017 , 7, e015620	3	15
Effect of eliminating CD4-count thresholds on HIV treatment initiation in South Africa: An empirical modeling study. <i>PLoS ONE</i> , 2017 , 12, e0178249	3.7	18
Changing the South African national antiretroviral therapy guidelines: The role of cost modelling. <i>PLoS ONE</i> , 2017 , 12, e0186557	3.7	34
Are we shifting attrition downstream in the HIV cascade?. <i>Lancet HIV,the</i> , 2016 , 3, e554-e555	7.8	14
Can Short-Term Use of Electronic Patient Adherence Monitoring Devices Improve Adherence in Patients Failing Second-Line Antiretroviral Therapy? Evidence from a Pilot Study in Johannesburg, South Africa. <i>AIDS and Behavior</i> , 2016 , 20, 2717-2728	4.3	9
CD4 count at antiretroviral therapy initiation and the risk of loss to follow-up: results from a multicentre cohort study. <i>Journal of Epidemiology and Community Health</i> , 2016 , 70, 549-55	5.1	28
The High Cost of HIV-Positive Inpatient Care at an Urban Hospital in Johannesburg, South Africa. <i>PLoS ONE</i> , 2016 , 11, e0148546	3.7	16
Treatment Outcomes and Costs of Providing Antiretroviral Therapy at a Primary Health Clinic versus a Hospital-Based HIV Clinic in South Africa. <i>PLoS ONE</i> , 2016 , 11, e0168118	3.7	9
Interventions to improve the rate or timing of initiation of antiretroviral therapy for HIV in sub-Saharan Africa: meta-analyses of effectiveness. <i>Journal of the International AIDS Society</i> , 2016 , 19, 20888	5.4	49
Developing a predictive risk model for first-line antiretroviral therapy failure in South Africa. Journal of the International AIDS Society, 2016 , 19, 20987	5.4	12
	Africa. PLoS Medicine, 2017, 14, e1002463 Varying intervals of antiretroviral medication dispensing to improve outcomes for HIV patients (The INTERVAL Study): study protocol for a randomized controlled trial. Trials, 2017, 18, 476 Recurrent Yeast Infections and Vulvodynia: Can We Believe Associations Based on Self-Reported Data?. Journal of Women's Health, 2017, 26, 1069-1076 Has the phasing out of stavudine in accordance with changes in WHO guidelines led to a decrease in single-drug substitutions in first-line antiretroviral therapy for HIV in sub-Saharan Africa?. Aids, 2017, 31, 147-157 Imputing HIV treatment start dates from routine laboratory data in South Africa: a validation study. BMC Health Services Research, 2017, 17, 41 Health facility and skilled birth deliveries among poor women with Jamkesmas health insurance in Indonesia: a mixed-methods study. BMC Health Services Research, 2017, 17, 105 Treatment outcomes of over 1000 patients on second-line, protease Inhibitor-based antiretroviral therapy from four public-sector HIV treatment Facilities across Johannesburg, South Africa. Tropical Medicine and International Health, 2017, 22, 221-231 HIV viral load as an independent risk factor for tuberculosis in South Africa: collaborative analysis of cohort studies. Journal of the International AIDS Society, 2017, 20, 21327 Cohort profile: the Right to Care Clinical HIV Cohort, South Africa. BMJ Open, 2017, 7, e015620 Effect of eliminating CD4-count thresholds on HIV treatment initiation in South Africa: An empirical modeling study. PLoS ONE, 2017, 12, e0178249 Changing the South African national antiretroviral therapy guidelines: The role of cost modelling. PLoS ONE, 2017, 12, e0186557 Are we shifting attrition downstream in the HIV cascade?. Lancet HIV, the, 2016, 3, e554-e555 Can Short-Term Use of Electronic Patient Adherence Monitoring Devices Improve Adherence in Patients Falling Second-Line Antiretroviral therapy? Evidence from a Pilot Study in Johannesburg, South Africa. AIDS and Behavior, 2	Africa. PLoS Medicine, 2017, 14, e1002463 Varying intervals of antiretroviral medication dispensing to improve outcomes for HIV patients (The INTERVAL Study): study protocol for a randomized controlled trial. Triols, 2017, 18, 476 Recurrent Yeast Infections and Vulvodynia: Can We Believe Associations Based on Self-Reported Data?. Journal of Women's Health, 2017, 26, 1069-1076 3. Has the phasing out of stavudine in accordance with changes in WHO guidelines led to a decrease in single-drug substitutions in first-line antiretroviral therapy for HIV in sub-Saharan Africa?. Aids, 2017, 31, 147-157 Imputing HIV treatment start dates from routine laboratory data in South Africa: a validation study, BMC Health Services Research, 2017, 17, 41 Health facility and skilled birth deliveries among poor women with Jamkesmas health insurance in Indonesia: a mixed methods study. BMC Health Services Research, 2017, 17, 105 Treatment outcomes of over 1000 patients on second-line, protease inhibitor-based antiretroviral therapy from four public-sector HIV treatment facilities across Johannesburg, South Africa. Tropical Medicine and International Health, 2017, 22, 221-231 HIV viral load as an independent risk factor for tuberculosis in South Africa: collaborative analysis of cohort studies. Journal of the International AIDS Society, 2017, 20, 21327 Cohort profile: the Right to Care Clinical HIV Cohort, South Africa. BMJ Open, 2017, 7, e015620 3. Effect of eliminating CD4-count thresholds on HIV treatment initiation in South Africa: An empirical modeling study. PLoS ONE, 2017, 12, e0178249 Changing the South African national antiretroviral therapy guidelines: The role of cost modelling. PLoS ONE, 2017, 12, e0186557 Are we shifting attrition downstream in the HIV cascade?. Lancet HIV, the, 2016, 3, e554-e555 7.8 Can Short-Term Use of Electronic Patient Adherence Monitoring Devices Improve Adherence in Patients Falling Second-Line Antiretroviral Therapy? Evidence From a Pilot Study in Johannesburg, South Africa. PLoS ONE,

117	Changes in second-line regimen durability and continuity of care in relation to national ART guideline changes in South Africa. <i>Journal of the International AIDS Society</i> , 2016 , 19, 20675	5.4	5
116	Marginal Structural Models to Assess Delays in Second-Line HIV Treatment Initiation in South Africa. <i>PLoS ONE</i> , 2016 , 11, e0161469	3.7	28
115	Life expectancy trends in adults on antiretroviral treatment in South Africa. Aids, 2016, 30, 2545-2550	3.5	11
114	Insights into Adherence among a Cohort of Adolescents Aged 12-20 Years in South Africa: Reported Barriers to Antiretroviral Treatment. <i>AIDS Research and Treatment</i> , 2016 , 2016, 4161738	2.3	14
113	Initiating Antiretroviral Therapy for HIV at a Patient® First Clinic Visit: The RapIT Randomized Controlled Trial. <i>PLoS Medicine</i> , 2016 , 13, e1002015	11.6	173
112	Accelerating the Uptake and Timing of Antiretroviral Therapy Initiation in Sub-Saharan Africa: An Operations Research Agenda. <i>PLoS Medicine</i> , 2016 , 13, e1002106	11.6	27
111	Quantitative Bias Analysis in Regulatory Settings. American Journal of Public Health, 2016, 106, 1227-30	5.1	19
110	Intensive adherence counselling for HIV-infected individuals failing second-line antiretroviral therapy in Johannesburg, South Africa. <i>Tropical Medicine and International Health</i> , 2016 , 21, 1131-7	2.3	32
109	Stressful events during pregnancy and postpartum depressive symptoms. <i>Journal of Womenls Health</i> , 2015 , 24, 384-93	3	44
108	Antibiotic trials for community-acquired pneumonia. Lancet Respiratory Medicine, the, 2015, 3, e4-5	35.1	3
107	Systematic review of retention of pediatric patients on HIV treatment in low and middle-income countries 2008-2013. <i>Aids</i> , 2015 , 29, 493-502	3.5	53
106	Linkage to care following a home-based HIV counselling and testing intervention in rural South Africa. <i>Journal of the International AIDS Society</i> , 2015 , 18, 19843	5.4	58
105	A comparison of death recording by health centres and civil registration in South Africans receiving antiretroviral treatment. <i>Journal of the International AIDS Society</i> , 2015 , 18, 20628	5.4	27
104	Maternal Recall Error in Retrospectively Reported Time-to-Pregnancy: an Assessment and Bias Analysis. <i>Paediatric and Perinatal Epidemiology</i> , 2015 , 29, 576-88	2.7	14
103	Retention of Adult Patients on Antiretroviral Therapy in Low- and Middle-Income Countries: Systematic Review and Meta-analysis 2008-2013. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2015 , 69, 98-108	3.1	206
102	Implementation and Operational Research: Risk Charts to Guide Targeted HIV-1 Viral Load Monitoring of ART: Development and Validation in Patients From Resource-Limited Settings. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2015 , 70, e110-9	3.1	7
101	Brief Report: Does Most Mortality in Patients on ART Occur in Care or After Lost to Follow-Up? Evidence From the Themba Lethu Clinic, South Africa. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2015 , 70, 323-8	3.1	6
100	Kaposiß Sarcoma in HIV-infected patients in South Africa: Multicohort study in the antiretroviral therapy era. <i>International Journal of Cancer</i> , 2014 , 135, 2644-52	7.5	40

99	Incidence and predictors of herpes zoster among antiretroviral therapy-naWe patients initiating HIV treatment in Johannesburg, South Africa. <i>International Journal of Infectious Diseases</i> , 2014 , 23, 56-62	10.5	16
98	CD4 criteria improves the sensitivity of a clinical algorithm developed to identify viral failure in HIV-positive patients on antiretroviral therapy. <i>Journal of the International AIDS Society</i> , 2014 , 17, 1913	9 ^{5.4}	10
97	Poor CD4 recovery and risk of subsequent progression to AIDS or death despite viral suppression in a South African cohort. <i>Journal of the International AIDS Society</i> , 2014 , 17, 18651	5.4	31
96	The relation between efavirenz versus nevirapine and virologic failure in Johannesburg, South Africa. <i>Journal of the International AIDS Society</i> , 2014 , 17, 19065	5.4	12
95	Immunodeficiency at the start of combination antiretroviral therapy in low-, middle-, and high-income countries. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2014 , 65, e8-16	3.1	122
94	Good practices for quantitative bias analysis. <i>International Journal of Epidemiology</i> , 2014 , 43, 1969-85	7.8	272
93	Mortality in patients with HIV-1 infection starting antiretroviral therapy in South Africa, Europe, or North America: a collaborative analysis of prospective studies. <i>PLoS Medicine</i> , 2014 , 11, e1001718	11.6	89
92	Impact of systematic HIV testing on case finding and retention in care at a primary care clinic in South Africa. <i>Tropical Medicine and International Health</i> , 2014 , 19, 1411-9	2.3	7
91	Impact of choice of NRTI in first-line antiretroviral therapy: a cohort analysis of stavudine vs. tenofovir. <i>Tropical Medicine and International Health</i> , 2014 , 19, 490-8	2.3	9
90	Mortality among adults transferred and lost to follow-up from antiretroviral therapy programmes in South Africa: a multicenter cohort study. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2014 , 67, e67-75	3.1	43
89	Attrition through multiple stages of pre-treatment and ART HIV care in South Africa. <i>PLoS ONE</i> , 2014 , 9, e110252	3.7	50
88	Low rates of treatment failure in children aged 2-59 months treated for severe pneumonia: a multisite pooled analysis. <i>Clinical Infectious Diseases</i> , 2013 , 56, 978-87	11.6	11
87	Tuberculosis and the risk of opportunistic infections and cancers in HIV-infected patients starting ART in Southern Africa. <i>Tropical Medicine and International Health</i> , 2013 , 18, 194-8	2.3	17
86	Increases in regimen durability associated with the introduction of tenofovir at a large public-sector clinic in Johannesburg, South Africa. <i>Journal of the International AIDS Society</i> , 2013 , 16, 18794	5.4	16
85	Loss to follow-up before and after delivery among women testing HIV positive during pregnancy in Johannesburg, South Africa. <i>Tropical Medicine and International Health</i> , 2013 , 18, 451-60	2.3	82
84	Kaposi sarcoma-associated herpes virus and response to antiretroviral therapy: a prospective study of HIV-infected adults. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2013 , 63, 442-8	3.1	10
83	Antiretroviral therapy, labor productivity, and sex: a longitudinal cohort study of tea pluckers in Kenya. <i>Aids</i> , 2013 , 27, 115-23	3.5	13
82	Patient retention from HIV diagnosis through one year on antiretroviral therapy at a primary health care clinic in Johannesburg, South Africa. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2013 , 62, e39-46	3.1	76

81	Rates and cost of hospitalization before and after initiation of antiretroviral therapy in urban and rural settings in South Africa. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2013 , 62, 322-32	28 ^{3.1}	27
80	CD4 count slope and mortality in HIV-infected patients on antiretroviral therapy: multicohort analysis from South Africa. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2013 , 63, 34-41	3.1	21
79	Life expectancies of South African adults starting antiretroviral treatment: collaborative analysis of cohort studies. <i>PLoS Medicine</i> , 2013 , 10, e1001418	11.6	277
78	Effect of home based HIV counselling and testing intervention in rural South Africa: cluster randomised trial. <i>BMJ, The</i> , 2013 , 346, f3481	5.9	62
77	Cohort profile: the Themba Lethu Clinical Cohort, Johannesburg, South Africa. <i>International Journal of Epidemiology</i> , 2013 , 42, 430-9	7.8	71
76	The interplay between CD4 cell count, viral load suppression and duration of antiretroviral therapy on mortality in a resource-limited setting. <i>Tropical Medicine and International Health</i> , 2013 , 18, 619-31	2.3	22
75	Same-Day CD4 Testing to Improve Uptake of HIV Care and Treatment in South Africa: Point-of-Care Is Not Enough. <i>AIDS Research and Treatment</i> , 2013 , 2013, 941493	2.3	28
74	The early effects of stavudine compared with tenofovir on adipocyte gene expression, mitochondrial DNA copy number and metabolic parameters in South African HIV-infected patients: a randomized trial. <i>HIV Medicine</i> , 2013 , 14, 217-25	2.7	7
73	Gender differences in mortality and CD4 count response among virally suppressed HIV-positive patients. <i>Journal of Womenls Health</i> , 2013 , 22, 113-20	3	67
72	Cough aerosols of Mycobacterium tuberculosis predict new infection: a household contact study. American Journal of Respiratory and Critical Care Medicine, 2013 , 187, 1007-15	10.2	95
72 71		3.5	95
	American Journal of Respiratory and Critical Care Medicine, 2013 , 187, 1007-15 Initiating antiretroviral therapy when presenting with higher CD4 cell counts results in reduced loss		
71	American Journal of Respiratory and Critical Care Medicine, 2013, 187, 1007-15 Initiating antiretroviral therapy when presenting with higher CD4 cell counts results in reduced loss to follow-up in a resource-limited setting. Aids, 2013, 27, 645-50 The feasibility of using screening criteria to reduce clinic visits for stable patients on antiretroviral	3.5	44
71 70	American Journal of Respiratory and Critical Care Medicine, 2013, 187, 1007-15 Initiating antiretroviral therapy when presenting with higher CD4 cell counts results in reduced loss to follow-up in a resource-limited setting. Aids, 2013, 27, 645-50 The feasibility of using screening criteria to reduce clinic visits for stable patients on antiretroviral therapy in South Africa. Journal of Acquired Immune Deficiency Syndromes (1999), 2013, 62, e82-6 Treatment response and mortality among patients starting antiretroviral therapy with and without	3.5	44
7 ¹ 7 ⁰ 69	American Journal of Respiratory and Critical Care Medicine, 2013, 187, 1007-15 Initiating antiretroviral therapy when presenting with higher CD4 cell counts results in reduced loss to follow-up in a resource-limited setting. Aids, 2013, 27, 645-50 The feasibility of using screening criteria to reduce clinic visits for stable patients on antiretroviral therapy in South Africa. Journal of Acquired Immune Deficiency Syndromes (1999), 2013, 62, e82-6 Treatment response and mortality among patients starting antiretroviral therapy with and without Kaposi sarcoma: a cohort study. PLoS ONE, 2013, 8, e64392 A novel approach to accounting for loss to follow-up when estimating the relationship between	3.5 3.1 3.7	44331
71 70 69 68	American Journal of Respiratory and Critical Care Medicine, 2013, 187, 1007-15 Initiating antiretroviral therapy when presenting with higher CD4 cell counts results in reduced loss to follow-up in a resource-limited setting. Aids, 2013, 27, 645-50 The feasibility of using screening criteria to reduce clinic visits for stable patients on antiretroviral therapy in South Africa. Journal of Acquired Immune Deficiency Syndromes (1999), 2013, 62, e82-6 Treatment response and mortality among patients starting antiretroviral therapy with and without Kaposi sarcoma: a cohort study. PLoS ONE, 2013, 8, e64392 A novel approach to accounting for loss to follow-up when estimating the relationship between CD4 Count at ART initiation and mortality. PLoS ONE, 2013, 8, e69300 The impact of choice of NNRTI on short-term treatment outcomes among HIV-infected patients	3.5 3.1 3.7	443314
71 70 69 68 67	American Journal of Respiratory and Critical Care Medicine, 2013, 187, 1007-15 Initiating antiretroviral therapy when presenting with higher CD4 cell counts results in reduced loss to follow-up in a resource-limited setting. Aids, 2013, 27, 645-50 The feasibility of using screening criteria to reduce clinic visits for stable patients on antiretroviral therapy in South Africa. Journal of Acquired Immune Deficiency Syndromes (1999), 2013, 62, e82-6 Treatment response and mortality among patients starting antiretroviral therapy with and without Kaposi sarcoma: a cohort study. PLoS ONE, 2013, 8, e64392 A novel approach to accounting for loss to follow-up when estimating the relationship between CD4 Count at ART initiation and mortality. PLoS ONE, 2013, 8, e69300 The impact of choice of NNRTI on short-term treatment outcomes among HIV-infected patients prescribed tenofovir and lamivudine in Johannesburg, South Africa. PLoS ONE, 2013, 8, e71719 Effectiveness and safety of 30 mg versus 40 mg stavudine regimens: a cohort study among HIV-infected adults initiating HAART in South Africa. Journal of the International AIDS Society, 2012,	3.5 3.1 3.7 3.7	4433148

(2011-2012)

63	Low rates of nucleoside reverse transcriptase inhibitor resistance in a well-monitored cohort in South Africa on antiretroviral therapy. <i>Antiviral Therapy</i> , 2012 , 17, 313-20	1.6	20
62	Poorer ART outcomes with increasing age at a large public sector HIV clinic in Johannesburg, South Africa. <i>Journal of the International Association of Providers of AIDS Care</i> , 2012 , 11, 57-65		36
61	Response to Lawn et al <i>Aids</i> , 2012 , 26, 1728-1729	3.5	
60	Rapid point-of-care CD4 testing at mobile HIV testing sites to increase linkage to care: an evaluation of a pilot program in South Africa. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2012 , 61, e13-7	3.1	87
59	Rates and predictors of failure of first-line antiretroviral therapy and switch to second-line ART in South Africa. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2012 , 60, 428-37	3.1	107
58	Prevalent tuberculosis and mortality among HAART initiators. <i>Aids</i> , 2012 , 26, 770-3	3.5	7
57	Treatment outcomes after 7 years of public-sector HIV treatment. Aids, 2012, 26, 1823-8	3.5	33
56	Frequency and trajectory of abnormalities in respiratory rate, temperature and oxygen saturation in severe pneumonia in children. <i>Pediatric Infectious Disease Journal</i> , 2012 , 31, 863-5	3.4	5
55	The net cost of incorporating resistance testing into HIV/AIDS treatment in South Africa: a Markov model with primary data. <i>Journal of the International AIDS Society</i> , 2011 , 14, 24	5.4	22
54	Community case management of severe pneumonia with oral amoxicillin in children aged 2-59 months in Haripur district, Pakistan: a cluster randomised trial. <i>Lancet, The</i> , 2011 , 378, 1796-803	40	69
53	Relationship between renal dysfunction, nephrotoxicity and death among HIV adults on tenofovir. <i>Aids</i> , 2011 , 25, 1603-9	3.5	67
52	Outcomes of stable HIV-positive patients down-referred from a doctor-managed antiretroviral therapy clinic to a nurse-managed primary health clinic for monitoring and treatment. <i>Aids</i> , 2011 , 25, 2027-36	3.5	68
51	Tuberculosis in HIV programmes in lower-income countries: practices and risk factors. <i>International Journal of Tuberculosis and Lung Disease</i> , 2011 , 15, 620-7	2.1	30
50	Response to: defaulting from antiretroviral treatment programmes in sub-Saharan Africa: a problem of definition. <i>Tropical Medicine and International Health</i> , 2011 , 16, 392-392	2.3	
49	Outpatient treatment of children with severe pneumonia with oral amoxicillin in four countries: the MASS study. <i>Tropical Medicine and International Health</i> , 2011 , 16, 995-1006	2.3	49
48	Prevalence and predictors of kaposi sarcoma herpes virus seropositivity: a cross-sectional analysis of HIV-infected adults initiating ART in Johannesburg, South Africa. <i>Infectious Agents and Cancer</i> , 2011 , 6, 22	3.5	17
47	Effect of training traditional birth attendants on neonatal mortality (Lufwanyama Neonatal Survival Project): randomised controlled study. <i>BMJ, The</i> , 2011 , 342, d346	5.9	107
46	Outcomes of antiretroviral treatment in programmes with and without routine viral load monitoring in Southern Africa. <i>Aids</i> , 2011 , 25, 1761-9	3.5	88

45	Correcting mortality for loss to follow-up: a nomogram applied to antiretroviral treatment programmes in sub-Saharan Africa. <i>PLoS Medicine</i> , 2011 , 8, e1000390	11.6	125
44	Retention in HIV care between testing and treatment in sub-Saharan Africa: a systematic review. <i>PLoS Medicine</i> , 2011 , 8, e1001056	11.6	576
43	Treatment outcomes and cost-effectiveness of shifting management of stable ART patients to nurses in South Africa: an observational cohort. <i>PLoS Medicine</i> , 2011 , 8, e1001055	11.6	92
42	Using vital registration data to update mortality among patients lost to follow-up from ART programmes: evidence from the Themba Lethu Clinic, South Africa. <i>Tropical Medicine and International Health</i> , 2010 , 15, 405-13	2.3	94
41	Patient retention in antiretroviral therapy programs up to three years on treatment in sub-Saharan Africa, 2007-2009: systematic review. <i>Tropical Medicine and International Health</i> , 2010 , 15 Suppl 1, 1-15	2.3	407
40	Early loss to follow up after enrolment in pre-ART care at a large public clinic in Johannesburg, South Africa. <i>Tropical Medicine and International Health</i> , 2010 , 15 Suppl 1, 43-7	2.3	85
39	Economic outcomes of patients receiving antiretroviral therapy for HIV/AIDS in South Africa are sustained through three years on treatment. <i>PLoS ONE</i> , 2010 , 5, e12731	3.7	31
38	Re: Promoting healthy skepticism in the news: helping journalists get it right. <i>Journal of the National Cancer Institute</i> , 2010 , 102, 829-30; author reply 830	9.7	2
37	Improvements in physical wellbeing over the first two years on antiretroviral therapy in western Kenya. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2010 , 22, 137-45	2.2	16
36	Lost opportunities to complete CD4+ lymphocyte testing among patients who tested positive for HIV in South Africa. <i>Bulletin of the World Health Organization</i> , 2010 , 88, 675-80	8.2	51
35	High rates of survival, immune reconstitution, and virologic suppression on second-line antiretroviral therapy in South Africa. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2010 , 53, 500-6	3.1	65
34	Nurse versus doctor management of HIV-infected patients receiving antiretroviral therapy (CIPRA-SA): a randomised non-inferiority trial. <i>Lancet, The</i> , 2010 , 376, 33-40	40	186
33	The high cost of second-line antiretroviral therapy for HIV/AIDS in South Africa. <i>Aids</i> , 2010 , 24, 915-9	3.5	68
32	Temporal changes in programme outcomes among adult patients initiating antiretroviral therapy across South Africa, 2002-2007. <i>Aids</i> , 2010 , 24, 2263-70	3.5	181
31	Initiating patients on antiretroviral therapy at CD4 cell counts above 200 cells/microl is associated with improved treatment outcomes in South Africa. <i>Aids</i> , 2010 , 24, 2041-50	3.5	58
30	The importance of clinic attendance in the first six months on antiretroviral treatment: a retrospective analysis at a large public sector HIV clinic in South Africa. <i>Journal of the International AIDS Society</i> , 2010 , 13, 49	5.4	56
29	Barriers to initiation of antiretroviral treatment in rural and urban areas of Zambia: a cross-sectional study of cost, stigma, and perceptions about ART. <i>Journal of the International AIDS Society</i> , 2010 , 13, 8	5.4	57
28	Role of breastfeeding cessation in mediating the relationship between maternal HIV disease stage and increased child mortality among HIV-exposed uninfected children. <i>International Journal of Epidemiology</i> , 2009 , 38, 569-76	7.8	20

27	Applying Quantitative Bias Analysis to Epidemiologic Data 2009 ,		251
26	High frequency of multidrug-resistant gram-negative rods in 2 neonatal intensive care units in the Philippines. <i>Infection Control and Hospital Epidemiology</i> , 2009 , 30, 543-9	2	36
25	Creating a demand for bias analysis in epidemiological research. <i>Journal of Epidemiology and Community Health</i> , 2009 , 63, 91	5.1	12
24	Global strategies to reduce the price of antiretroviral medicines: evidence from transactional databases. <i>Bulletin of the World Health Organization</i> , 2009 , 87, 520-8	8.2	76
23	Ambulatory short-course high-dose oral amoxicillin for treatment of severe pneumonia in children: a randomised equivalency trial. <i>Lancet, The</i> , 2008 , 371, 49-56	40	127
22	Early effects of antiretroviral therapy on work performance: preliminary results from a cohort study of Kenyan agricultural workers. <i>Aids</i> , 2008 , 22, 421-5	3.5	58
21	Reduced mortality associated with breast-feeding-acquired HIV infection and breast-feeding among HIV-infected children in Zambia. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2008 , 48, 90-6	3.1	15
20	Cost and cost-effectiveness of switching from stavudine to tenofovir in first-line antiretroviral regimens in South Africa. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2008 , 48, 334-44	3.1	38
19	Under utilization of surveillance mammography among older breast cancer survivors. <i>Journal of General Internal Medicine</i> , 2008 , 23, 158-63	4	74
18	Mammography surveillance and mortality in older breast cancer survivors. <i>Journal of Clinical Oncology</i> , 2007 , 25, 3001-6	2.2	102
17	Patient retention in antiretroviral therapy programs in sub-Saharan Africa: a systematic review. <i>PLoS Medicine</i> , 2007 , 4, e298	11.6	572
16	Improved diagnostic testing and malaria treatment practices in Zambia. <i>JAMA - Journal of the American Medical Association</i> , 2007 , 297, 2227-31	27.4	188
15	Using probabilistic corrections to account for abstractor agreement in medical record reviews. <i>American Journal of Epidemiology</i> , 2007 , 165, 1454-61	3.8	22
14	The impact of AIDS on government service delivery: the case of the Zambia Wildlife Authority. <i>Aids</i> , 2007 , 21 Suppl 3, S53-9	3.5	7
13	Resolving design problems in equivalency trials. <i>Journal of Pediatrics</i> , 2006 , 149, 12-6	3.6	10
12	Reduced mortality rate associated with annual mammograms after breast cancer therapy. <i>Breast Journal</i> , 2006 , 12, 2-6	1.2	27
11	Adherence to tamoxifen over the five-year course. Breast Cancer Research and Treatment, 2006, 99, 215	-204	257
10	Failure of standard antimicrobial therapy in children aged 3-59 months with mild or asymptomatic HIV infection and severe pneumonia. <i>Bulletin of the World Health Organization</i> , 2006 , 84, 269-75	8.2	25

9	A sensitivity analysis of a randomized controlled trial of zinc in treatment of falciparum malaria in children. <i>Contemporary Clinical Trials</i> , 2005 , 26, 281-9	2.3	3	
8	A method to automate probabilistic sensitivity analyses of misclassified binary variables. <i>International Journal of Epidemiology</i> , 2005 , 34, 1370-6	7.8	197	
7	The cost of HIV/AIDS to businesses in southern Africa. <i>Aids</i> , 2004 , 18, 317-24	3.5	40	
6	The impact of HIV/AIDS on labour productivity in Kenya. <i>Tropical Medicine and International Health</i> , 2004 , 9, 318-24	2.3	105	
5	AIDS is your business. <i>Harvard Business Review</i> , 2003 , 81, 80-7, 125		31	
4	Assessing Knowledge, Attitudes, and Practices towards Causal Directed Acyclic Graphs: a qualitative research project		1	
3	Use of directed acyclic graphs (DAGs) in applied health research: review and recommendations		15	
2	Building a National HIV Cohort from Routine Laboratory Data: Probabilistic Record-Linkage with Graph	S	6	
1	The impact of delayed switch to second-line antiretroviral therapy on mortality, depending on failure time definition and CD4 count at failure		2	