Masakatsu Tsurusaki

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2344028/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Predictive factors of truncation artifacts in the arterial phase of Gd-EOB-DTPA-enhanced MRI: a nationwide multicenter study. Japanese Journal of Radiology, 2021, 39, 165-177.	1.0	5
2	Pre-Operative Imaging and Pathological Diagnosis of Localized High-Grade Pancreatic Intra-Epithelial Neoplasia without Invasive Carcinoma. Cancers, 2021, 13, 945.	1.7	14
3	Three-Dimensional Radiological Assessment of Ablative Margins in Hepatocellular Carcinoma: Pilot Study of Overlay Fused CT/MRI Imaging with Automatic Registration. Cancers, 2021, 13, 1460.	1.7	5
4	Higher Enhancement Intrahepatic Nodules on the Hepatobiliary Phase of Gd-EOB-DTPA-Enhanced MRI as a Poor Responsive Marker of Anti-PD-1/PD-L1 Monotherapy for Unresectable Hepatocellular Carcinoma. Liver Cancer, 2021, 10, 615-628.	4.2	31
5	Dual-Energy Computed Tomography of the Liver: Uses in Clinical Practices and Applications. Diagnostics, 2021, 11, 161.	1.3	16
6	Analysis of Progression Time in Pancreatic Cancer including Carcinoma In Situ Based on Magnetic Resonance Cholangiopancreatography Findings. Diagnostics, 2021, 11, 1858.	1.3	4
7	Usefulness of respiratory-gated PET acquisition during delayed F-FDG PET/CT scanning for patients with liver metastases. Asia Oceania Journal of Nuclear Medicine and Biology, 2021, 9, 12-149.	0.1	0
8	Patterns of bone metastases from head and neck squamous cell carcinoma. Auris Nasus Larynx, 2020, 47, 262-267.	0.5	12
9	Dualâ€frequency MR elastography to differentiate between inflammation and fibrosis of the liver: Comparison with histopathology. Journal of Magnetic Resonance Imaging, 2020, 51, 1053-1064.	1.9	6
10	Transcatheter Arterial Embolization Treatment for Bleeding Visceral Artery Pseudoaneurysms in Patients with Pancreatitis or following Pancreatic Surgery. Cancers, 2020, 12, 2733.	1.7	8
11	Prediction of post-hepatectomy liver failure using gadoxetic acid-enhanced magnetic resonance imaging for hepatocellular carcinoma with portal vein invasion. European Journal of Radiology, 2020, 130, 109189.	1.2	18
12	Exploratory Analysis of Lenvatinib Therapy in Patients with Unresectable Hepatocellular Carcinoma Who Have Failed Prior PDâ^'1/PD-L1 Checkpoint Blockade. Cancers, 2020, 12, 3048.	1.7	37
13	Assessment of Liver Metastases Using CT and MRI Scans in Patients with Pancreatic Ductal Adenocarcinoma: Effects of Observer Experience on Diagnostic Accuracy. Cancers, 2020, 12, 1455.	1.7	6
14	Efficiency of a computer-aided diagnosis (CAD) system with deep learning in detection of pulmonary nodules on 1-mm-thick images of computed tomography. Japanese Journal of Radiology, 2020, 38, 1052-1061.	1.0	24
15	Partial Pancreatic Parenchymal Atrophy Is a New Specific Finding to Diagnose Small Pancreatic Cancer (â‰⊉0 mm) Including Carcinoma in Situ: Comparison with Localized Benign Main Pancreatic Duct Stenosis Patients. Diagnostics, 2020, 10, 445.	1.3	24
16	The technical aspects of a feasible new technique for ipsilateral percutaneous transhepatic portal vein embolization. British Journal of Radiology, 2018, 91, 20180124.	1.0	4
17	Dualâ€energy computed tomography for nonâ€invasive staging of liver fibrosis: Accuracy of iodine density measurements from contrastâ€enhanced data. Hepatology Research, 2018, 48, 1008-1019.	1.8	45
18	Magnetic resonance elastography in the assessment of hepatic fibrosis: a study comparing transient elastography and histological data in the same patients. Abdominal Radiology, 2017, 42, 1659-1666.	1.0	17

#	Article	IF	CITATIONS
19	Comparison of gadoxetic acid-enhanced dynamic MR imaging and contrast-enhanced computed tomography for preoperative evaluation of colorectal liver metastases. Japanese Journal of Radiology, 2017, 35, 197-205.	1.0	26
20	Prospective Comparison of Gd-EOB-DTPA-Enhanced MRI with Dynamic CT for Detecting Recurrence of HCC after Radiofrequency Ablation. Liver Cancer, 2017, 6, 349-359.	4.2	29
21	Utility of Amplatzer Vascular Plug with Preoperative Common Hepatic Artery Embolization for Distal Pancreatectomy with En Bloc Celiac Axis Resection. CardioVascular and Interventional Radiology, 2017, 40, 445-449.	0.9	9
22	Present and future roles of FDG-PET/CT imaging in the management of gastrointestinal cancer: an update. Nagoya Journal of Medical Science, 2017, 79, 527-543.	0.6	43
23	Current evidence for the diagnostic value of gadoxetic acidâ€enhanced magnetic resonance imaging for liver metastasis. Hepatology Research, 2016, 46, 853-861.	1.8	34
24	Comparison of gadoxetic acid-enhanced magnetic resonance imaging and contrast-enhanced computed tomography with histopathological examinations for the identification of hepatocellular carcinoma: a multicenter phase III study. Journal of Gastroenterology, 2016, 51, 71-79.	2.3	30
25	Surgical and Locoregional Therapy of HCC: TACE. Liver Cancer, 2015, 4, 165-175.	4.2	154
26	Clinical utility of imaging for evaluation of hepatocellular carcinoma. Journal of Hepatocellular Carcinoma, 2014, 1, 101.	1.8	0
27	Hypervascular Benign and Malignant Liver Tumors That Require Differentiation from Hepatocellular Carcinoma: Key Points of Imaging Diagnosis. Liver Cancer, 2014, 3, 85-96.	4.2	70
28	Can low-dose CT with iterative reconstruction reduce both the radiation dose and the amount of iodine contrast medium in a dynamic CT study of the liver?. European Journal of Radiology, 2014, 83, 684-691.	1.2	27
29	Clinical application of 18F-fluorodeoxyglucose positron emission tomography for assessment and evaluation after therapy for malignant hepatic tumor. Journal of Gastroenterology, 2014, 49, 46-56.	2.3	32
30	Feasible and technical aspects of transcatheter arterial chemoembolization for non-resectable hepatocellular carcinoma using a 3.5-French catheter system. Abdominal Imaging, 2014, 39, 1304-1308.	2.0	2
31	Does Gadoxetic acid-enhanced 3.0T MRI in addition to 64-detector-row contrast-enhanced CT provide better diagnostic performance and change the therapeutic strategy for the preoperative evaluation of colorectal liver metastases?. European Radiology, 2014, 24, 2532-2539.	2.3	42
32	Gd-EOB-DTPA-enhanced 3.0ÂT MR imaging: quantitative and qualitative comparison of hepatocyte-phase images obtained 10Âmin and 20Âmin after injection for the detection of liver metastases from colorectal carcinoma. European Radiology, 2011, 21, 2336-2343.	2.3	48
33	Highlights on Ultrasound-Guided Subclavian Vein Access. CardioVascular and Interventional Radiology, 2011, 34, 215-216.	0.9	0
34	Detection of hepatic metastases by superparamagnetic iron oxide-enhanced MR imaging: prospective comparison between 1.5-T and 3.0-T images in the same patients. European Radiology, 2010, 20, 2265-2273.	2.3	6
35	Ultrasound-Guided Radiological Placement of Central Venous Port via the Subclavian Vein: A Retrospective Analysis of 500 Cases at a Single Institute. CardioVascular and Interventional Radiology, 2010, 33, 989-994.	0.9	43
36	Quantitative and qualitative comparison of 1.5 and 3.0 tesla MRI in patients with chronic liver diseases. Journal of Magnetic Resonance Imaging, 2009, 29, 869-879.	1.9	39

Masakatsu Tsurusaki

#	Article	IF	CITATIONS
37	Atypical hemangioma mimicking hepatocellular carcinoma with a special note on radiological and pathological findings. Japanese Journal of Radiology, 2009, 27, 156-160.	1.0	6
38	Quantitative and qualitative comparison of 3.0T and 1.5T MR imaging of the liver in patients with diffuse parenchymal liver disease. European Journal of Radiology, 2009, 72, 314-320.	1.2	39
39	3.0-T MRI evaluation of patients with chronic liver diseases: initial observations. Magnetic Resonance Imaging, 2008, 26, 650-660.	1.0	24
40	Prospective comparison of high- and low-spatial-resolution dynamic MR imaging with sensitivity encoding (SENSE) for hypervascular hepatocellular carcinoma. European Radiology, 2008, 18, 2206-2212.	2.3	3
41	Neurilemoma of the renal capsule: MR imaging and pathologic correlation. European Radiology, 2001, 11, 1834-1837.	2.3	14