

Thomas Kaufmann

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2349906/thomas-kaufmann-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

69
papers

7,622
citations

34
h-index

79
g-index

79
ext. papers

9,273
ext. citations

10.6
avg, IF

5.2
L-index

#	Paper	IF	Citations
69	Hexokinase 3 enhances myeloid cell survival via non-glycolytic functions.. <i>Cell Death and Disease</i> , 2022 , 13, 448	9.8	2
68	Granule Leakage Induces Cell-Intrinsic, Granzyme B-Mediated Apoptosis in Mast Cells. <i>Frontiers in Cell and Developmental Biology</i> , 2021 , 9, 630166	5.7	0
67	Loss of BOK Has a Minor Impact on Acetaminophen Overdose-Induced Liver Damage in Mice. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	3
66	A novel functional mast cell assay for the detection of allergies. <i>Journal of Allergy and Clinical Immunology</i> , 2021 ,	11.5	1
65	Impact of BH3-mimetics on Human and Mouse Blood Leukocytes: A Comparative Study. <i>Scientific Reports</i> , 2020 , 10, 222	4.9	3
64	The Multifaceted Roles of the BCL-2 Family Member BOK. <i>Frontiers in Cell and Developmental Biology</i> , 2020 , 8, 574338	5.7	13
63	IgA Triggers Cell Death of Neutrophils When Primed by Inflammatory Mediators. <i>Journal of Immunology</i> , 2020 , 205, 2640-2648	5.3	1
62	TNFR2 induced priming of the inflammasome leads to a RIPK1-dependent cell death in the absence of XIAP. <i>Cell Death and Disease</i> , 2019 , 10, 700	9.8	14
61	BCL-2 family protein BOK is a positive regulator of uridine metabolism in mammals. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 15469-15474	11.5	20
60	Molecular mechanisms of cell death: recommendations of the Nomenclature Committee on Cell Death 2018. <i>Cell Death and Differentiation</i> , 2018 , 25, 486-541	12.7	2160
59	IL-4 enhances survival of in vitro-differentiated mouse basophils through transcription-independent signaling downstream of PI3K. <i>Cell Death and Disease</i> , 2018 , 9, 713	9.8	4
58	Loss of BID Delays FASL-Induced Cell Death of Mouse Neutrophils and Aggravates DSS-Induced Weight Loss. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	3
57	BH3 mimetics efficiently induce apoptosis in mouse basophils and mast cells. <i>Cell Death and Differentiation</i> , 2018 , 25, 204-216	12.7	11
56	BOK promotes chemical-induced hepatocarcinogenesis in mice. <i>Cell Death and Differentiation</i> , 2018 , 25, 708-720	12.7	15
55	Negative Regulation of BOK Expression by Recruitment of TRIM28 to Regulatory Elements in Its 3' Untranslated Region. <i>Science</i> , 2018 , 9, 461-474	6.1	5
54	FcεRI cross-linking and IL-3 protect human basophils from intrinsic apoptotic stress. <i>Journal of Allergy and Clinical Immunology</i> , 2018 , 142, 1647-1650.e3	11.5	3
53	PU.1 supports TRAIL-induced cell death by inhibiting NF-κB-mediated cell survival and inducing DR5 expression. <i>Cell Death and Differentiation</i> , 2017 , 24, 866-877	12.7	16

52	BOK displays cell death-independent tumor suppressor activity in non-small-cell lung carcinoma. <i>International Journal of Cancer</i> , 2017 , 141, 2050-2061	7.5	18
51	IVIG regulates the survival of human but not mouse neutrophils. <i>Scientific Reports</i> , 2017 , 7, 1296	4.9	19
50	Balance between IL-3 and type I interferons and their interrelationship with FasL dictates lifespan and effector functions of human basophils. <i>Clinical and Experimental Allergy</i> , 2017 , 47, 71-84	4.1	6
49	Impact of inhibitor of apoptosis proteins on immune modulation and inflammation. <i>Immunology and Cell Biology</i> , 2017 , 95, 236-243	5	11
48	Interrogating the relevance of mitochondrial apoptosis for vertebrate development and postnatal tissue homeostasis. <i>Genes and Development</i> , 2016 , 30, 2133-2151	12.6	42
47	Is BOK required for apoptosis induced by endoplasmic reticulum stress?. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016 , 113, E492-3	11.5	22
46	Bok Is Not Pro-Apoptotic But Suppresses Poly ADP-Ribose Polymerase-Dependent Cell Death Pathways and Protects against Excitotoxic and Seizure-Induced Neuronal Injury. <i>Journal of Neuroscience</i> , 2016 , 36, 4564-78	6.6	35
45	In Vitro Differentiation of Mouse Granulocytes. <i>Methods in Molecular Biology</i> , 2016 , 1419, 95-107	1.4	6
44	Loss of XIAP facilitates switch to TNF-induced necroptosis in mouse neutrophils. <i>Cell Death and Disease</i> , 2016 , 7, e2422	9.8	56
43	Survival control of malignant lymphocytes by anti-apoptotic MCL-1. <i>Leukemia</i> , 2016 , 30, 2152-2159	10.7	24
42	NET formation can occur independently of RIPK3 and MLKL signaling. <i>European Journal of Immunology</i> , 2016 , 46, 178-84	6.1	72
41	The generation of neutrophils in the bone marrow is controlled by autophagy. <i>Cell Death and Differentiation</i> , 2015 , 22, 445-56	12.7	70
40	Essential versus accessory aspects of cell death: recommendations of the NCCD 2015. <i>Cell Death and Differentiation</i> , 2015 , 22, 58-73	12.7	643
39	Basophils exhibit antibacterial activity through extracellular trap formation. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2015 , 70, 1184-8	9.3	45
38	NADPH oxidase-independent formation of extracellular DNA traps by basophils. <i>Journal of Immunology</i> , 2014 , 192, 5314-23	5.3	107
37	BID-dependent release of mitochondrial SMAC dampens XIAP-mediated immunity against Shigella. <i>EMBO Journal</i> , 2014 , 33, 2171-87	13	37
36	XIAP restricts TNF- and RIP3-dependent cell death and inflammasome activation. <i>Cell Reports</i> , 2014 , 7, 1796-808	10.6	172
35	TREM-1 deficiency can attenuate disease severity without affecting pathogen clearance. <i>PLoS Pathogens</i> , 2014 , 10, e1003900	7.6	83

34	The tumor suppressor gene DAPK2 is induced by the myeloid transcription factors PU.1 and C/EBP β during granulocytic differentiation but repressed by PML-RAR α in APL. <i>Journal of Leukocyte Biology</i> , 2014 , 95, 83-93	6.5	14
33	Intracellular localization of the BCL-2 family member BOK and functional implications. <i>Cell Death and Differentiation</i> , 2013 , 20, 785-99	12.7	93
32	Foxo-mediated Bim transcription is dispensable for the apoptosis of hematopoietic cells that is mediated by this BH3-only protein. <i>EMBO Reports</i> , 2013 , 14, 992-8	6.5	25
31	The Bcl-2 protein family member Bok binds to the coupling domain of inositol 1,4,5-trisphosphate receptors and protects them from proteolytic cleavage. <i>Journal of Biological Chemistry</i> , 2013 , 288, 25340-25349 ⁶⁶	5.4	66
30	In vitro differentiation of near-unlimited numbers of functional mouse basophils using conditional Hoxb8. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2013 , 68, 604-13	9.3	22
29	Consequences of the combined loss of BOK and BAK or BOK and BAX. <i>Cell Death and Disease</i> , 2013 , 4, e650	9.8	52
28	Fas death receptor signalling: roles of Bid and XIAP. <i>Cell Death and Differentiation</i> , 2012 , 19, 42-50	12.7	253
27	TRAIL enhances paracetamol-induced liver sinusoidal endothelial cell death in a Bim- and Bid-dependent manner. <i>Cell Death and Disease</i> , 2012 , 3, e447	9.8	16
26	Death receptor-induced apoptosis signalling - essential guardian against autoimmune disease. <i>Arthritis Research and Therapy</i> , 2012 , 14,	5.7	78
25	The ubiquitin ligase XIAP recruits LUBAC for NOD2 signaling in inflammation and innate immunity. <i>Molecular Cell</i> , 2012 , 46, 746-58	17.6	272
24	Novel insights into mechanisms of food allergy and allergic airway inflammation using experimental mouse models. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2012 , 67, 1483-90	9.3	17
23	BCL-2 family member BOK is widely expressed but its loss has only minimal impact in mice. <i>Cell Death and Differentiation</i> , 2012 , 19, 915-25	12.7	82
22	Role of TRAIL and the pro-apoptotic Bcl-2 homolog Bim in acetaminophen-induced liver damage. <i>Cell Death and Disease</i> , 2011 , 2, e171	9.8	29
21	A novel TNFR1-triggered apoptosis pathway mediated by class IA PI3Ks in neutrophils. <i>Blood</i> , 2011 , 117, 5953-62	2.2	64
20	MEK/ERK-mediated phosphorylation of Bim is required to ensure survival of T and B lymphocytes during mitogenic stimulation. <i>Journal of Immunology</i> , 2009 , 183, 261-9	5.3	66
19	Fatal hepatitis mediated by tumor necrosis factor TNF α requires caspase-8 and involves the BH3-only proteins Bid and Bim. <i>Immunity</i> , 2009 , 30, 56-66	32.3	108
18	XIAP discriminates between type I and type II FAS-induced apoptosis. <i>Nature</i> , 2009 , 460, 1035-9	50.4	344
17	Puma indirectly activates Bax to cause apoptosis in the absence of Bid or Bim. <i>Cell Death and Differentiation</i> , 2009 , 16, 555-63	12.7	64

16	Proapoptotic BH3-only protein Bid is essential for death receptor-induced apoptosis of pancreatic beta-cells. <i>Diabetes</i> , 2008 , 57, 1284-92	0.9	78
15	Switch from type II to I Fas/CD95 death signaling on in vitro culturing of primary hepatocytes. <i>Hepatology</i> , 2008 , 48, 1942-53	11.2	46
14	Loss of the BH3-only protein Bid does not rescue RelA-deficient embryos from TNF-R1-mediated fatal hepatocyte destruction. <i>Cell Death and Differentiation</i> , 2007 , 14, 637-9	12.7	5
13	The BH3-only protein bid is dispensable for DNA damage- and replicative stress-induced apoptosis or cell-cycle arrest. <i>Cell</i> , 2007 , 129, 423-33	56.2	170
12	Response: Does Bid Play a Role in the DNA Damage Response?. <i>Cell</i> , 2007 , 130, 10-11	56.2	10
11	Apoptosis initiated when BH3 ligands engage multiple Bcl-2 homologs, not Bax or Bak. <i>Science</i> , 2007 , 315, 856-9	33.3	937
10	The BH3-only protein Puma plays an essential role in cytokine deprivation induced apoptosis of mast cells. <i>Blood</i> , 2007 , 110, 3209-17	2.2	94
9	Apaf-1 and caspase-9 are required for cytokine withdrawal-induced apoptosis of mast cells but dispensable for their functional and clonogenic death. <i>Blood</i> , 2006 , 107, 1872-7	2.2	26
8	Chronic inflammation and pain inside the mandibular jaw and a 10-year forgotten amalgam filling in an alveolar cavity of an extracted molar tooth. <i>Ultrastructural Pathology</i> , 2005 , 29, 405-13	1.3	7
7	Conformational control of Bax localization and apoptotic activity by Pro168. <i>Journal of Cell Biology</i> , 2004 , 164, 1021-32	7.3	124
6	Bcl-w(edding) with mitochondria. <i>Trends in Cell Biology</i> , 2004 , 14, 8-12	18.3	35
5	Bcl-2 family members: integrators of survival and death signals in physiology and pathology [corrected]. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2004 , 1644, 95-105	4.9	113
4	Bcl-x(S) induces an NGF-inhibitable cytochrome c release. <i>Experimental Cell Research</i> , 2004 , 297, 392-403	4.2	9
3	Characterization of the signal that directs Bcl-x(L), but not Bcl-2, to the mitochondrial outer membrane. <i>Journal of Cell Biology</i> , 2003 , 160, 53-64	7.3	272
2	Inhibition of tumour cell growth by hyperforin, a novel anticancer drug from St. John's wort that acts by induction of apoptosis. <i>Oncogene</i> , 2002 , 21, 1242-50	9.2	202
1	Bcl-2 is a monomeric protein: prevention of homodimerization by structural constraints. <i>EMBO Journal</i> , 2000 , 19, 1534-44	13	34