Florian Kopp

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/2371857/florian-kopp-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

14
papers2,598
citations11
h-index17
g-index17
ext. papers3,540
ext. citations18.8
avg, IF6.4
L-index

| # | Paper | IF | Citations |
|----|--|---------------|-----------|
| 14 | Molecular functions and biological roles of long non-coding RNAs in human physiology and disease. <i>Journal of Gene Medicine</i> , 2019 , 21, e3104 | 3.5 | 24 |
| 13 | PUMILIO hyperactivity drives premature aging of -deficient mice. <i>ELife</i> , 2019 , 8, | 8.9 | 40 |
| 12 | PUMILIO, but not RBMX, binding is required for regulation of genomic stability by noncoding RNA. <i>ELife</i> , 2019 , 8, | 8.9 | 27 |
| 11 | Functional Classification and Experimental Dissection of Long Noncoding RNAs. Cell, 2018, 172, 393-40 | 0756.2 | 1584 |
| 10 | An Argonaute phosphorylation cycle promotes microRNA-mediated silencing. <i>Nature</i> , 2017 , 542, 197-2 | 03 0.4 | 140 |
| 9 | Noncoding RNA NORAD Regulates Genomic Stability by Sequestering PUMILIO Proteins. <i>Cell</i> , 2016 , 164, 69-80 | 56.2 | 496 |
| 8 | Salinomycin treatment reduces metastatic tumor burden by hampering cancer cell migration. <i>Molecular Cancer</i> , 2014 , 13, 16 | 42.1 | 48 |
| 7 | Sequential Salinomycin Treatment Results in Resistance Formation through Clonal Selection of Epithelial-Like Tumor Cells. <i>Translational Oncology</i> , 2014 , 7, 702-11 | 4.9 | 10 |
| 6 | The proto-oncogene KRAS is targeted by miR-200c. <i>Oncotarget</i> , 2014 , 5, 185-95 | 3.3 | 59 |
| 5 | Generation of a tumor- and tissue-specific episomal non-viral vector system. <i>BMC Biotechnology</i> , 2013 , 13, 49 | 3.5 | 11 |
| 4 | De-targeting by miR-143 decreases unwanted transgene expression in non-tumorigenic cells. <i>Gene Therapy</i> , 2013 , 20, 1104-9 | 4 | 9 |
| 3 | miR-200c sensitizes breast cancer cells to doxorubicin treatment by decreasing TrkB and Bmi1 expression. <i>PLoS ONE</i> , 2012 , 7, e50469 | 3.7 | 90 |
| 2 | Combinatorial treatment of mammospheres with trastuzumab and salinomycin efficiently targets HER2-positive cancer cells and cancer stem cells. <i>International Journal of Cancer</i> , 2012 , 131, 2808-19 | 7.5 | 59 |
| 1 | PLIMILIO but not RBMX binding is required for regulation of genomic stability by noncoding RNA NOR | RAD | 7 |