

# Anne Sofie Gram

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2373845/publications.pdf>

Version: 2024-02-01

11  
papers

222  
citations

1163117

8  
h-index

1281871

11  
g-index

11  
all docs

11  
docs citations

11  
times ranked

466  
citing authors

#	ARTICLE	IF	CITATIONS
1	Effect of exercise training on skeletal muscle protein expression in relation to insulin sensitivity: Perâ€protocol analysis of a randomized controlled trial (GOâ€ACTIVE). <i>Physiological Reports</i> , 2021, 9, e14850.	1.7	2
2	Structured exercise alters the gut microbiota in humans with overweight and obesityâ€A randomized controlled trial. <i>International Journal of Obesity</i> , 2020, 44, 125-135.	3.4	76
3	PPARG Pro12Ala Ala carriers exhibit greater improvements in peripheral insulin sensitivity in response to 12 weeks of aerobic exercise training. <i>Physiological Genomics</i> , 2019, 51, 254-260.	2.3	3
4	How does 6 months of active bike commuting or leisure-time exercise affect insulin sensitivity, cardiorespiratory fitness and intra-abdominal fat? A randomised controlled trial in individuals with overweight and obesity. <i>British Journal of Sports Medicine</i> , 2019, 53, 1183-1192.	6.7	28
5	Effects of Exercise Domain and Intensity on Sleep in Women and Men with Overweight and Obesity. <i>Journal of Obesity</i> , 2019, 2019, 1-12.	2.7	8
6	Effects of active commuting and leisure-time exercise on appetite in individuals with overweight and obesity. <i>Journal of Applied Physiology</i> , 2019, 126, 941-951.	2.5	16
7	Effects of 6 Months of Active Commuting and Leisure-Time Exercise on Fibrin Turnover in Sedentary Individuals with Overweight and Obesity: A Randomised Controlled Trial. <i>Journal of Obesity</i> , 2018, 2018, 1-10.	2.7	12
8	The GO-ACTIVE randomized controlled trial - An interdisciplinary study designed to investigate the health effects of active commuting and leisure time physical activity. <i>Contemporary Clinical Trials</i> , 2017, 53, 122-129.	1.8	22
9	Anti-inflammatory effects of active commuting and leisure time exercise in overweight and obese women and men: A randomized controlled trial. <i>Atherosclerosis</i> , 2017, 265, 318-324.	0.8	25
10	Three months of strictly controlled daily endurance exercise reduces thrombin generation and fibrinolytic risk markers in younger moderately overweight men. <i>European Journal of Applied Physiology</i> , 2015, 115, 1331-1338.	2.5	22
11	Compliance with physical exercise: Using a multidisciplinary approach within a dose-dependent exercise study of moderately overweight men. <i>Scandinavian Journal of Public Health</i> , 2014, 42, 38-44.	2.3	8